

Review

Recent Achievements in Microalgal Photobiological Hydrogen Production

Eleftherios Touloupakis ¹, Cecilia Faraloni ², Ana Margarita Silva Benavides ^{3,4} and Giuseppe Torzillo ^{2,3,*}¹ Istituto di Ricerca sugli Ecosistemi Terrestri, CNR, Via Madonna del Piano 10, 50019 Sesto Fiorentino, Italy; eleftherios.touloupakis@cnr.it² Istituto per la Bioeconomia, CNR, Via Madonna del Piano 10, 50019 Sesto Fiorentino, Italy; cecilia.faraloni@ibe.cnr.it³ Centro de Investigacion en Ciencias Del Mar y Limnología, Universidad de Costa Rica, San José 2060, Costa Rica; ana.silva@ucr.ac.cr⁴ Escuela de Biología, Universidad de Costa Rica, San José 2060, Costa Rica

* Correspondence: giuseppe.torzillo@cnr.it

Abstract: It is well known that over the last 60 years the trend of long-lived greenhouse gas emissions have shown a strong acceleration. There is an increasing concern and a mounting opposition by public opinion to continue with the use of fossil energy. Western countries are presently involved in a so-called energy transition with the objective of abandoning fossil energy for renewable sources. In this connection, hydrogen can play a central role. One of the sustainable ways to produce hydrogen is the use of microalgae which possess two important natural catalysts: photosystem II and hydrogenase, used to split water and to combine protons and electrons to generate gaseous hydrogen, respectively. For about 20 years of study on photobiological hydrogen production, our scientific hopes were based on the application of the sulfur protocol, which indisputably represented a very important advancement in the field of hydrogen production biotechnology. However, as reported in this review, there is increasing evidence that this strategy is not economically viable. Therefore, a change of paradigm for the photobiological production of hydrogen based on microalgae seems mandatory. This review points out that an increasing number of microalgal strains other than *Chlamydomonas reinhardtii* are being tested and are able to produce sustainable amount of hydrogen without nutrient starvation and to fulfill this goal including the application of co-cultures.

Keywords: biohydrogen; microalgae; *Chlamydomonas reinhardtii*; *Chlorella* sp.; photobioreactors; light conversion efficiency



Citation: Touloupakis, E.; Faraloni, C.; Silva Benavides, A.M.; Torzillo, G. Recent Achievements in Microalgal Photobiological Hydrogen Production. *Energies* **2021**, *14*, 7170. <https://doi.org/10.3390/en14217170>

Academic Editor: Wei-Hsin Chen

Received: 6 October 2021

Accepted: 27 October 2021

Published: 1 November 2021

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Microalgae are capable of converting light energy into chemical energy. Biofuels such as biodiesel, biohydrogen, and bioethanol can be derived from microalgae [1]. Photosynthesis in microalgae is coupled to the splitting of water and the evolution of oxygen (O₂). This process is catalyzed by the membrane-bound multi-protein complex photosystem II (PSII) [2].

It has been known since 1942, when Gaffron and co-workers noticed that under anaerobic conditions *Scenedesmus obliquus* cells can transiently produce hydrogen (H₂) upon illumination when deprived of oxygen [3]. In microalgae, hydrogenase enzyme catalyzes H₂ production in a light-dependent process [4]. Upon illumination, after a dark incubation period, due to light-driven electron transport from ferredoxin to hydrogenase, H₂ production is observed. H₂ production in microalgae can be divided into direct or indirect processes [5]. A direct process occurs when electrons (e[−]) from water splitting are transferred via PSII and ferredoxin to hydrogenase. An indirect process occurs when e[−] are derived from the metabolism of carbohydrates, previously accumulated during the (light) aerobic phase, and then utilized for H₂ production via both a photo-fermentation

process involving photosystem I (PSI) and in a process in the dark, involving the enzyme pyruvate:ferredoxin oxidoreductase (PFR). In *Chlamydomonas reinhardtii* (hereafter *C. reinhardtii*), PFR enzyme catalyzes the reduction of ferredoxin (Fdx) and the transfer of e^- to hydrogenase in a similar pathway to that utilized by bacteria (Figure 1) [6]. In *C. reinhardtii*, up to 92% of the final H_2 output comes from the direct photolysis coupled to the water oxidation operated by PSII [7]. Contribution of dark fermentation to the overall H_2 output is considered negligible (about 4%) in *C. reinhardtii*, but it can be significant in other microalgae such as *Chlorella*, as recently shown [8]. Microalgal hydrogenase enzymes are inactivated by the presence of molecular oxygen, and their expression is induced under anaerobic conditions.




Figure 1. Metabolic hydrogen production pathways used by *Chlamydomonas reinhardtii*. FDX: ferredoxin; H_2 ase: hydrogenase; NPQR: NADPH–plastoquinone oxidoreductase; PFR: pyruvate:ferredoxin oxidoreductase; PSI: photosystem I; PSII: photosystem II.

In recent years, energy-related H_2 demands have prompted scientists to develop methods that greatly enhance the H_2 -evolving ability of microalgae. The most promising approach has been the so-called “two-stage process” of photosynthesis (stage 1) and H_2 production (stage 2) [9]. In this process, there is a separation of the reactions of oxygen and hydrogen production. This bypasses the sensitivity of the hydrogenase enzyme to oxygen. Under such conditions, it was possible to produce significant volumes of H_2 by *C. reinhardtii* in a sustained process.

Several microalgae species have been studied for H_2 production, especially *C. reinhardtii*, *Chlorella vulgaris*, and *Chlorella pyrenoidosa* [10–13]. Among them, *C. reinhardtii* is a model microorganism widely recognized as an H_2 producer, presenting a hydrogenase with an enzymatic activity 10 to 100 times higher than other species [14]. H_2 production requires many optimization steps in order to reach a sustainable process [8,14–17]. Some of these parameters include choosing a proper microalgae strain and selecting appropriate culture conditions (growth media, light, pH, temperature, chlorophyll concentration) and proper photobioreactor (PBR) designs [18–20].

Many works have reported improved H_2 production in many microalgal strains by using sulfur, phosphorus, or nitrogen-depleted media [12,21–23]. In such culture conditions, microalgae sustain H_2 production only for some days since macro/micro-nutrient depletion in the culture compromises cell viability. This is the major drawback in microalgal H_2 production processes carried out by nutrient deprivation. Microalgae-based H_2 production requires anaerobic conditions due to the sensitivity of hydrogenase to O_2 [24]. O_2 sensitivity of hydrogenase is a major issue for H_2 production; therefore, there are many studies on oxygen suppression in order to improve H_2 production yield. Genetic and metabolic engineering of microalgae [25,26], nutrient stresses [27,28], light conditions optimization [29], and elimination of competing pathways for electrons [30] are examples of strategies used to improve H_2 evolution in microalgae.

This review provides an overview of the most relevant achievements in the photobiological production of H₂ by microalgae, and proposes a change of paradigm for the future research in the field.

2. Genetic Modification

Krishna et al. reported that sustained H₂ production is achieved by altering the ratio between PSI and PSII [31]. In this work, a *C. reinhardtii* C3 mutant with a modified PSI/PSII ratio (0.33) produced H₂ with a rate of 3 mL H₂/L/d for 42 days. Chen et al. identified a *C. reinhardtii* mutant strain hpm91 lacking proton gradient regulation 5, with 30-fold H₂ production yield compared to wild type (WT) [32]. Characterization of the hpm91 strain revealed an increased reactive-oxygen-species-scavenging capacity. This translates into an enhanced stability of PSII complex and increased H₂ production yield. Steinbeck et al. investigated the capacity of *C. reinhardtii* *pgr5* and *pgr5 pgrl1* double mutant to produce H₂ [33]. The *pgr* mutants showed four times higher maximal enhanced H₂ production rate (7 mL/L/h) than the WT. Pinto et al. studied a *Chlamydomonas* mutant with reduced rubisco levels, activity, and stability [34]. This mutant was used to reduce carbon fixation by Calvin cycle activity, which is the main competitor for the reducing power required by the hydrogenase. In this work, the rubisco mutant presented 15 times higher H₂ production than the WT. Eilenberg et al. studied the in vivo H₂ production efficiency of a *C. reinhardtii* strain Fd-HydA containing ferredoxin fused to HydA. H₂ production rate was 4.5 times higher than that of the native HydA in vivo [35]. Torzillo et al. showed that the in vivo H₂ production of the *C. reinhardtii* mutant strain L159I-N230Y was up to 5-fold higher (16 nmoles H₂/μg_{chl}/h) than that of *C. reinhardtii* CC 124 [36,37]. Batyrova et al. developed a genetically modified *C. reinhardtii* strain that activates photosynthesis in a cyclical manner. In this strain, the low O₂ production benefits H₂ production [38]. In comparison with the WT, this genetically modified strain presented higher H₂ production levels. Kosourov et al. showed that a truncated light antenna *C. reinhardtii* mutant could produce six times more H₂ compared to the WT strain [39]. Xu et al. introduced a catalase gene from *Synechococcus elongatus* PCC7942 and an *Escherichia coli* pyruvate oxidase gene, both driven by a HSP70A/RBCS2 promoter, into the chloroplast of *C. reinhardtii* [40]. Under low light, these microalgal cells consumed more O₂ than WT, resulting in a lower O₂ content and increased H₂ production [40]. Kruse et al. used the *Chlamydomonas* strain Stm6, which has a modified respiratory metabolism and large starch reserves compared with the WT [41]. *Chlamydomonas* strain Stm6 presented 5–13 times increased H₂ production rate (540 mL H₂/L_{culture}) compared to the WT [41]. Later, Volgusheva et al. obtained similar results by using the *Chlamydomonas* Stm6 mutant [42]. They attained an anaerobic condition much faster in the Stm6 strain than in the WT. This was a result of the higher respiration rate and lower initial O₂ production rate. H₂ production was four times higher in the Stm6 strain compared to the WT. Oey and co-workers reported the knock-down of the LHCMB 1, 2, and 3 proteins in the *C. reinhardtii* strain Stm6Glc4 [43]. The produced *C. reinhardtii* mutant exhibited increased light-to-H₂ and biomass conversion efficiencies of 180% and 165%, respectively. Wu et al. introduced a leghemoglobin gene (*lba*) into chloroplasts of *C. reinhardtii*. The genetically modified *Chlamydomonas* with *lba* gene consumed O₂ faster than WT, thus improving H₂ production [44]. Noone et al. introduced the clostridial hydrogenase gene into *C. reinhardtii* that contains insertionally inactivated hydrogenase genes. The presence of the more O₂-tolerant clostridial hydrogenase led to more sustained H₂ production [45].

Nowadays, the primary current challenge of such a process is the development of an oxygen-resistant hydrogenase. However, other bottlenecks may also be of significant importance, such as the oxygen sensitivity of hydrogenases. In this case, a number of other scientific and engineering issues are very likely to arise. They may include: (1) maximizing photosynthetic light-conversion efficiency (LCE); finding the proper redox potential balance in the organism to facilitate H₂ production; (2) preventing the effect of the buildup of high relative H₂ partial pressure restricting the process by feedback

inhibition; (3) addressing inefficient metabolic processes such as unneeded ATP generation during H₂ production in microalgae; (4) examining issues associated with the generation of destructive, active-oxygen species; and (5) minimizing the production of alternative, carbon-containing products that drain usable reducing power from the system. Recently, an increased H₂ output was attained by bioengineering photosynthesis [46].

In the following paragraphs, some of the most recent strategies used for sustained photobiological H₂ production by microalgae are summarized.

3. O₂ Removal

The use of inert gas (such as N₂ or Ar) is another type of strategy to remove the O₂ in microalgal cultures [47,48]. Alternatively, O₂ scavengers can be employed to remove the O₂ in order to induce anaerobiosis in the culture. Paramesh and Chandrasekhar screened three O₂ scavengers individually in order to improve H₂ production in *Chlorococcum minutum* [49]. In the presence of all three O₂ scavengers, efficient H₂ generation was found. They found that sodium sulfite was the best one for enhancement of H₂ production. Nagy et al. showed that the simultaneous addition of glucose, glucose oxidase, and ascorbate to the *C. reinhardtii* culture resulted in reduced O₂ content in the headspace and tenfold-increased H₂ production [30]. Su et al. created an O₂-consuming sandwich-like layer by using tannic acid, polydopamine, and laccase, in order to generate anaerobiosis around the *Chlorella pyrenoidosa* cells [50]. This layer enabled the encapsulated cell to switch from O₂ production to H₂ production. Márquez-Reyes et al. found that the chemical reducing agent cysteine induced anaerobic H₂ production in cultures of *Chlamydomonas gloeopara* and *Scenedesmus obliquus* cultures [51]. In the presence of cysteine, H₂ production was 5 times higher compared to the sulfur-starvation protocol. Chen and coworkers found that *C. reinhardtii* produce H₂ at a rate of 0.44 µmol H₂/h/mg_{chl} per month by using a chemoenzymatic cascade system (CEC). The CEC system contained four components: glucose oxidase, catalase, glucose, and Mg(OH)₂. In this CEC, they combined O₂ consumption, cell aggregation, and pH maintenance to activate hydrogenase [52]. Nagy et al. showed that the application of an iron-based O₂ absorbent (O2O_{TM}) in *C. reinhardtii* cultures, in which the activation of the Calvin–Benson–Bassham cycle in the light was prevented, presented a H₂ production yield of 2.58 mL/L/h, to which corresponded a mean LCE (light to H₂) of 0.27% [30].

4. Co-Cultures

Another approach to create an anaerobic environment is the addition of living aerobic bacteria to the microalgae cultures (co-culture) [53]. Many works have proven the possibility of increasing H₂ production by co-culturing microalgae and bacteria [28,54–57]. The main advantage of co-culturing microalgae with heterotrophic bacteria is the efficient removal of the O₂ from the growth media. Simultaneously, the CO₂ released during bacterial fermentation of an organic substrate can support microalgae growth. Moreover, many metabolites can be exchanged between these microorganisms, such as carbon, nitrogen, phosphorous, and sulfur sources, and vitamins [58,59]. The presence of bacteria inside the microalgal culture enhances starch accumulation [60]. Different *Chlamydomonas* WT co-cultures incubated in sulfur depleted TAP medium employing *Pseudomonas* sp. or *Bradyrhizobium japonicum* have achieved high H₂ production rates (165–170 mL H₂/L) [54,60]. Fakhimi et al. evaluated H₂ production by *C. reinhardtii* in co-cultures with different bacteria strains [28]. They found that co-culturing *Pseudomonas* spp. with *Chlamydomonas* significantly improved microalgal H₂ production. Interestingly, the integration of the photobiological and the fermentative H₂ production in *Chlamydomonas* and *Escherichia coli* co-cultures resulted in H₂ production 60% higher than the sum of the respective monocultures [28]. *Chlamydomonas* co-cultures with *Pseudomonas* sp. and *Bradyrhizobium japonicum* (not H₂-producing bacteria) in sulfur-depleted TAP medium improved H₂ production by 22.7 times and 32.3 times compared to the pure microalgal cultures, respectively [57]. Furthermore, the production of H₂ by *C. reinhardtii* in nutrient-replete cultures is strongly limited by the O₂ release, unless it is performed under very low

light irradiance (lower than $20 \mu\text{mol photons/m}^2/\text{s}$), but it may become feasible under higher light irradiance by using different consortia, which allow the maintenance of anaerobiosis conditions, thus creating an opportunity to use full medium and much higher light irradiance, enhancing the H_2 output. Finally, it must be pointed out that large-scale production of H_2 with *Chlamydomonas* and other microalgae will be necessarily carried out not with axenic cultures, but rather a microalgae–bacteria consortium, therefore understanding the complex interplay between microalgae H_2 producers and bacteria is important for the economic exploitation of an industrial H_2 production process.

5. Immobilization

Microalgae immobilization can increase the H_2 production yield [61]. The main reason for their higher performance is that such experiments are usually carried out by using a much higher chlorophyll concentration compared to liquid cultures. The process of switching between oxygenic photosynthesis (aerobiosis) and H_2 production (anaerobiosis) can be facilitated by using cell immobilization systems [62]. One of the most used materials for microalgae encapsulation is calcium alginate [63]. Immobilization in calcium alginate matrix allows high cell density and protection from mechanical stress and contamination and is easy to scale-up [64]. Immobilization of microalgal cells could increase their LCE [62]. Using *Calothrix*, *Anabaena*, and *Chlamydomonas* cells immobilized on thin calcium alginate films gave an LCE of 2.5% of the photosynthetically active radiation [48,65]. *C. reinhardtii* immobilized on calcium alginate films in nutrient-depleted cultures (-P,-S) presented a H_2 production rate of $12.5 \mu\text{mol/mgChl/h}$ [66]. Ruiz-Marin et al. proposed immobilization of *Chlorella vulgaris* and *Scenedesmus obliquus* cells in calcium alginate for the production of H_2 [11]. These microalgae were grown in urban wastewater under sulfur starvation and blue or purple light conditions. The maximum H_2 production obtained under red light was $204.8 \text{ mL H}_2/\text{L/d}$ for *Scenedesmus obliquus* and $39.1 \text{ mL H}_2/\text{L/d}$ for *Chlorella vulgaris* [11]. Maswanna et al. studied H_2 production by *Tetraspora* sp. CU2551 cells immobilized in a 4% w/v calcium alginate matrix in their recent work. They obtained a maximum H_2 production rate of $182 \pm 20 \text{ nmol/mg of cell dry weight/h}$ [67].

Our group recently tested the capability of immobilized *Chlorella vulgaris* (BEIJ G-120 strain) cells in a calcium alginate (3%w/v) gel matrix to produce H_2 in a direct light-driven process under continuous illumination. Calcium alginate beads were stable, showing minimal cell leakage, and they measured $4.69 \pm 0.02 \text{ mm}$ in diameter and $54.01 \pm 0.03 \mu\text{L}$ in volume, carrying $145.5 \pm 8.9 \mu\text{g}$ of microalgal cells (biomass dry weight/bead) (Figure 2). Immobilized cells retained their viability for more than 30 days (Figure 3A). Immobilized *Chlorella* cells were capable of generating H_2 without nutrient deprivation with a maximum rate of 162 mL/L (Figure 3B). Anaerobiosis was maintained by the presence of glucose and the high respiration rate of the strain.




Figure 2. *Chlorella vulgaris* (BEIJ G-120 strain) immobilized in calcium alginate beads. (A) Calcium-alginate bead; (B) *Chlorella vulgaris* cells inside the calcium-alginate beads.




Figure 3. (A) F_v/F_m of immobilized *Chlorella vulgaris* cells as a function of time; (B) H_2 production of the immobilized *Chlorella vulgaris* cells as a function of time.

6. Hydrogen Production without Nutrient Starvation

The H_2 production protocol by Melis and coworkers based on sulfur starvation greatly improved light-driven, algal H_2 production, and particularly the possibility for researchers to study the process [68]. However, in the recent years it has become clear that it is not adequate for an industrial development of the process since it requires one to eliminate sulfur residues. Moreover, the severe reduction of PSII activity caused by the sulfur deprivation greatly reduces the H_2 production and thus the viability of the process. Awareness of these limits has prompted several workers to eliminate the sulfur deprivation phase by selecting strains with high respiration-to-photosynthesis ratios.

Liu et al. presented a work on H_2 production of *Chlorella pyrenoidosa* using $NaHCO_3$ as a carbon source and $N'-(3,4\text{-Dichlorophenyl})\text{-}N,N\text{-dimethylurea}$ (DCMU) [69]. In this work, *Chlorella pyrenoidosa* cells showed an overall H_2 production of 93.86 mL/L. In a recent work, Li et al. constructed a transgenic *C. reinhardtii* strain (amiRNA-D1) with a heat-inducible expression system targeting D1 gene (*psbA*). After a heat-shock, the transgenic *C. reinhardtii* strain presented a 73% decrease of *psbA* gene expression and a 60% increase of H_2 content compared to the WT strain [70]. Ben-Zvi et al. explored the in vivo H_2 production of HydA–SOD fusion phenotype in *C. reinhardtii* and found that expression of an active hydrogenase superoxide dismutase fusion protein resulted in sustained H_2 production with a rate of 20 mL H_2 /L/d for 8 days [71]. Hwang et al. showed that the over-expression of the hydrogenase gene in *Chlorella vulgaris* resulted in H_2 production under aerobic conditions with continuous illumination using CO_2 as the sole source of carbon [72]. Under 5% O_2 and 10% CO_2 , *Chlorella vulgaris* strains YSL01 and YSL16 produced 1.9 mL H_2 /h and 1.2 mL H_2 /h in 3 and 4 days, respectively. In another of their works, this group studied and compared the photosynthetic activities of *C. reinhardtii* and *Chlorella sorokiniana* with different acetate/ Cl^- ratios [73]. They found that maintaining acetate/ Cl^- ratios greater than 60–100 led to continuous O_2 depletion. Using fermenter effluents, at an acetate/ Cl^- ratio of 150, *Chlorella sorokiniana* and *C. reinhardtii* presented an H_2 production rate of 0.25–0.33 mmol/L/min and 0.20–0.38 mmol/L/min, respectively. Kosourov et al. demonstrated sustained H_2 production by *C. reinhardtii* by shifting the culture light conditions from continuous illumination to a set of light pulses interrupted by longer dark phases [29]. In a recent work, Sirawattamongkol et al. demonstrated that *Chlorella* sp. strain KLS Sc59 was able to produce up to 750 mL H_2 /L in the presence of reducing agents such as ethanol and dithionite [74]. H_2 production rates in various microalgae strains are summarized in Table 1.


Table 1. Comparison of H₂ production rates in various microalgae strains.

Microalgal Species	Growth Mode	H ₂ Production	References
<i>C. reinhardtii</i>	TAP	25 µmol/mg _{Chl} /h	[29]
<i>Chlorella</i> sp. AARL G014	TAP-S	0.49 mmol/mg _{Chl} /h	[75]
<i>C. reinhardtii</i> CC-503	TAP co-culture	255 mmol/mg _{Chl}	[55]
<i>Chlorella vulgaris</i> strains YSL01	BBM-EDTANa ₂	1.9 mL/L	[72]
<i>Chlorella lewinii</i> KU201	TAP-S	13.03 mL/L	[18]
<i>Chlorella</i> sp. IOAC707S	TAP-NaCl	38.00 mL/L	[38]
<i>Chlorella sorokiniana</i> KU204	TAP-P	69.00 mL/L	[18]
<i>Chlorella protothecoides</i>	TAP-NS	82.50 mL/L	[76]
<i>Chlorella sorokiniana</i> KU204	TAP-S	89.64 mL/L	[18]
<i>Chlorella pyrenoidosa</i>	TCP + DCMU	93.86 mL/L	[69]
<i>C. reinhardtii</i> Stm6	TAP-S	540 mL/L	[41]
<i>C. reinhardtii</i> C3	TAP	3.0 mL/L/d	[31]
<i>C. reinhardtii</i> (HS-14)	TAP	20 mL/L/d	[71]
Immobilized <i>Chlorella vulgaris</i>	Artificial wastewater-S	39.1 mL/L/d	[11]
<i>Chlorella vulgaris</i> MACC360	TAP co-culture	56.0 mL/L/d	[77]
Immobilized <i>Scenedesmus obliquus</i>	Artificial wastewater-S	204.8 mL/L/d	[11]
<i>Chlorella salina</i> Mt	TAP-S	0.5 mL/L/h	[78]
<i>C. reinhardtii</i> CC124	TAP-S	0.6 mL/L/h	[79]
<i>C. reinhardtii</i> CC-124	TAP-S	3.3 mL/L/h	[80]
<i>C. reinhardtii</i> pgr5/pgr11	TAP-S	7.0 mL/L/h	[33]
<i>C. reinhardtii</i> L159I-N230Y	TAP-S	11.1 mL/L/h	[37]
<i>Chlorella vulgaris</i> BEIJ (G-120)	HM + glucose	5.0 mL/L/h	[8]
Immobilized <i>Chlorella vulgaris</i> NIER-10003	MA-S + glucose	238 mL/L/h	[81]
<i>Chlorella sorokiniana</i>	150 of acetate/Cl ⁻ ratio	0.33 mmol/L/min	[73]
<i>C. reinhardtii</i>	150 of acetate/Cl ⁻ ratio	0.38 mmol/L/min	[73]
<i>C. reinhardtii</i> (ΨH1)	TAP	3.6 mL/L/h	[46]
<i>C. reinhardtii</i>	HSM + O ₂ absorbent	2.58 mL/L/h	[30]

In our recent work, we reported H₂ production by *Chlorella vulgaris* (strain BEIJ G-120) without the use of nutrient deprivation [8]. This *Chlorella* strain presents two main properties: high respiration rate and high light compensation point. By exploiting these two properties, it was possible to efficiently consume the photosynthetically produced O₂, thus maintaining anaerobiosis, even under light conditions. In this work, *Chlorella* cells presented a maximum H₂ production rate of 12 mL/L/h and an average rate of 4.98 mL/L/h. The strain was capable of producing H₂ in the dark as well, by fermentation of glucose. The excessive accumulation of byproducts of the fermentation (e.g., acetate, formate, lactate, ethanol) may inhibit H₂ production. However, the possibility of also producing H₂ in the dark by microalgae is desirable for the development of the process under natural light/dark cycle. On the other hand, some of the byproducts of dark fermentation, such as acetate, can be used as substrate for mixotrophic grown during the following light phase.

7. Theoretical Limit for Biological Hydrogen Production

Only a small fraction of the total solar light radiation (>1,100,000 EJ per year) can potentially be transformed into H₂ energy using the process of photosynthesis, according to the following general equation:



Step 1 indicates the total incident radiation received at the surface of the culture (100%) (Figure 4). It follows that:

1. Approximately 10% is lost by reflection and scattering (90% of initial remaining).
2. Approximately 55% of radiation is not available to drive photosynthesis since it falls outside of the photosynthetically active radiation (400–700 nm) and thus is not utilized

- by photosynthetic pigments. As a result, the total amount of available light drops to 41%.
- About 20.4% of the radiation is lost as heat [82].
 - Assuming as quantum requirement that 8 photons are required to produce 2 mol of H_2 , and considering that 1 mol of H_2 is 286 KJ, and the mean energy for charge separation at PSII and PSI is 173.5 KJ/mol, it follows that the efficiency of the process will be the following: $(286 \text{ KJ/mol} \times 2) / (173.5 \text{ KJ/mol} \times 8) \times 100 = 41.2\%$, with a corresponding loss of energy of 59%. Consequently, the theoretical LCE for H_2 production, attainable by direct biophotolysis is about 13.4% of incident solar light [83].
 - With a LCE of about 10%, assuming that approximately 20% of the energy can be lost for cell maintenance, it might be possible to produce about 600,000 $m^3/ha/y$ of H_2 in sunny areas.




Figure 4. The energy losses of the incident solar light during the different steps of the photobiological H_2 production process.

8. Chlorophyll Fluorescence Measurements as a Tool for Monitoring Changes of Photochemical Efficiency during the Hydrogen Production Process

Chlorophyll fluorescence is a fast and non-invasive tool for monitoring residual photosynthetic activity during the H_2 production process through changes in the maximum quantum yield of PSII (F_v/F_m) and the effective quantum yield ($\Delta F/F'_m$) [38]. In particular, it has been observed that in *C. reinhardtii*, at the occurrence of anaerobiosis, the value of $\Delta F/F'_m$ rapidly declined, and this drop could be ascribed to the state 1 to state 2 transition, controlled by the redox state of plasto-quinone (PQ)-pool [36,84–86]. This mechanism regulates the migration of the light-harvesting complex (LHC) from PSII (state 1) to PSI (state 2), and it is induced under high level of PQ-pool reduction and excess of light energy. The start of the H_2 production induces a partial oxidation of the photosynthetic electron chains, comprising PQ-pool, with a partial recovery of $\Delta F/F'_m$. In *C. reinhardtii*, migration of LHC can involve up to 80% of the total LHC. The redox potential of the cells represents another important parameter related to the cell physiology under anaerobiosis, as it is the result of a balance between starch degradation, the capacity of PSII to perform photosynthesis, and the ability of cell to dissipate electrons from PQ-pool. Indeed, after establishing anaerobiosis, the value of the redox potential changes from a positive initial value to a very low value (about -550mV in *C. reinhardtii*). The changes of the values of the redox potential lag behind the changes in the yield and are less rapid than the chlorophyll fluorescence changes. In Figure 5, an example of the typical kinetics of chlorophyll fluorescence yield and redox potential in the different phases of the H_2 production process is reported.




Figure 5. *C. reinhardtii* kinetics of chlorophyll fluorescence and redox potential during induction of H_2 production under sulfur deprivation. (A) Time courses in dissolved oxygen (empty circle) and output of hydrogen (H_2) (filled square). (B) The time courses in the effective quantum yield of PSII ($\Delta F/F'_m$) (filled circle) and the redox potential (Eh) (empty diamond) in *C. reinhardtii* under sulfur deprivation with 70 $\mu\text{mol photons/m}^2/\text{s}$, supplied on both sides of the reactor.

Other important information on the changes of the photosynthetic efficiency can be provided by the chlorophyll fluorescence rise kinetics (OJIP curve), strictly reflecting the progressive reduction of the photosynthetic electron transport chain [60,87], which can indicate and quantify the reduction of electrons transport for each step [88]. The most evident change occurs at the J-step level, indicating the reduced transfer of electrons further than Q_A , measured by V_J parameter, and thus, an accumulation of reduced Q_A^- [88]. An example of the changes of the shape of the OJIP curve during the occurrence of anaerobiosis in *C. reinhardtii* is reported in Figure 6.




Figure 6. Effect of sulfur deprivation on the chlorophyll a fluorescence transient in *C. reinhardtii* cultures. Start (dark squares); 40 h of sulfur deprivation (blue triangles); 110 h of sulfur deprivation (red diamonds). Relative variable fluorescence ($V_t = (F_t - F_0)/(F_m - F_0)$) [89].

In *Chlorella sorokiniana*, nutrient starvation conditions reduce photosynthetic activity and induce anaerobiosis and H_2 production, indicated by the decrease of both F_v/F_m and $\Delta F/F'_m$. Similarly, to what was observed in *C. reinhardtii*, the maintenance of a residual PSII activity provides electron to hydrogenase enzyme [90]. The same behavior was observed in *Chlorella vulgaris* without nutrient starvation with a strain able to reach anaerobiosis in complete medium, showing a decline of F_v/F_m and $\Delta F/F'_m$ within 24 h [8].

In conclusion, the use of fluorescence measurements to monitor changes in photosynthetic activity can help us to better understand the physiological status of microalgae during the H_2 production process, making it easier to interfere in the cell metabolism or enhance the production process. Moreover, the application of chlorophyll fluorescence helps in selecting strains more resistant to the stress imposed by anaerobic conditions, and with higher potential H_2 output.

9. Photobiological Hydrogen Production in Outdoor Photobioreactors

Until now, H_2 production experiments using *C. reinhardtii* have been carried out mostly under laboratory conditions. Mean LCE in sulfur-deprived laboratory cultures grown in well-mixed PBR has hardly surpassed 1% (light to H_2). The necessity to downregulate the PSII activity to the level of the respiration is considered the main reason for such a low efficiency. As a matter of fact, the LCE strongly increased when it was possible to use microalgal strains with high respiration-to-photosynthesis ratio. This was the case of the *Chlorella* strain G-120, which averaged 3.2%, over the 8-day period [8].

The utilization of solar energy is mandatory for the economical scale-up of the H_2 production process. However, under solar light, the light energy received by microalgae cells exceeds their ability of light conversion into valuable biomass. This leads to either energy dissipation as heat or to photodamage and cell death, which strongly reduce the LCE. To reduce the “saturation effect”, a number of PBR designs have been proposed [91]. Torzillo and coworkers reported H_2 production of about 21% of that attained under laboratory in an outdoor 50 L tubular PBR using *C. reinhardtii* under sulfur deprivation (Figure 7) [92]. The PBR consisted of ten glass tubes (2.0 m length and 4.85 cm internal diameter) placed horizontally and connected by polyvinylchloride U-bends (Figure 7). The PBR was placed in a stainless-steel container with temperature-controlled water. A polyvinyl chloride pump allowed the culture to circulate.




Figure 7. The tubular photobioreactor (working volume 50 L) used for the outdoor H_2 production experiments. The photobioreactor was equipped with probes for measurement and control of pH and temperature. Culture speed can be adjusted to reach the desired turbulence.

The low performance of the culture was explained by the rapid PSII inactivation by the high light irradiation, which during the experiments reached as much as $1850 \mu\text{mol photons/m}^2/\text{s}$ in the middle of day. In order to avoid the problem of light

saturation, Giannelli and Torzillo, 2012 [79] proposed a 110 L PBR in which the culture tubes were immersed in water with light-scattering silica nanoparticles. The PBR contained 64 glass tubes (length 2.0 m, internal diameter 27.5 mm) arranged on an 8×8 square pitch cell and connected by polyvinylchloride U-bends. The PBR was immersed in a rectangular parallelepiped tank made of isotactic polypropylene, except for the opposite square faces, which were made of transparent Plexiglas. The culture was circulated with a peristaltic pump. The light scattering promoted by nanoparticles permitted a homogeneous distribution of light on the surface of the PBR (Figure 8). Solar light was collected by two sun-tracking mirrors, which delivered light to the opposite faces of the reactors through two light ducts (Figure 8c).




Figure 8. The 110 L photobioreactor utilized for the hydrogen production experiments. (a) General view of the photobioreactor; (b) frontal view of the photobioreactor; (c) sun-tracking mirrors collecting light; (d) light ducts delivering light to the photobioreactor from the opposite sides; (e) nocturnal view of the photobioreactor with two light bulbs (1000 W each) placed in the light duct to provide illumination to the culture during the night.

The total amount of H_2 collected reached 3.5 L, which was almost 2-fold higher than that obtained with the 50 L tubular PBR per unit of volume of reactor. Furthermore, in the scaled-up PBR, the LCE of the process increased from 0.055% in the 50 L horizontal PBR to 0.213% in the 110 L PBR, which was much closer to that attained in the laboratory with sulfur-deprived *C. reinhardtii* cultures. This still-low efficiency was the result of a number of

factors: (i) the necessity to carry out the process according to the two-step protocol (sulfur starvation), which strongly reduces the contribution of PSII; (ii) the need to attain sulfur starvation by culture dilution, which normally yields lower H₂ output; and (iii) the longer mixing time of tubular PBRs, which delays reaching H₂-saturation levels in the cultures.

10. Optimal Photobioreactor Design for the Hydrogen Production

The experience acquired from experiments with different indoor and outdoor PBRs has provided some guidelines useful for the optimal design of future PBRs for H₂ production [93,94]. An in-depth discussion on the influence of the PBR design on the H₂ output can be found in a recent book on the subject of H₂ production [94].

The development of an economically feasible PBR system is the most important factor for successful H₂ production. Closed PBRs (such as flat-panel and tubular) are mandatory for H₂ production. Tubular PBRs contain one or more glass tubes arranged in various configurations and orientations. Flat-panel PBRs consist of one or more transparent panels containing the culture, which is circulated between the panels by a pump. Flat-panel PBRs probably meet most of the above-mentioned requirements for H₂ production. They present high area-to-volume ratio and good biomass productivity, and shorter mixing time compared to tubular PBRs, which reduces the risk of H₂ oversaturation in the reactor. A drawback is the difficulty of controlling the temperature, and the high power consumption for mixing. Although the setup costs of closed PBRs are high, they provide several advantages, such as optimal growth, minimizing the risk of culture contamination, and reduced water and CO₂ consumption.

The International Energy Agency established the commercial cost target for H₂ production at 0.3 USD/kg. According to James et al., with 10% LCE, the cost could be 2.99 USD per gallon of gasoline equivalent [95]. The authors estimated that by using robust microalgae strains presenting 1.5% LCE, the cost of H₂ would be 8.44 USD per gallon of gasoline equivalent. Greater costs for PBR construction are materials, manufacturing, and personnel costs. H₂ production prices should be more promising than market prices. Show et al. showed that the costs of fabrication materials and chemical nutrients are the main expenses (84% of the total cost) for PBR development [96]. Recycling metabolic products of PBRs (such as organic acids) and/or considering potentially cheaper nutrient sources are possible ways to reduce the chemical cost of nutrients necessary for microalgal growth [97]. Finally, production cost based on direct bio-photolysis, were estimated to be, 18.45 \$/kgH₂, for the Netherlands, which is expected to drop significantly in the future (potential cost of 3.10 \$/kgH₂) [98].

11. Concluding Remarks

This review provides evidence that photobiological H₂ production by microalgae and cyanobacteria might be a viable option. The discovery of the sulfur starvation method has allowed maintenance of H₂ production for several days, and thus, it has represented an opportunity to study the process beyond scientific curiosity.

Nowadays, however, a substantial change of paradigm in photobiological H₂ production is necessary. The opportunity to improve the economic feasibility of the process could come from employing strains which do not need sulfur starvation. These strains feature a high respiration-to-photosynthesis ratio and a higher level of saturation irradiance compared to the *C. reinhardtii* strains currently available. These characteristics are usually found in microalgal strains with reduced antenna size, which is a very important biotechnological condition to allow penetration of the light deep through the culture layers [99–102]. Important achievements could be expected from microalgal cultures growing both mixotrophically and heterotrophically in PBRs and fermenter, respectively, in the presence of glucose. A *Chlorella* strain with such characteristics was recently studied by us under laboratory conditions, but its better H₂ performance needs to be proved under solar light in PBRs. Of course, the use of an expensive source such as glucose as a respiratory substrate to maintain anaerobiosis, and thus the functioning of the hydrogenase, halves

the efficiency of the process and strongly reduces its sustainability. Therefore, it will be important to consider the potential of much cheaper sources of organic substrates such as wastewater from sugar factories, baker's yeast, and breweries. The O₂ consumption, through respiration of organic substrates, produces H₂ with high purity (close to 98%), which strongly reduces the investment cost for H₂ purification. The use of molasses, which is very rich in glucose and sucrose, could represent an option.

In conclusion, until a hydrogenase resistant to oxygen is discovered, the selection of strains with higher resistance to oxygen and/or with high respiration-to-photosynthesis ratio represent nowadays the only realistic possibility for the success of photobiological H₂ production. This research should proceed in parallel with efforts to engineer organisms with O₂-resistant hydrogenases. Success in either direction will lead to expected improvements in technologies to: (1) increase effective conversion efficiency of photosynthesis; (2) reduce or possibly eliminate competing pathways, such as CO₂ fixation; (3) increase starch biosynthesis.

Author Contributions: Writing—original draft preparation, E.T., G.T., C.F. and A.M.S.B.; writing—review and editing, E.T. and G.T. All authors have read and agreed to the published version of the manuscript.

Funding: This work received no funds by any research agency.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: The data that support the findings of this study are available from the corresponding author upon reasonable request.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Moshood, T.D.; Nawani, G.; Mahmud, F. Microalgae biofuels production: A systematic review on socioeconomic prospects of microalgae biofuels and policy implications. *Environ. Chall.* **2021**, *5*, 100207. [\[CrossRef\]](#)
2. Masojidek, J.; Torzillo, G.; Kobližek, M. Photosynthesis in microalgae. In *Handbook of Microalgal Culture: Applied Phycology and Biotechnology*, 2nd ed.; John Wiley & Sons: Chichester, UK, 2013; pp. 21–36.
3. Gaffron, H.; Rubin, J. Fermentative and photochemical production of hydrogen in algae. *J. Gen. Physiol.* **1942**, *26*, 219–240. [\[CrossRef\]](#)
4. Martinez-Burgos, W.J.; de Souza Candeo, E.; Pedroni Medeiros, A.B.; Cesar de Carvalho, J.; Oliveira de Andrade Tanobe, V.; Soccol, C.R.; Sydney, E.B. Hydrogen: Current advances and patented technologies of its renewable production. *J. Clean. Prod.* **2021**, *286*, 124970. [\[CrossRef\]](#)
5. Wang, Y.; Yang, H.; Zhang, X.; Han, F.; Tu, W.; Yang, W. Microalgal hydrogen production. *Small Methods* **2020**, *4*, 1900514. [\[CrossRef\]](#)
6. Dębowski, M.; Dudek, M.; Zieliński, M.; Nowicka, A.; Kazimierowicz, J. Microalgal hydrogen production in relation to other biomass-based technologies—A review. *Energies* **2021**, *14*, 6025. [\[CrossRef\]](#)
7. Kosourov, S.; Nagy, V.; Shevela, D.; Jokel, M.; Messinger, J.; Allahverdiyeva, Y. Water oxidation by photosystem II is the primary source of electron for sustained H₂ photoproduction in nutrient-replete green algae. *Proc. Natl. Acad. Sci. USA* **2020**, *117*, 29629–29636. [\[CrossRef\]](#) [\[PubMed\]](#)
8. Touloupakis, E.; Faraloni, C.; Silva Benavides, A.M.; Masojidek, J.; Torzillo, G. Sustained photobiological hydrogen production by *Chlorella vulgaris* without nutrient starvation. *Int. J. Hydrog. Energy* **2021**, *46*, 3684–3694. [\[CrossRef\]](#)
9. Melis, A. Photosynthetic H₂ metabolism in *Chlamydomonas reinhardtii* (unicellular green algae). *Planta* **2007**, *226*, 1075–1086. [\[CrossRef\]](#)
10. Liu, J.Z.; Ge, Y.M.; Sun, J.Y.; Chen, P.; Addy, M.; Huo, S.H.; Li, K.; Cheng, P.F.; Ruan, R. Exogenic glucose as an electron donor for algal hydrogenases to promote hydrogen photoproduction by *Chlorella pyrenoidosa*. *Bioresour. Technol.* **2019**, *289*, 121762. [\[CrossRef\]](#) [\[PubMed\]](#)
11. Ruiz-Marin, A.; Canedo-López, Y.; Chávez-Fuentes, P. Biohydrogen production by *Chlorella vulgaris* and *Scenedesmus obliquus* immobilized cultivated in artificial wastewater under different light quality. *AMB Express* **2020**, *10*, 191. [\[CrossRef\]](#)
12. Li, L.; Litao, Z.; Jianguo, L. Proteomic analysis of hydrogen production in *Chlorella pyrenoidosa* under nitrogen deprivation. *Algal. Res.* **2021**, *53*, 102143. [\[CrossRef\]](#)
13. Grechanik, V.; Naidov, I.; Bolshakov, M.; Tsygankov, A. Photoautotrophic hydrogen production by nitrogen-deprived *Chlamydomonas reinhardtii* cultures. *Int. J. Hydrog. Energy* **2021**, *46*, 3565–3575. [\[CrossRef\]](#)

14. Amaro, H.M.; Esquivel, M.G.; Pinto, T.S.; Malcata, F.X. Hydrogen Production by Microalgae. In *Natural and Artificial Photosynthesis: Solar Power as an Energy Source*, 1st ed.; Razeghifard, R., Ed.; John Wiley and Sons, Inc.: Hoboken, NJ, USA, 2013; pp. 231–241.
15. Oliveira, F.; Araujo, A.P.C.; Romao, B.B.; Cardoso, V.L.; Ferreira, J.S.; Batista, F.R.X. Hydrogen photo-production using *Chlorella* sp. through sulfur-deprived and hybrid system strategy. *Chem. Eng. Trans.* **2015**, *43*, 301–306.
16. Jimenez-Llanos, J.; Ramirez-Carmona, M.; Rendon-Castrillon, L.; Ocampo-Lopez, C. Sustainable biohydrogen production by *Chlorella* sp. microalgae: A review. *Int. J. Hydrog. Energy* **2020**, *45*, 8310–8328. [\[CrossRef\]](#)
17. Nagarajan, D.; Dong, C.D.; Chen, C.Y.; Lee, D.J.; Chang, J.S. Biohydrogen production from microalgae-Major bottlenecks and future research perspectives. *Biotechnol. J.* **2021**, *16*, e2000124. [\[CrossRef\]](#) [\[PubMed\]](#)
18. Pongpadung, P.; Liu, J.; Yokthongwattana, K.; Techapinyawat, S.; Juntawong, N. Screening for hydrogen producing strains of green microalgae in phosphorus or sulfur deprived medium under nitrogen limitation. *Sci. Asia* **2015**, *41*, 97. [\[CrossRef\]](#)
19. Rashid, N.; Lee, K.; Han, J.I.; Gross, M. Hydrogen production by immobilized *Chlorella vulgaris*: Optimizing pH, carbon source and light. *Bioproc. Biosyst. Eng.* **2013**, *36*, 867–872. [\[CrossRef\]](#)
20. Alalayah, W.M.; Alhamed, Y.A.; Al-Zahrani, A.; Edris, G. Influence of culture parameters on biological hydrogen production using green algae *Chlorella vulgaris*. *Rev. Chim.* **2015**, *66*, 788–791.
21. Melis, A. Green alga hydrogen production: Progress, challenges and prospects. *Int. J. Hydrog. Energy* **2002**, *27*, 1217–1228. [\[CrossRef\]](#)
22. Tsygankov, A.A.; Kosourov, S.N.; Tolstygina, I.V.; Ghirardi, M.L.; Seibert, M. Hydrogen production by sulfur-deprived *Chlamydomonas reinhardtii* under photoautotrophic conditions. *Int. J. Hydrog. Energy* **2006**, *31*, 1574–1584. [\[CrossRef\]](#)
23. Batyrova, K.; Gavrisheva, A.; Ivanova, E.; Liu, J.G.; Tsygankov, A. Sustainable hydrogen photoproduction by phosphorus-deprived marine green microalgae *Chlorella* sp. *Int. J. Mol. Sci.* **2015**, *16*, 2705–2716. [\[CrossRef\]](#)
24. Rashid, N.; Lee, K.; Mahmood, Q. Bio-hydrogen production by *Chlorella vulgaris* under diverse photoperiods. *Bioresour. Technol.* **2011**, *102*, 2101–2104. [\[CrossRef\]](#)
25. Khetkorn, W.; Rastogi, R.P.; Incharoensakdi, A.; Lindblad, P.; Madamwar, D.; Pandey, A.; Larroche, C. Microalgal hydrogen production—A review. *Bioresour. Technol.* **2017**, *243*, 1194–1206. [\[CrossRef\]](#)
26. Kosourov, S.; Böhm, M.; Senger, M.; Berggren, G.; Stensjö, K.; Mamedov, F.; Lindblad, P.; Allahverdiyeva, Y. Photosynthetic hydrogen production: Novel protocols, promising engineering approaches and application of semi-synthetic hydrogenases. *Physiol. Plant* **2021**, in press. [\[CrossRef\]](#) [\[PubMed\]](#)
27. Gonzalez-Ballester, D.; Jurado-Oller, J.L.; Fernandez, E. Relevance of nutrient media composition for hydrogen production in *Chlamydomonas*. *Photosynth. Res.* **2015**, *125*, 395–406. [\[CrossRef\]](#)
28. Fakhimi, N.; Dubini, A.; Tavakoli, O.; González-Ballester, D. Acetic acid is key for synergetic hydrogen production in *Chlamydomonas*-bacteria co-cultures. *Bioresour. Technol.* **2019**, *289*, 121648. [\[CrossRef\]](#) [\[PubMed\]](#)
29. Kosourov, S.; Jokel, M.; Aro, E.M.; Allahverdiyeva, Y. A new approach for sustained and efficient H₂ photoproduction by: *Chlamydomonas reinhardtii*. *Energy Environ. Sci.* **2018**, *11*, 1431–1436. [\[CrossRef\]](#)
30. Nagy, V.; Podmaniczki, A.; Vidal-Meireles, A.; Tengölics, R.; Kovács, L.; Rákhely, G.; Scoma, A.; Tóth, S.Z. Water-splitting-based, sustainable and efficient H₂ production in green algae as achieved by substrate limitation of the Calvin–Benson–Bassham cycle. *Biotechnol. Biofuels* **2018**, *11*, 69. [\[CrossRef\]](#)
31. Krishna, P.S.; Styring, S.; Mamedov, F. Photosystem ratio imbalance promotes direct sustainable H₂ production in *Chlamydomonas reinhardtii*. *Green Chem.* **2019**, *21*, 4683–4690. [\[CrossRef\]](#)
32. Chen, M.; Zhang, J.; Zhao, L.; Xing, J.; Peng, L.; Kuang, T.; Rochaix, J.D.; Huang, F. Loss of algal Proton Gradient Regulation 5 increases reactive oxygen species scavenging and H₂ evolution. *J. Integr. Plant Biol.* **2016**, *58*, 943–946. [\[CrossRef\]](#)
33. Steinbeck, J.; Nikolova, D.; Weingarten, R.; Johnson, X.; Richaud, P.; Peltier, G.; Hermann, M.; Magneschi, L.; Hippler, M. Deletion of proton gradient regulation 5 (PGR5) and PGR5-Like 1 (PGRL1) proteins promote sustainable light-driven hydrogen production in *Chlamydomonas reinhardtii* due to increased PSII activity under sulfur deprivation. *Front. Plant Sci.* **2015**, *6*, 1–11. [\[CrossRef\]](#)
34. Pinto, T.S.; Malcata, F.X.; Arrabaça, J.D.; Silva, J.M.; Spreitzer, R.J.; Esquivel, M.G. Rubisco mutants of *Chlamydomonas reinhardtii* enhance photosynthetic hydrogen production. *Appl. Microbiol. Biotechnol.* **2013**, *97*, 5635–5643. [\[CrossRef\]](#)
35. Eilenberg, H.; Weiner, I.; Ben-Zvi, O.; Pundak, C.; Marmari, A.; Liran, O.; Wecker, M.S.; Milrad, Y.; Yacoby, I. The dual effect of a ferredoxin-hydrogenase fusion protein in vivo: Successful divergence of the photosynthetic electron flux towards hydrogen production and elevated oxygen tolerance. *Biotechnol. Biofuels* **2016**, *9*, 182. [\[CrossRef\]](#) [\[PubMed\]](#)
36. Torzillo, G.; Scoma, A.; Faraloni, C.; Ena, A.; Johanningmeier, U. Increased hydrogen photoproduction by means of a sulfur-deprived *Chlamydomonas reinhardtii* D1 protein mutant. *Int. J. Hydrog. Energy* **2009**, *34*, 4529–4536. [\[CrossRef\]](#)
37. Scoma, A.; Krawietz, D.; Faraloni, C.; Giannelli, L.; Happe, T.; Torzillo, G. Sustained H₂ production in a *Chlamydomonas reinhardtii* D1 protein mutant. *J. Biotechnol.* **2012**, *157*, 613–619. [\[CrossRef\]](#)
38. Batyrova, K.; Hallenbeck, P.C. Hydrogen Production by a *Chlamydomonas reinhardtii* strain with inducible expression of photosystem II. *Int. J. Mol. Sci.* **2017**, *18*, 647. [\[CrossRef\]](#) [\[PubMed\]](#)
39. Kosourov, S.N.; Ghirardi, M.L.; Seibert, M. A truncated antenna mutant of *Chlamydomonas reinhardtii* can produce more hydrogen than the parental strain. *Int. J. Hydrog. Energy* **2011**, *36*, 2044–2048. [\[CrossRef\]](#)
40. Xu, F.Q.; Ma, W.M.; Zhu, X.G. Introducing pyruvate oxidase into the chloroplast of *Chlamydomonas reinhardtii* increases oxygen consumption and promotes hydrogen production. *Int. J. Hydrog. Energy* **2011**, *36*, 10648. [\[CrossRef\]](#)

41. Kruse, O.; Rupprecht, J.; Bader, K.P.; Thomas-Hall, S.; Schenk, P.M.; Finazzi, G.; Hankamer, B. Improved photobiological H₂ production in engineered green algal cells. *J. Biol. Chem.* **2005**, *280*, 34170–34177. [\[CrossRef\]](#)
42. Volgusheva, A.; Styring, S.; Mamedov, F. Increased photosystem II stability promotes H₂ production in sulfur-deprived *Chlamydomonas reinhardtii*. *Proc. Natl. Acad. Sci. USA* **2013**, *110*, 7223. [\[CrossRef\]](#) [\[PubMed\]](#)
43. Oey, M.; Ross, I.L.; Stephens, E.; Steinbeck, J.; Wolf, J.; Radzun, K.A.; Kugler, J.; Ringsmuth, A.K.; Kruse, O.; Hankamer, B. RNAi knock-down of LHCBM1, 2 and 3 increases photosynthetic H₂ production efficiency of the green alga *Chlamydomonas reinhardtii*. *PLoS ONE* **2013**, *8*, e61375. [\[CrossRef\]](#)
44. Wu, S.X.; Yan, G.Y.; Xu, L.L.; Wang, Q.X.; Liu, X.L. Improvement of hydrogen production with expression of lba gene in chloroplast of *Chlamydomonas reinhardtii*. *Int. J. Hydrog. Energy* **2010**, *35*, 13419. [\[CrossRef\]](#)
45. Noone, S.; Ratcliff, K.; Davis, R.; Subramanian, V.; Meuser, J.; Posewitz, M.C.; Ghirardi, M.L. Expression of a clostridial [FeFe]-hydrogenase in *Chlamydomonas reinhardtii* prolongs photo-production of hydrogen from water splitting. *Algal. Res.* **2017**, *22*, 116–121. [\[CrossRef\]](#)
46. Kanygin, A.; Milrad, Y.; Thummala, C.; Reischneider, K.; Baker, P.; Pini, M.; Jacoby, I.; Redding, K.E. Rewriting photosynthesis, I-hydrogenase chimera that makes H₂ in vivo. *Energy Environ. Sci.* **2020**, *13*, 2903. [\[CrossRef\]](#)
47. Touloupakis, E.; Silva Benavides, A.M.; Cicchi, N.; Torzillo, G. Growth and hydrogen production of outdoor cultures of *Synechocystis* PCC 6803. *Algal. Res.* **2016**, *16*, 78–85. [\[CrossRef\]](#)
48. Kosourov, S.; Murukesan, G.; Seibert, M.; Allahverdiyeva, Y. Evaluation of light energy to H₂ energy conversion efficiency in thin films of cyanobacteria and green alga under photoautotrophic conditions. *Algal. Res.* **2017**, *28*, 253–263. [\[CrossRef\]](#)
49. Paramesh, K.; Chandrasekhar, T. Improvement of photobiological hydrogen production in *Chlorococcum minutum* using various oxygen scavengers. *Int. J. Hydrog. Energy* **2020**, *45*, 7641–7646. [\[CrossRef\]](#)
50. Su, D.; Qi, J.; Liu, X.; Wang, L.; Zhang, H.; Xie, H.; Huang, X. Enzyme-modulated anaerobic encapsulation of *Chlorella* cells allows switching from O₂ to H₂ production. *Angew. Chem. Int. Ed. Engl.* **2019**, *58*, 3992–3995. [\[CrossRef\]](#) [\[PubMed\]](#)
51. Márquez-Reyes, L.A.; Sánchez-Saavedra, M.P.; Valdez-Vazquez, I. Improvement of hydrogen production by reduction of the photosynthetic oxygen in microalgae cultures of *Chlamydomonas gloeopara* and *Scenedesmus obliquus*. *Int. J. Hydrog. Energy* **2015**, *40*, 7291–7300. [\[CrossRef\]](#)
52. Chen, J.; Li, J.; Li, Q.; Wang, S.; Wang, L.; Liu, H.; Fan, C. Engineering a chemoenzymatic cascade for sustainable photobiological hydrogen production with green algae. *Energy Environ. Sci.* **2020**, *13*, 2064. [\[CrossRef\]](#)
53. Scognamiglio, V.; Giardi, M.T.; Zappi, D.; Touloupakis, E.; Antonacci, A. Photoautotrophs–bacteria co-cultures: Advances, challenges and applications. *Materials* **2021**, *14*, 3027. [\[CrossRef\]](#) [\[PubMed\]](#)
54. Ban, S.; Lin, W.; Wu, F.; Luo, J. Algal-bacterial cooperation improves algal photolysis-mediated hydrogen production. *Bioresour. Technol.* **2018**, *251*, 350–357. [\[CrossRef\]](#) [\[PubMed\]](#)
55. He, J.; Xi, L.; Sun, X.; Ge, B.; Liu, D.; Han, Z.; Pu, X.; Huang, F. Enhanced hydrogen production through co-cultivation of *Chlamydomonas reinhardtii* CC-503 and a facultative autotrophic sulfide-oxidizing bacterium under sulfurated conditions. *Int. J. Hydrog. Energy* **2018**, *43*, 15005–15013. [\[CrossRef\]](#)
56. Fakhimi, N.; Tavakoli, O.; Marashi, S.-A.; Moghimi, H.; Mehrnia, M.R.; Dubini, A.; González-Ballester, D. Acetic acid uptake rate controls H₂ production in *Chlamydomonas*-bacteria co-cultures. *Algal. Res.* **2019**, *42*, 101605. [\[CrossRef\]](#)
57. Fakhimi, N.; Gonzalez-Ballester, D.; Fernández, E.; Galván, A.; Dubini, A. Algae-bacteria consortia as a strategy to enhance H₂ production. *Cells* **2020**, *9*, 1353. [\[CrossRef\]](#)
58. Hom, E.; Aiyar, P.; Schaeme, D.; Mittag, M.; Sasso, S. A chemical perspective on microalgal-microbial interactions. *Trends Plant Sci.* **2015**, *20*, 689–693. [\[CrossRef\]](#) [\[PubMed\]](#)
59. Fuentes, J.L.; Nores, I.G.; Cuaresma, M.; Montero, Z.; Del Valle, M.G.; Vélchez, C. Impact of microalgae-bacteria interactions on the production of algal biomass and associated compounds. *Mar. Drugs* **2016**, *14*, 100. [\[CrossRef\]](#)
60. Xu, L.; Li, D.; Wang, Q.; Wu, S. Improved hydrogen production and biomass through the co-cultivation of *Chlamydomonas reinhardtii* and *Bradyrhizobium japonicum*. *Int. J. Hydrog. Energy* **2016**, *41*, 9276–9283. [\[CrossRef\]](#)
61. Khosravitar, F. Microalgal biohydrogen photoproduction: Scaling up challenges and the ways forward. *J. Appl. Phycol.* **2020**, *32*, 277–289. [\[CrossRef\]](#)
62. Kosourov, S.N.; He, M.; Allahverdiyeva, Y.; Seibert, M. Immobilization of microalgae as a tool for efficient light utilization in H₂ production and other biotechnology applications. In *Microalgal Hydrogen Production*; Royal Society of Chemistry: London, UK, 2018; pp. 355–384.
63. Touloupakis, E.; Rontogiannis, G.; Silva Benavides, A.M.; Cicchi, B.; Ghanotakis, D.F.; Torzillo, G. Hydrogen production by immobilized *Synechocystis* sp. PCC 6803. *Int. J. Hydrog. Energy* **2016**, *41*, 15181–15186. [\[CrossRef\]](#)
64. Tsygankov, A.; Kosourov, S. Microbial BioEnergy: Hydrogen production, advances in photosynthesis and respiration. In *Immobilization of Photosynthetic Microorganisms for Efficient Hydrogen Production*; Zannoni, D., De Philippis, R., Eds.; Springer Science & Business Media: Dordrecht, The Netherlands, 2014; pp. 321–347.
65. Laurinavichene, T.V.; Fedorov, A.S.; Ghirardi, M.L.; Seibert, M.; Tsygankov, A.A. Demonstration of sustained hydrogen photoproduction by immobilized, sulfur-deprived *Chlamydomonas reinhardtii* cells. *Int. J. Hydrog. Energy* **2006**, *31*, 659–667. [\[CrossRef\]](#)
66. Kosourov, S.N.; Seibert, M. Hydrogen photoproduction by nutrient-deprived *Chlamydomonas reinhardtii* cells immobilized within thin alginate films under aerobic and anaerobic conditions. *Biotechnol. Bioeng.* **2009**, *102*, 50–58. [\[CrossRef\]](#) [\[PubMed\]](#)

67. Maswanna, T.; Lindblad, P.; Maneeruttanarungroj, C. Improved biohydrogen production by immobilized cells of the green alga *Tetraspora* sp. CU2551 incubated under aerobic condition. *J. Appl. Phycol.* **2020**, *32*, 2937–2945. [\[CrossRef\]](#)
68. Melis, A.; Zhang, L.; Forestier, M.; Ghirardi, M.L.; Seibert, M. Sustained photobiological hydrogen gas production upon reversible inactivation of oxygen evolution in the green alga *Chlamydomonas reinhardtii*. *Plant Physiol.* **2000**, *122*, 127–136. [\[CrossRef\]](#)
69. Liu, J.-Z.; Ge, Y.-M.; Xia, S.-Y.; Sun, J.-Y.; Mu, J. Photoautotrophic hydrogen production by *Chlorella pyrenoidosa* without sulfur deprivation. *Int. J. Hydrog. Energy* **2016**, *41*, 8427–8432. [\[CrossRef\]](#)
70. Li, H.; Liu, Y.; Wang, Y.; Chen, M.; Zhuang, X.; Wang, C.; Wang, J.; Hu, Z. Improved photobio-H₂ production regulated by artificial miRNA targeting psbA in green microalga *Chlamydomonas reinhardtii*. *Biotechnol. Biofuels* **2018**, *11*, 36. [\[CrossRef\]](#)
71. Ben-Zvi, O.; Dafni, E.; Feldman, Y.; Yacoby, I. Re-routing photosynthetic energy for continuous hydrogen production in vivo. *Biotechnol. Biofuels* **2019**, *12*, 266. [\[CrossRef\]](#)
72. Hwang, J.H.; Kim, H.C.; Choi, J.A.; Abou-Shanab, R.A.; Dempsey, B.A.; Regan, J.M.; Kim, J.R.; Song, H.; Nam, I.H.; Kim, S.N.; et al. Photoautotrophic hydrogen production by eukaryotic microalgae under aerobic conditions. *Nat. Commun.* **2014**, *5*, 3234. [\[CrossRef\]](#)
73. Hwang, J.-H.; Lee, M.; Kang, E.H.; Lee, W.H. Renewable algal photo H₂ production without S control using acetate enriched fermenter effluents. *Int. J. Hydrog. Energy* **2021**, *46*, 1740–1751. [\[CrossRef\]](#)
74. Sirawattanamongkol, T.; Maswanna, T.; Maneeruttanarungroj, C. A newly isolated green alga *Chlorella* sp. KLS59: Potential for biohydrogen production. *J. Appl. Phycol.* **2020**, *32*, 2927–2936. [\[CrossRef\]](#)
75. Duangjan, K.; Nakkhuntho, W.; Pekkoh, J.; Pumas, C. Comparison of hydrogen production in microalgae under autotrophic mixotrophic media. *Bot. Lith.* **2017**, *23*, 169–177.
76. Zhang, L.; He, M.; Liu, J.; Li, L. Role of the mitochondrial alternative oxidase pathway in hydrogen photoproduction in *Chlorella protothecoides*. *Planta* **2015**, *241*, 1005–1014. [\[CrossRef\]](#)
77. Shetty, P.; Boboescu, I.Z.; Pap, B.; Wirth, R.; Kovacs, K.L.; Biro, T.; Futo, Z.; White, R.A.; Maroti, G. Exploitation of algal-bacterial consortia in combined biohydrogen generation and wastewater treatment. *Front. Energy Res.* **2019**, *7*. [\[CrossRef\]](#)
78. Chader, S.; Hacene, H.; Agathos, S.N. Study of hydrogen production by three strains of *Chlorella* isolated from the soil in the Algerian Sahara. *Int. J. Hydrog. Energy* **2009**, *34*, 4941–4946. [\[CrossRef\]](#)
79. Giannelli, L.; Torzillo, G. Hydrogen production with the microalga *Chlamydomonas reinhardtii* grown in a compact tubular photobioreactor immersed in a scattering light nanoparticle suspension. *Int. J. Hydrog. Energy* **2012**, *37*, 16951–16961. [\[CrossRef\]](#)
80. Faraloni, C.; Ena, A.; Pintucci, C.; Torzillo, G. Enhanced hydrogen production by means of sulfur-deprived *Chlamydomonas reinhardtii* cultures grown in pretreated olive mill wastewater. *Int. J. Hydrog. Energy* **2011**, *36*, 5920–5931. [\[CrossRef\]](#)
81. Song, W.; Rashid, N.; Choi, W.; Lee, K. Biohydrogen production by immobilized *Chlorella* sp. using cycles of oxygenic photosynthesis and anaerobiosis. *Bioresour. Technol.* **2011**, *102*, 8676–8681. [\[CrossRef\]](#)
82. Zhu, X.-G.; Long, S.P.; Ort, D.R. What is the maximum efficiency with which photosynthesis can convert solar energy into biomass? *Curr. Opin. Biotechnol.* **2008**, *19*, 153–159. [\[CrossRef\]](#)
83. Melis, A. Solar energy conversion efficiencies in photosynthesis: Minimizing the chlorophyll antennae to maximize efficiency. *Plant Sci.* **2009**, *177*, 272–280. [\[CrossRef\]](#)
84. Finazzi, G.; Barbagallo, R.P.; Bergo, E.; Barbato, R.; Forti, G. Photoinhibition of *Chlamydomonas reinhardtii* in State 1 and State 2: Damages to the photosynthetic apparatus under linear and cyclic electron flow. *J. Biol. Chem.* **2001**, *276*, 22251. [\[CrossRef\]](#)
85. Antal, T.K.; Krendeleva, T.E.; Laurinavichene, T.V.; Makarova, W.; Ghirardi, M.L.; Rubin, A.B.; Tsygankov, A.A.; Seibert, M. The dependence of algal H₂ production on photosystem II and O₂ consumption activities in sulphur-deprived *Chlamydomonas reinhardtii* cells. *Biochem. Biophys. Acta* **2003**, *1607*, 153–160.
86. Burlacot, A.; Sawyer, A.; Cuiné, S.; Auroy-Tarrago, P.; Blangy, S.; Happe, T.; Peltiera, G. Flavodiiron-mediated O₂ photoreduction links H₂ production with CO₂ fixation during the anaerobic induction of photosynthesis. *Plant Physiol.* **2018**, *177*, 1639–1649. [\[CrossRef\]](#)
87. Antal, T.K.; Kukarskikh, G.P.; Volgusheva, A.A.; Krendeleva, T.E.; Tyystjärvi, E.; Rubin, A.B. Hydrogen photoproduction by immobilized S-deprived *Chlamydomonas reinhardtii*: Effect of light intensity and spectrum, and initial medium pH. *Algal. Res.* **2016**, *17*, 38–45. [\[CrossRef\]](#)
88. Strasser, R.; Srivastava, A.; Govindjee. Polyphasic chlorophyll a fluorescence transient in plants and cyanobacteria. *Photochem. Photobiol.* **1995**, *61*, 32–42. [\[CrossRef\]](#)
89. Faraloni, C.; Torzillo, G. Xanthophyll cycle induction by anaerobic conditions under low light in *Chlamydomonas reinhardtii*. *J. Appl. Phycol.* **2013**, *25*, 1457–1471. [\[CrossRef\]](#)
90. Pongpadung, P.; Zhang, L.; Sathasivam, R.; Yokthongwattana, K.; Juntawon, N.; Liu, J. Stimulation of hydrogen photoproduction in *Chlorella sorokiniana* subjected to simultaneous nitrogen limitation and sulfur- and/or phosphorus-deprivation. *J. Pure Appl. Microbiol.* **2018**, *12*, 1719–1727. [\[CrossRef\]](#)
91. Torzillo, G.; Chini Zittelli, G. Tubular photobioreactors. Products and biorefinery design. In *Algal Biorefineries*; Prokop, A., Bajpai, R.K., Zappi, M.E., Eds.; Springer International Publishing: Cham, Switzerland, 2015; Volume 2.
92. Scoma, A.; Giannelli, L.; Faraloni, C.; Torzillo, G. Outdoor H₂ production in a 50-L tubular photobioreactor by means of a sulfur-deprived culture of the microalga *Chlamydomonas reinhardtii*. *J. Biotechnol.* **2012**, *157*, 620–627. [\[CrossRef\]](#) [\[PubMed\]](#)

93. Lindblad, P.; Fuente, D.; Borbe, F.; Cicchi, B.; Conejero, J.A.; Couto, N.; Čelešnik, H.; Diano, M.; Dolinar, M.; Esposito, S.; et al. CyanoFactory, a European consortium to develop technologies needed to advance cyanobacteria as chassis for production of chemicals and fuels. *Algal. Res.* **2019**, *41*, 101510. [\[CrossRef\]](#)
94. Siebert, M.; Torzillo, G. *Microalgal Hydrogen Production: Achievements and Perspectives*; The Royal Society of Chemistry: Cambridge, UK, 2018.
95. James, B.D.; Baum, G.N.; Perez, J.; Baum, K.N. *Technoeconomic Analysis of Photoelectrochemical (PEC) Hydrogen Production*; (US DOE Contract no. GS-10F-009J); Directed Technologies Inc.: Arlington, VA, USA, 2009.
96. Show, K.-Y.; Yan, Y.; Zong, C.; Guo, N.; Chang, J.-S.; Lee, D.-J. State of the art and challenges of biohydrogen from microalgae. *Bioresour. Technol.* **2019**, *289*, 121747. [\[CrossRef\]](#)
97. Silva Benavides, A.M.; Campos Rudin, M.; Villalobos, N.; Touloupakis, E.; Torzillo, G. Growth and hydrogen production by three *Chlamydomonas* strains cultivated in a commercial fertilizer. *Int. J. Hydrog. Energy* **2019**, *44*, 9849–9855. [\[CrossRef\]](#)
98. Frowijn, L.S.F.; van Sark, W.G.J.H.M. Analysis of photon-driven solar-to-hydrogen production methods in the Netherlands, *Sustain. Energy Technol. Assess.* **2021**, *48*, 101631.
99. Perrine, Z.; Negi, S.; Sayre, R. Optimization of photosynthetic light energy utilization by microalgae. *Algal. Res.* **2012**, *1*, 134–142. [\[CrossRef\]](#)
100. Cazzaniga, S.; Dall'Osto, L.; Szaub, J.; Ballottari, M.; Purton, S.; Bassi, R. Domestication of the green alga *Chlorella sorokiniana*: Reduction of antenna size improves light-use efficiency in a photobioreactor. *Biotechnol. Biofuels* **2014**, *7*, 157–170. [\[CrossRef\]](#)
101. Shin, W.-S.; Lee, B.; Jeong, B.-R.; Chang, Y.K. Truncated light-harvesting chlorophyll antenna size in *Chlorella vulgaris* improves biomass productivity. *J. Appl. Phycol.* **2016**, *28*, 3193–3202. [\[CrossRef\]](#)
102. Hu, G.-R.; Fan, Y.; Zhen, Y.-L.; Xu, F.; Zhang, L.; Li, F.-L. Photoprotection capacity of microalgae improved by regulating the antenna size of high-harvesting complexes. *J. Appl. Phycol.* **2020**, *32*, 1027–1039. [\[CrossRef\]](#)