

Microbial residue distribution in microaggregates decreases with stand age in subtropical plantations

Yanli Jing^{1,2}, Xuechao Zhao^{2,3}, Shengen Liu⁴, Peng Tian⁵, Zhaolin Sun⁵, Longchi Chen² and Qingkui Wang^{2,*}

¹ Ecology Security and Protection Key Laboratory of Sichuan Province, Mianyang Normal University, Mianyang, 621000, China

² Huitong Experimental Station of Forest Ecology, CAS Key Laboratory of Forest Ecology and Management, Institute of Applied Ecology, Shenyang, 110016, China

³ University of Chinese Academy of Sciences, Beijing 100049, China

⁴ College of Biological and Pharmaceutical Sciences, China Three Gorges University, Yichang, 443000, China.

⁵ School of Forestry & Landscape Architecture, Anhui Agricultural University, Hefei 230036, China

*Corresponding author: Qingkui Wang

E-mail address: qwang@iae.ac.cn

Tel: +8624 8397 0206




Figure S1. Microbial residues in SOC across the Chinese fir chronosequence in southern China. Values are the mean \pm SE.




Figure S2. Microbial residues in topsoil and subsoil at a global scale. Data are from Ni et al. [1].

Reference

1. Ni, X.; Liao, S.; Tan, S.; Peng, Y.; Wang, D.; Yue, K.; Wu, F.Z.; Yang, Y. The vertical distribution and control of microbial necromass carbon in forest soils. *Glob. Ecol. Biogeogr.* **2020**, *29*, 1829–1839.