

Supplemental Table S1.

	Reference cohort (N = 20)	Non-COVID-19 (N = 44)	Non-hospitalized	Hospitalized			
			COVID-19 (N = 44)	COVID-19 (N = 53)	p-value	p-value	p-value
			Group 1	Group 2	Group 3	Group 4	1 vs. 2
Demographics							
Age [years]	55.6 (44.9, 67.6)	63.1 (50.1, 77.5)	37.2 (27.7, 52.7)	71.2 (54.1, 79.2)	0.143	0.040	<0.001
Female	12 (60%)	23 (52%)	19 (43%)	27 (51%)	0.600	0.716	0.541
Comorbidity							
CKD	1/19 (5%)	9 (20%)	3 (7%)	11 (21%)	0.258	0.461	0.080
CHD	1/19 (5%)	5 (11%)	4 (9%)	11 (21%)	0.658	0.610	0.160
Hypertension	5/19 (26%)	18 (41%)	8 (18%)	28 (53%)	0.394	0.711	<0.001
T2DM	2/19 (11%)	3 (7%)	6 (14%)	13 (25%)	0.633	0.078	0.207
COPD	0/19 (0%)	3 (7%)	0/43 (0%)	6 (11%)	0.547	1.000	0.031
Asthma	1/19 (5%)	1 (2%)	2 (5%)	9 (17%)	0.516	0.104	0.104
MD	1/19 (5%)	7/43 (16%)	1 (2%)	1 (2%)	0.416	0.004	1.000
OAC	1/19 (5%)	9/43 (21%)	0/43 (0%)	10 (19%)	0.155	0.112	0.002
Treatment							
Length of stay	1.0 (1.0, 4.2)	3.0 (1.0, 9.0)	1.0 (1.0, 1.0)	12.0 (8.0, 20.0)	0.204	0.129	<0.001

CKD = chronic kidney disease; CHD=coronary heart disease; T2DM=type 2 diabetes mellitus; COPD=chronic obstructive pulmonary disease; MD=malignant disease; OAC=oral anticoagulation.