

**Table S1. Key demographic characteristics of the participants, not all participants responded to all questions**

	<b>Persistent oral HPV16 group (n=13)</b>	<b>Transient oral HPV16 group (n=26)</b>
<b>Age, mean (range)</b>	25.3 (18-29)	27.0 (19-39)
<b>Smoking, current</b>		
No smoking	3	11
1-10 /day	5	7
11-20/day	3	6
>20/day	-	1
<b>Alcohol use</b>		
never	1	4
2-3 dose (12g)/week	1	3
1 dose/week	2	9
1 dose or less/month	6	9
<b>Mode of delivery (this pregnancy)</b>		
alatie	10	17
keisarin leikkaus	3	9
<b>Age of first pregnancy (mean <math>\pm</math>SD)</b>	22.8 $\pm$ 3.8	23.3 $\pm$ 3.5
<b>Number of given births (mean <math>\pm</math> SD)</b>	1.27 $\pm$ 0.47	0.32 $\pm$ 0.48
<b>Age at starting oral contraceptives</b>		
never used	1	1
14-16	3	13
17-19	6	9
$\geq$ 20	1	2
<b>Age (years) at first sexual intercourse</b>		
$\leq$ 16	6	17
> 16	5	8
<b>Practice of oral sex</b>		
yes	10	20
no	1	5
<b>Practise of anal sex</b>		
yes	3	5
no	8	20
<b>Number of sexual intercourse/mo</b>		
2-4	5	13
5-10	4	9
>10	2	3
<b>Number of lifetime sexual partners</b>		
0-2	4	7
3-5	3	4
6-10	2	7
>10	2	7
<b>History of STDs (genital warts excluded)</b>		
yes	1	8
no	12	18
<b>History of genital warts</b>		
yes	6	11
no	5	14