



Supplementary Materials: Deep Eutectic Solvents for Improving the Solubilization and Delivery of Dapsone

Sonia Trombino, Carlo Siciliano, Debora Procopio, Federica Curcio, Annarita S. Laganà, Di Gioia Maria Luisa, and Cassano Roberta

Table S1. Experimental POM images of the prepared DESs.

| DESs | HBA | HBD | HBA/HBD molar ratio | Aspect | POM images |
|--------|-----|-----|---------------------|--|------------|
| CC:AA | CC | AA | 2:1 | Viscous yellow liquid | |
| CC:PG | CC | PG | 1:3 | Transparent liquid | |
| | | | 1:1 | Viscous liquid that rapidly solidified at rt | |
| CC:DAP | CC | DAP | 1:2 | Solid | |




Figure S1. FTIR spectra of choline chloride, ascorbic acid and CC:AA DES (2:1).







Figure S2. FTIR spectra of choline chloride, propylene glycol and CC:PG DES (1:3).

a



b**c**




Figure S3. ¹³C NMR spectra of choline chloride (a), dapsone (b), ChCl-PG DES (c) and DES-DAP ternary system (d).




Figure S4. POM images of (DRUG + PG) and (DRUG + DESs) at rt.




Figure S5. DSC thermograms of (1) CC:AA DES, (2) pure CC and (3) pure AA.