

Article

New Contributions to the *Ericion umbellatae* Alliance in the Central Iberian Peninsula

José C. Piñar Fuentes ¹, Mauro Raposo ² , Carlos J. Pinto Gomes ² , Sara del Río González ³, Giovanni Spampinato ⁴  and Eusebio Cano ^{1,*} 

¹ Department of Animal and Plant Biology and Ecology, Section of Botany, University of Jaén, Las Lagunillas s/n, 23071 Jaén, Spain; jcpfuentes@gmail.com

² Department of Landscape, Environment and Planning, Institute for Mediterranean Agrarian and Environmental Sciences (ICAAM), School of Science and Technology, University of Évora (Portugal), Rua Romão Ramalho, n° 59, 7000-671 Évora, Portugal; mrraposo@uevora.pt (M.R.); cpgomes@uevora.pt (C.J.P.G.)

³ Department of Biodiversity and Environmental Management (Botany), Faculty of Biological and Environmental Sciences, Campus de Vegazana s/n, University of León, 24071 León, Spain; sara.delrio@unileon.es

⁴ Department of Agraria, “Mediterranea” University of Reggio Calabria, Loc. Feo di Vito, 89122 Reggio Calabria, Italy; gspampinato@unirc.it

* Correspondence: ecano@ujaen.es



Citation: Piñar Fuentes, J.C.; Raposo, M.; Pinto Gomes, C.J.; del Río González, S.; Spampinato, G.; Cano, E. New Contributions to the *Ericion umbellatae* Alliance in the Central Iberian Peninsula. *Sustainability* **2021**, *13*, 5639. <https://doi.org/10.3390/su13105639>

Academic Editor: Imre J. Holb

Received: 19 January 2021

Accepted: 30 April 2021

Published: 18 May 2021

Publisher’s Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Abstract: The study of heathlands dominated by *Erica australis*, *E. umbellata* and *Cistus populifolius* in the centre and west of the Iberian Peninsula allows us to separate the eight shrubland communities. The taxonomic analysis of *E. australis* distinguishes two subspecies: *E. australis* subsp. *australis* and *E. australis* subsp. *ragonensis*. The statistical treatment confirms the differences between the sub-alliances *Ericenion aragonensis* and *Ericenion umbellatae*. This ecological, bioclimatic, biogeographical and floristic study has allowed us to differentiate three new associations from the remaining five: TCp = *Teucro oxylepis-Cistetum populifolii* nova. HEau = *Halimio ocymoidis-Ericetum australis* nova. DEu = *Drosophyllo lusitanicae-Ericetum umbellatae* nova. ECp = *Erico australis-Cistetum populifolii* Rivas Goday 1964. PCp = *Polygalo microphyllae-Cistetum populifolii* Rivas Goday 1964. HEa = *Halimio ocymoidis-Ericetum aragonensis* Rivas-Martínez 1979. HEu = *Halimio ocymoidis-Ericetum umbellatae* Rivas Goday 1964. UEu = *Ulici eriocladi-Ericetum umbellatae*.

Keywords: association; taxonomy; heathlands; habitats; shrubs

1. Introduction

Humid and subhumid areas contain woodlands of *Quercus broteroi* (Cout.) Rivas-Martínez & C.Sáenz (Portuguese oak), *Quercus canariensis* and *Quercus mariannica* Vicioso, *Quercus suber* L. (cork oak) and *Quercus pyrenaica* Willd. (Pyrenean oak.) [1–3], all originating within the vegetation community dynamic of the Ericaceae family: *Arbutus unedo* L., *Erica australis* subsp. *australis* L., *Erica umbellata* Loefl. ex L., *Erica scoparia* L., and some communities of the Cistaceae family, *Cistus populifolius* L. Until now these communities in the central-western Iberian Peninsula have been included in the associations *Halimio ocymoidis-Ericetum aragonensis*, *Halimio ocymoidis-Ericetum umbellatae*, *Ulici eriocladi-Ericetum umbellatae*, and *Polygalo microphyllae-Cistetum populifolii* [4–13].

Heathlands are one of the most extensive and important habitats on the planet [14,15], and are among the most representative plant communities on the European continent with an Atlantic bioclimate and influence [5,16]. These formations are linked to oceanic environments, with abundant rainfall and short periods of summer drought in the case of the southern Iberian Peninsula [14,17–20], and in acidic, oligotrophic soils subject to erosion and fire [21].

From a conservation point of view, these communities have received a different treatment on the Iberian Peninsula [15] and were considered as mere replacement stages [5,21], whereas in the rest of Europe these plant formations—which harbour less diversity—are more highly regarded [15] in conservation terms. In the Habitats Directive, *Erica umbellata*-dominated heaths correspond to code 4030 “European dry heaths”.

The associations dominated by *Erica* spp., *Ulex* spp. and *Cistus populifolius* are typical of subhumid and humid ombro-types and are consequently fragile in the face of climate change, since they are affected by the increase in temperature and rainfall irregularity, as are the endemic taxa in these associations, and consequently the endemic taxa present in mountain areas [22–24]. These thickets dominated by the genus *Erica* are typical of rainy environments and are at their optimum in temperate Europe, although they extend to the sub-Mediterranean mountains, and connect catenally towards dry-subhumid ombro-types, with micro-forests of *Juniperus oxycedrus* subsp. *badia* (H. Gay) according to Debeaux [25,26].

On the Iberian Peninsula, this type of habitat has encountered problems in its interpretation, largely due to its extension and its bioclimatic and biogeographical diversity. The classifications proposed have disregarded the typical Mediterranean heaths, with Atlantic nuances, of the southwest Iberian Peninsula, which are of enormous floristic and biogeographical interest [15]. It should be noted that, of the three heathland subtypes considered, these Mediterranean dry heaths are the richest in endemisms.

The associations studied should be considered habitats of interest for conservation, since they dominate the middle and high mountains and include rocky habitats and wetlands with a predominance of endemic, rare and in some cases threatened species such as *Coincya longirostra* (Boiss.) Greuter & Burdet, *Digitalis mariana* Boiss., *Jasione crispa* subsp. *tomentosa*, *Jasione crispa* subsp. *mariana* (A.DC.) Rivas Mart., *Dianthus lusitanus* Brot., *Adenocarpus hispanicus* subsp. *argyrophyllus* (Rivas Goday), *Drosophyllum lusitanicum* (L.) *Solenopsis laurentia* (L.) C. Presl, *Pinguicula lusitanica* L.

According to [22], *Erica australis* L. is highly variable; the species *Erica aragonensis* Willk. and *Erica australis* subsp. *aragonensis* (Willk.) Cout. Are not recognized and claimed to be a montane ecotype, although any convincing arguments to synonymise these taxa to *E. australis* are not provided; however, [23] the subsp. *Aragonensis* is accepted. In our opinion the character of extrorse anthers in the subsp. *aragonensis* compared to non-extrorse anthers in the subsp. *australis* is clear. We, therefore, maintain the subsp. *aragonensis*, which has a more northerly distribution than the subsp. *australis*.

The main objective of the present work is to highlight the plant diversity of scrubs dominated by different species of the genus *Erica* in the centre-south of the Iberian Peninsula within the Mediterranean climate domain. On the other hand, from the point of view of the sustainability of forests and accompanying shrublands, it is important to know how these heathlands provide the environmental conditions necessary for the establishment of plant stages of greater biomass. It is, therefore, necessary to know the diversity and structure of Mediterranean dry heaths for their conservation and sustainability over time.

2. Materials and Methods

2.1. Study Area

The territory in the study is located in the central-western Iberian Peninsula (Figure 1) and is dominated by siliceous materials, Palaeozoic slates, metamorphic quartzites and plutonic rocks (granite); it has a Mediterranean bioclimate with a thermo-type that ranges between the thermo- and the supra-Mediterranean, and an ombro-type between dry and humid.




Figure 1. Location of the study area including the biogeographical provinces.

2.2. Sampling

An analysis was made of 232 heathland relevés dominated by *Erica umbellata*, *E. australis* and *Cistus populifolius* in the centre and west of the Iberian Peninsula, obtained from our own field research (56 samples) and various publications (176 samples) (Table 1). These relevés belong to ten associations: SDI = *Saturejo salzmannii-Drosophylletum lusitanicii*; TCl = *Teucrio mariani-Cistetum laurifolii*; TCp = *Teucrio oxylepis-Cistetum populifolii* (Figure 2); HEau = *Halimio ocymoidis-Ericetum australis*; ECp = *Erico australis-Cistetum populifolii*; PCp = *Polygalo microphylla-Cistetum populifolii*; HEa = *Halimio ocymoidis-Ericetum aragonensis*; DEu = *Drosophyllo lusitanicae-Ericetum umbellatae*; HEu = *Halimio ocymoidis-Ericetum umbellatae*; UEu = *Ulici eriocladi-Ericetum umbellatae*. For the authorship of the taxa, we have followed Flora Ibérica, Flora Western Andalusia and Eastern Andalusia Flora. For the authorship of syntaxa, we have followed [12].

Our own samplings were taken following the Braun-Blanquet method, modified by Van der Maarel [24].

Table 1. Origin of the vegetation samples used in this work.

Plant Community	Bibliography	Nº of Relevés
SD1	Galán de Mera, A. (1993)	5
	Junta de Andalucía	2
	Quezel P. et. al. (1988)	5
TCp	Own relevés	18
	Rodríguez Marzal (2006)	3
TCI	Junta de Andalucía	1
	Velasco Negueruela, A. (1981)	6
	Own relevés	4
HEa	Own relevés	12
ECp	Rivas Goday, S. (1964)	5
	Belmonte M.D. (2008)	13
	Muñoz, A.F.; Santa-Bárbara, C. & Vicent, C. (2008)	24
	Rufo Nieto (2009)	13
PCp	Melendo Luque, M. (1998)	7
	Rivas Goday, S. (1964)	8
	Belmonte M.D. (2008)	13
HEu	Rivas Goday, S. (1964)	5
	Belmonte M.D. (2008)	16
	Melendo Luque, M. (1998)	4
	Muñoz, A.F.; Santa-Bárbara, C. & Vicent, C. (2008)	9
Ueu	Muñoz, A.F.; Santa-Bárbara, C. & Vicent, C. (2008)	4
	Rivas-Martínez, S. (1979)	8
	Rufo Nieto (2009)	4
	Junta de Andalucía	21
DEu	Own relevés	22

SD1 = *Saturejo salzmannii*-*Drosophylletum lusitanici*. TCI = *Teucrio mariani*-*Cistetum laurifolii*. TCp = *Teucrio mariani*-*Cistetum populifolii*. HEau = *Halimio ocyoidis*-*Ericetum australis*. ECp = *Erico australis*-*Cistetum populifolii*. PCp = *Polygalo microphyllae*-*Cistetum populifolii*. HEa = *Halimio ocyoidis*-*Ericetum aragonensis*. DEu = *Drosophylo lusitanicae*-*Ericetum umbellatae*. HEu = *Halimio ocyoidis*-*Ericetum umbellatae*. Ueu = *Ulici eriocladi*-*Ericetum umbellatae*.

2.3. Data Analyses

Subsequently, and based on the geographic and topographic information contained in these relevés, each sampling was geo-referenced and implemented in a geographic information system (GIS) using the corresponding maps (Appendix A). They were then matched with the place names and the rest of the data from the samplings in the bibliography in order to obtain the different climatic and bioclimatic variables for each sampling. The figures for altitude, orientation and slope were determined by plotting the previously geo-referenced points on a digital terrain map (DTM) with a pixel resolution of 30 × 30 metres for the points in Portugal and Morocco, and on a digital terrain model with a resolution of 10 × 10 metres for the points in Spain. The numerical values for altitude, orientation and slope were obtained from the DTM for each sampling point by means of numerical classification methods. An ascending hierarchical classification was applied for the ordination and comparison of the different samples using Ward's distance. For a better understanding of the results, a multivariate ordination analysis was carried out by means of detrended correspondence analysis.




Figure 2. Distribution of plant communities dominated by *Teucrium oxylepis* subsp. *marianum*. TCI = *Teucro mariani-Cistetum laurifolii* is located in the most inland areas of the Iberian Peninsula with a major continental influence. TCp = *Teucro mariani-Cistetum populifolii* is located in more peripheral zones with a greater Atlantic influence. TCp-Cl = *Teucro mariani-Cistetum populifolii* is a colder and more continental variant with *Cistus laurifolius*.

3. Results

We studied the communities of *Erica umbellata* Loefl. ex L., *Erica australis* L. subsp. *australis*, *Erica australis* L. subsp. *aragonensis* (Willk.) Cout. and *Cistus populifolius* L. in the central Iberian Peninsula. The statistical analysis of the communities dominated by these taxa (Figures 2–5) reveals the existence of two groups, G1 and G2, corresponding to the suballiances *Ericenion aragonensis* Rivas Martínez 1979 and *Ericenion umbellatae* Rivas Martínez 1979. The association *Polygalo microphyllae-Cistetum populifolii* (PCp1-PCp8) was described by [1] for shady Silurian sites in Extremadura, whose relevés PCp1-PCp8 are far removed from our group of relevés and [7] for the eastern Sierra Morena, and from the relevés of [1] from which he describes the association *Erico australis-Cistetum populifolii* (ECp1-ECp5), and which have a more thermophilous character than the former (Figure 4).




Figure 3. Cluster of heathland associations in the central-western Iberian Peninsula. The hierarchical classification used is based on Euclidean distance, using Ward's agglomerative method.




Figure 4. DECORANA (Detrended analyse of variance) ordination of heathland communities in the central-western Iberian Peninsula.




Figure 5. DCA ordination of heathland communities in the central-western Iberian Peninsula.

The relevés dominated by *C. populifolius* in eastern Sierra Morena differ floristically from the communities described by [4] in Extremadura, as they quite frequently include *Teucrium oxylepis* (endemic) and *Cistus laurifolius*, which grow in shady sites and foothills more influenced by the cold winds from the Manchegan plateau. We therefore propose the association *Teucro mariani-Cistetum populifolii nova* (Table 2 rel. TCp1-TCp18 typus rel. TCp15*).

The relevés corresponding to the association described by [5], *Ulici eriocladi-Ericetum umbellatae* (UEu1-UEu8), are clearly separated and floristically, ecologically, dynamically and biogeographically differentiated from the other communities of *E. australis*. The association *Halimio ocymoidis-Ericetum aragonensis* (HEa1-HEa15), described by [5] for the territories of Guadalajara and Segovia, shows significant differences from the heathlands of *E. australis* in the Marianica range due to the absence in Marianican territories of *E. australis* subsp. *aragonensis*, *Luzula lactea* (Link) E.Mey., *Genista pilosa* L., *Deschampsia flexuosa* subsp. *iberica* Rivas Mart. and *Helianthemum pulverulentum* Pers. We therefore propose the association *Halimio ocymoidis-Ericetum australis nova* (Table 3 rel. HEau1-HEau11, typus rel. HEau4*), which has a more southerly character than the previous association, for supra-Mediterranean environments in the eastern Mariánico-Monchiquensean sector; this heathland represents a dynamic stage of the woodlands of *Sorbo terminalis-Quercetum pyrenaicae*. This association is statistically close to *Erico australis-Cistetum populifolii*; however, the latter has a more eastern and thermophilous character ([4], Table 67, page 446).

In the group of communities of *E. umbellata*, the relevés given by [4,5] in the Sierra de San Pedro (Cáceres) and Herrera del Duque (Badajoz), León and Zamora are gathered in a single compact group (HEu12-HEu21) corresponding to *Halimio ocymoidis-Ericetum umbellatae*; this association was described by [4] in highly oligotrophic and nutrient-poor soils. It has floristic, dynamic and biogeographical differences with the communities of *Erica umbellata* in the eastern Sierra Morena; the Leonese and Zamoran territories include the species *Halimium alysoides*, *E. australis* subsp. *aragonensis*, *Ulex minor*, *Genista micrantha*, and *Agrostis setacea*, which have a more oceanic character and are not present in the eastern Sierra Morena. The relevés of [7,10] are grouped with those taken recently by us in the province of Ciudad Real, where they include *E. umbellata*, *Halimium ocymoides*, *Tuberaria lignosa*, *Pterospartum tridentatum*, *Drophyllum lusitanicum*, *Lavandula pedunculata* and occasionally *Festuca elegans*, *Coincyia longirostra* and *Juniperus oxycedrus* subsp. *badia*. We therefore propose the association *Drosophyllo lusitanicae-Ericetum umbellatae nova* (Table 4 rel. DEu1-DEu22 typus rel. DEu6 *) growing on highly washed soils (distic lithosols) with a total absence of organic matter. This association is distributed throughout the eastern Mariánico-Monchiquensean sector.

Table 2. Teucrio mariani-Cistetum populifolii nova

Table 2. *Cont*

Table 2. Cont.

<i>Quercus faginea</i> subsp. <i>faginea</i>								2	+
<i>Arenaria montana</i> subsp. <i>montana</i>								1	
<i>Tamus communis</i>		+						+	
<i>Epipactis helleborine</i> subsp. <i>helleborine</i>								+	
<i>Epipactis helleborine</i> subsp. <i>tremolsii</i>									+
<i>Quercus pyrenaica</i>								+	
<i>Sorbus torminalis</i>								+	
Companion species									
<i>Pinus pinaster</i>								5	4
<i>Rosmarinus officinalis</i>	1		2	+		1	1		5
<i>Thapsia villosa</i>	+		+	+		1	1	2	+
<i>Thymus mastichina</i>		1		1				+	+
<i>Carlina corymbosa</i> subsp. <i>corymbosa</i>								+	2
<i>Agrostis capillaris</i> x <i>castellana</i>								3	+
<i>Cistus albidus</i>	+		2				+		
<i>Halimum halimifolium</i>								+	2
<i>Castanea sativa</i>								2	+
<i>Dactylis lusitanica</i>	+	+	1						
<i>Holcus lanatus</i>								2	+
<i>Rubus ulmifolius</i>								+	2
<i>Asphodelus albus</i> subsp. <i>albus</i>		+						1	
<i>Briza maxima</i>									2
<i>Helichrysum stoechas</i>	+							+	+
<i>Sanguisorba verrucosa</i>								+	+
<i>Sedum forsterianum</i>					1		+		
<i>Centaurea melitensis</i>		1						1	+
<i>Festuca rothmaleri</i>									1
<i>Leucanthemum sylvaticum</i>	1								
<i>Magydaris panacifolia</i>	1								
<i>Origanum virens</i>								+	+
<i>Avenula marginata</i> subsp. <i>sulcata</i>								+	+
<i>Celtica gigantea</i>								+	+
<i>Dactylis glomerata</i> subsp. <i>hispanica</i>								+	+
<i>Ranunculus gramineus</i>							+	+	

Table 2. Cont.

Rosa pouzinii	+												+
Sanguisorba minor													+
Andryala integrifolia													+
Aristolochia pistolochia	+												+
Arrhenatherum elatius subsp. bulbosum		+											
Campanula rapunculus		+											
Cardamine hirsuta													+
Clinopodium vulgare subsp. arundanum													+
Conopodium mariannum		+											
Dorycnium pentaphyllum	+												
Eryngium tenue													+
Iris planifolia		+											
Linum suffruticosum subsp. suffruticosum		+											
Picris comosa		+											
Piptatherum miliaceum		+											
Srophularia scorodonia													+
Thapsia garganica		+											
Tulipa sylvestris													+

Typus *.

Table 3. Halimio ocymoidis-Ericetum australis nova.

	HEau1	HEau2	HEau3	HEau *	HEau5	HEau6	HEau7	HEau8	HEau9	HEau10	HEau11	HEau12
Altitude m.	820	1022	1061	1249	1229	742	752	800	760	1160	800	750
Area m ²	300	200	200	200	200	100	200	200	200	400	100	200
Coverage %	80	95	80	90	80	90	100	75	90	85	95	95
Slope %	15	8	8.5	8	10	8.5	4	20	20	25	20	20
Orientation	E	NE		NW	N		E-SE	N	N	S	N	S
Medium height veg. in m.	1	1.5	1.5	1.5	1.20	0.70	1.8	1.0	1.0	1.0	1.0	1.0
Characteristic association species and higher syn-taxonomic units												
<i>Erica australis</i>	4	5	4	5	4	5	5	5	5	4	5	5
<i>Halimium ocymoides</i>	1	1	1	2	1	1	1	1	+	2	1	1
<i>Cistus pulchellus</i>	+	+	1	1	1	1	4		1	1	1	2

Table 3. *Cont*

Table 3. Cont.

	HEau1	HEau2	HEau3	HEau *	HEau5	HEau6	HEau7	HEau8	HEau9	HEau10	HEau11	HEau12
<i>Arrhenatherum elatius</i> subsp. <i>bulbosum</i>				+								
<i>Narcissus triandrus</i>						+						
<i>Arrhenatherum album</i>						+						
<i>Cardamine hirsuta</i>								2				
<i>Scirpus holoschoenus</i>							+					
<i>Salix atrocinerea</i>							+					
<i>Osyris alba</i>								1				
<i>Teucrium haenseleri</i>								+				
<i>Cytisus multiflorus</i>								+				
<i>Pistacia terebinthus</i>										+		
<i>Juniperus oxycedrus</i>											+	

Typus *.

Table 4. Drosophyllo lusitanicae-Ericetum umbellatae nova.

	DEu1	DEu2	DEu3	DEu4	DEu5	DEu6 *	DEu7	DEu8	DEu9	DEu10	DEu11	DEu12	DEu13	DEu14	DEu15	DEu16	DEu17	DEu18	DEu19	DEu20	DEu21	DEu22
Altitude m.	795	1233	982	1042	1051	1061	1074	1095	742	599	830	1010	800	730	800	800	700	750	720	750	760	740
Area m ²	300	400	100	50	10	40	20	40	100	100	200	100	200	200	200	200	300	200	150	200	150	100
Coverage %	40	60	80	50	80	70	70	90	80	80	90	95	85	80	80	75	75	80	80	75	85	90
Slope %	12	20	4	6	8	8	6	4	10	8	30	5	15	15	15	30	35	30	25	10	10	15
Orientation	SW	N	S	NE	NE	NE	NE	NW	W	W	W	W	N-NE	N	N	S	S	W	SW	N	NE	NW
Medium height veg. in m.	0.80	1.5	0.70	0.40	0.60	0.50	0.40	0.50	0.25	0.35		0.60										
Characteristic association species and higher syntaxonomic units																						
<i>Erica umbellata</i>	4	5	4	4	5	5	5	5	4	4	5	5	4	4	5	5	4	5	4	4	4	4
<i>Halimium ocymoides</i>	4	4	4	4	4	4	1	4	1	4	1	4	1	4	4	5	4	4	5	4	1	5
<i>Erica australis</i>	1	+	4	1	1	1	1	1	1	1	1	1	1	1	1	1	+	1				
<i>Drosophyllum lusitanicum</i>	4	5	4	5	4	4	4															
<i>Cistus populifolius</i>				1	1	+		1												+		
<i>Tuberaria lignosa</i>	+																		1		+	1
<i>Lavandula luisieri</i>																						
<i>Pterospartum tridentatum</i> subsp. <i>lasianthum</i>	1																					1
<i>Calluna vulgaris</i>									1								5	1	1	1	+	1
<i>Erica lusitanica</i>										+										+	+	1
<i>Erica scoparia</i>											+											
Companion species																						
<i>Cistus ladanifer</i>	1	+	4	4	+	+	+	+	1	1	1	1	1	+	+	+	+	+	1	1	4	1

Table 4. Cont.

	DEu1	DEu2	DEu3	DEu4	DEu5	DEu6*	DEu7	DEu8	DEu9	DEu10	DEu11	DEu12	DEu13	DEu14	DEu15	DEu16	DEu17	DEu18	DEu19	DEu20	DEu21	DEu22
<i>Rosmarinus officinalis</i>	+											1	1	1	+	1	+	1	4	4	4	
<i>Lavandula stoechas</i> subsp. <i>sampaioana</i>																		+	1			
<i>Briza maxima</i>															1	+	1	+				
<i>Tolpis barbata</i>																1						
<i>Petrorhagia prolifera</i>																	1	+	+			
<i>Cistus salvifolius</i>																				+	+	+
<i>Asphodelus albus</i>	+					+															+	
<i>Arrhenatherum bulbosum</i> subsp. <i>elatius</i>	+																					
<i>Jasione montana</i>		+				+																
<i>Ranunculus gramineus</i>			+																			
<i>Xolanta guttata</i>			+																			
<i>Jasione crispa</i> subsp. <i>tomentosa</i>				+																		
<i>Lavandula pedunculata</i>					+																	
<i>Cytinus hypocistis</i>																						
<i>Scirpus holoschoenus</i>																						
<i>Salix atrocinerea</i>																						
<i>Genista hirsuta</i>																						
<i>Asphodelus microcarpus</i>																						
<i>Osyris alba</i>																						
<i>Gladiolus illyricus</i>																						
<i>Thymelaea villosa</i>																				+		+
<i>Adenocarpus telonensis</i>																						
<i>Dactylis hispanica</i>																						
<i>Daphne gnidium</i>																						
<i>Arbutus unedo</i>																						
<i>Erica arborea</i>																						
<i>Phillyrea angustifolia</i>	+																					

Typus*.

4. Discussion

The study of heathlands dominated by *Erica australis* subsp. *australis*, subsp. *aragonensis*, *E. umbellata* and *Cistus populifolius* contributes eight different vegetation communities. Until now the heathlands in Luso-Extremaduran territories have been included in the known associations [4,5]. However, our recent studies in the province of Ciudad Real reveal floristic differences to the associations described previously. In the taxonomic study, [22,25] do not recognise *E. australis* L. subsp. *aragonensis* (Willk.) Cout., although [5,23] recognise the *aragonensis* variety.

Species in the genus *Erica* are always found in sub-humid–humid environments and appear in the dynamics of various types of *Quercus* from temperate and Mediterranean European areas [27–29], but always in acid-neutral pH substrates, never in *Quercus* forests on basic substrates [30]. In the degradation of these scrubs, either by human action or climate change [31–34]. They are replaced by species of the genus *Cistus* [24]; *Cistus ladanifer* is the species that replaces *Erica australis* and *Erica umbellata* in the studied territory.

The association TCp is a shrubland dominated by *C. populifolius* and is a cold vicariant community of PCp described by [4] for the more oceanic Extremaduran territories. The new association occupies the easternmost areas of the Mariánica mountain range, which receives the influence of the climate of the Manchegan plateau and comprises differential species such as *C. laurifolius* and the endemism *Teucrium oxylepis* subsp. *marianum*. These communities cannot therefore be included in the association PCp. [5] Lecto typifies the association HEu described by Rivas Goday in the province of Cáceres. The floristic and ecological differences make it inadvisable to include the communities dominated by *E. umbellata* from the eastern Mariánica range in this association, HEu, as they grow on extremely nitrogen-poor distric lithosols that allow the presence of *Drosophyllum lusitanicum*; the new association is therefore a soil vicariant of HEu. This same author describes the association HEa for the provinces of Segovia and Guadalajara, and gives the typus in the locality of Puerto de San Benito (Guadalajara) at an altitude of 1840 m. The floristic, bioclimatic and biogeographical differences lead us to propose the association HEau for the Luso-Extremaduran territories (Table 5).

Table 5. Synoptic table: TCp = Teucrio oxylepis-Cistetum populifolii; HEau = Halimio ocymoidis-Ericetum australis; ECp = Erico australis-Cistetum populifolii; PCp = Polygalo microphyllae-Cistetum populifolii; HEa = Halimio ocymoidis-Ericetum aragonensis; DEu = Drosophylo lusitanicae-Ericetum umbellatae; HEu = Halimio ocymoidis-Ericetum umbellatae.

	TCp	HEau	ECp	PCp	HEa	DEu	HEu	UEu
<i>Halimium ocymoides</i>	I	V	III	II	II	V	V	V
<i>Erica australis</i>	II	V	V			IV		II
<i>Cistus ladanifer</i>	IV	V	IV			V	II	II
<i>Erica umbellata</i>	I	II				V	V	V
<i>Tuberaria lignosa</i>	I	II				I	III	II
<i>Cistus crispus</i>	I	I	II					V
<i>Lavandula luisieri</i>	II	III	III			I		V
<i>Erica scoparia</i>	I	I		I		I	I	I
<i>Cistus salvifolius</i>	II	I				I		IV
<i>Quercus rotundifolia</i>	I	I	II	II			II	I
<i>Calluna vulgaris</i>	I	II		II	V	II	IV	
<i>Genista tridentata</i>	I	I	III	II	I	II	V	
<i>Polygala microphylla</i>	I	I		IV			I	
<i>Halimium umbellatum</i> subsp. <i>viscosum</i>	I		II		IV		I	
<i>Quercus pyrenaica</i>	I	II			I		I	
<i>Erica arborea</i>	III	II		IV	I	I		
<i>Cistus populifolius</i>	V	V	V	V		I		
<i>Arbutus unedo</i>	III	II	III	IV		I		
<i>Phillyrea angustifolia</i>	III	III	IV	III		I		
<i>Daphne gnidium</i>	II	I		II		I		
<i>Rosmarinus officinalis</i>	II	III	I			II		

Table 5. Cont.

	TCp	HEau	ECp	PCp	HEa	DEu	HEu	UEu
<i>Asphodelus albus</i>	I	I				I		
<i>Arrhenatherum elatius</i> subsp <i>bulbosum</i>	I					I		
<i>Ranunculus gramineus</i>	I					I		
<i>Adenocarpus telonensis</i>	I			III		I		
<i>Lavandula stoechas</i> subsp <i>sampaioana</i>	I			III		I		
<i>Cistus laurifolius</i>	I			II	I			
<i>Quercus suber</i>	II	I	IV	III				
<i>Pistacia terebinthus</i>	I	I	I	IV				
<i>Viburnum tinus</i>	I		I	III				
<i>Dactylis lusitanica</i>	I		II	I				
<i>Lonicera periclymenum</i>	I		I	III				
<i>Paeonia broteroi</i>	I		I	I				
<i>Rubia peregrina</i>	I		I	II				
<i>Lonicera implexa</i>	I			II				
<i>Doronicum plantagineum</i>	I			II				
<i>Cytisus grandiflorus</i>	I			I				
<i>Cephalanthera longifolia</i>	I			II				
<i>Arenaria montana</i>	I			I				
<i>Tamus communis</i>	I			III				
<i>Teucrium fruticans</i>	I			II				
<i>Magydaris panacifolia</i>	I			II				
<i>Astragalus lusitanicus</i>	I			III				
<i>Conopodium mariannum</i>	I			II				
<i>Jasminum fruticans</i>	I			I				
<i>Thapsia villosa</i>	I							
<i>Cytisus scoparius</i> subsp <i>bourgaei</i>	II	I	IV					
<i>Cardamine hirsuta</i>	I	I	IV					
<i>Helichrysum stoechas</i>	I		I					
<i>Srophularia scorodonia</i>	I		I					
<i>Origanum vulgare</i> subsp <i>virens</i>	I		I					
<i>Cistus albidus</i>	I		I					
<i>Quercus broteroii</i>	III	II						
<i>Thymus mastichina</i>	I	I						
<i>Juniperus oxycedrus</i> subsp <i>badia</i>	I							
<i>Teucrium oxylepis</i> subsp <i>mariannum</i>	I							
<i>Sedum forsterianum</i>	I							
<i>Narcissus pallidulus</i>	I							
<i>Cistus laurifolius</i>	I							
<i>Geum sylvaticum</i>	I							
<i>Festuca rothmaleri</i>	I							
<i>Rubus ulmifolius</i>	I							
<i>Rosa pouzinii</i>	I							
<i>Epipactis helleborine</i>	I							
<i>Sorbus torminalis</i>	I							
<i>Tulipa sylvestris</i>	I							
<i>Dorycnium pentaphyllum</i>	I							
<i>Aristolochia pistolochia</i>	I							
<i>Piptatherum miliaceum</i>	I							
<i>Helminthotheca comosa</i>	I							
<i>Thapsia garganica</i>	I							
<i>Leucanthemum sylvaticum</i>	I							
<i>Iberis linifolia</i>	I							
<i>Centaurea melitensis</i>	I							
<i>Campanula rapunculus</i>	I							
<i>Iris planifolia</i>	I							
<i>Linum suffruticosum</i>	I							
<i>Cistus monspeliensis</i>	I							
<i>Sanguisorba verrucosa</i>	I							

Table 5. Cont.

	TCp	HEau	ECp	PCp	HEa	DEu	HEu	UEu
<i>Teucrium haenseleri</i>		I		I			I	
<i>Osyris alba</i>		I		I		I		
<i>Narcissus triandrus</i>		I				I		
<i>Scirpus holoschoenus</i>		I				I		
<i>Cytisus multiflorus</i>		I		I				
<i>Festuca elegans</i>		I						
<i>Ranunculus sardous</i>		I						
<i>Klasea integrifolia</i>		I						
<i>Arrhenatherum album</i>		I						
<i>Helictochloa marginata</i>		I						
<i>Salix atrocinerea</i>		I						
<i>Halimium lasianthum</i>			III					I
<i>Xolantha guttata</i>			IV			I	I	
<i>Pteridium aquilinum</i>		I		III				I
<i>Urginea maritima</i>			IV	IV			I	
<i>Sanguisorba hybrida</i>			I					I
<i>Phillyrea latifolia</i>			I	III				
<i>Asparagus acutifolius</i>			III	I				
<i>Senecio minutus</i>			I	II				
<i>Phlomis purpurea</i>			I					
<i>Genista polyanthos</i>			III					
<i>Smilax mauritanica</i>			III					
<i>Euphorbia nicaeensis</i>			II					
<i>Myrtus communis</i>			II					
<i>Pistacia lentiscus</i>			I					
<i>Bupleurum fruticosum</i>			I					
<i>Genista falcata</i>			I					
<i>Ruscus aculeatus</i>			I					
<i>Quercus canariensis</i>			I					
<i>Digitalis purpurea</i> subsp. <i>tomentosa</i>			I					
<i>Saxifraga granulata</i> subsp. <i>glaucescens</i>			I					
<i>Asplenium adiantum-nigrum</i>			I					
<i>Selaginella denticulata</i>			I					
<i>Lunularia cruciata</i>			I					
<i>Genista triacanthos</i>				I				V
<i>Thymelaea villosa</i>				V			I	II
<i>Drosophyllum lusitanicum</i>					III		I	
<i>Quercus lusitanica</i>					IV			
<i>Teucrium scorodonia</i>					III			
<i>Anthericum baeticum</i>					IV			
<i>Ornithogalum pyrenaicum</i>					IV			
<i>Silene psammitis</i>					IV			
<i>Thapsia nitida</i>					IV			
<i>Cistus psilosepalus</i>					II			
<i>Genista tournefortii</i>					II			
<i>Smilax aspera</i>					II			
<i>Pterocephalus papposus</i>					II			
<i>Halimium halimifolium</i>					I			
<i>Halimium atriplicifolium</i>					I			
<i>Quercus coccifera</i>					I			
<i>Blechnum spicant</i>					I			
<i>Agrostis castellana</i>					I		II	II
<i>Erica australis</i> subsp. <i>aragonensis</i>					V		I	
<i>Polytrichum piliferum</i>					II		II	
<i>Avenula sulcata</i>					III		I	
<i>Cladonia furcata</i>					I		I	
<i>Jasione montana</i>					I	I		

Table 5. Cont.

	TCp	HEau	ECp	PCp	HEa	DEu	HEu	UEu
<i>Luzula lactea</i>					V			
<i>Deschampsia flexuosa subsp iberica</i>					V			
<i>Genista pilosa</i>					III			
<i>Artostaphylos uva-ursi subsp crassifolia</i>					II			
<i>Sedum brevifolium</i>					II			
<i>Helianthemum apenninum</i>					II			
<i>Lotus corniculatus subsp carpetanus</i>					II			
<i>Vaccinium myrtillus</i>					I			
<i>Jasione laevis subsp carpetana</i>					I			
<i>Erica cinerea</i>					I			
<i>Cetraria islandica</i>					I			
<i>Fagus sylvatica</i>					I			
<i>Nardus stricta</i>					I			
<i>Hieracium pilosella</i>					I			
<i>Plantago radicata</i>					I			
<i>Festuca durandii</i>					I			
<i>Juniperus nana</i>					I			
<i>Thymus bracteatus</i>					I			
<i>Genista hirsuta</i>					I			
<i>Cytinus hypocistis</i>					I			
<i>Jasione crispa subsp tomentosa</i>					I			
<i>Lavandula pedunculata</i>					I			
<i>Erica lusitanica</i>					I			
<i>Asphodelus microcarpus</i>					I			
<i>Gladiolus illyricus</i>					I			
<i>Briza maxima</i>					I			
<i>Tolpis barbata</i>					I			
<i>Petrorhagia prolifera</i>					I			
<i>Dactylis hispanica</i>					I			
<i>Halimium alyssoides</i>					I			
<i>Cladonia verticillata</i>					I			
<i>Cladonia mediterranea</i>					I			
<i>Agrostis setacea</i>					I			
<i>Ulex minor</i>					I			
<i>Genista micrantha</i>					I			
<i>Agrostis truncatula</i>					I			
<i>Ulex parviflorus subsp eriocladus</i>								V
<i>Pterospartum tridentatum</i>								V
<i>Festuca ampla</i>								I

*Syntaxonomical Checklist**CALLUNO-ULICETEA* Br.-Bl. & Tüxen ex Klika & Hadac 1944*ULICETALIA MINORIS* Quantin 1935*Ericion umbellatae* Br.-Bl., P. Silva, Rozeira & Fontes 1952*Ericenion umbellatae* Rivas-Martínez 1979*Halimio ocymoidis-Ericetum umbellatae* Rivas Goday 1964*Ullici eriocladi-Ericetum umbellatae* Rivas-Martínez 1979*Drosophyllo lusitanicae-Ericetum umbellatae nova**Ericenion aragonensis* Rivas-Martínez 1979*Polygalo microphyllae-Cistetum populifolii* Rivas Goday 1964*Erico australis-Cistetum populifolii* Rivas Goday 1964*Halimio ocymoidis-Ericetum aragonensis* Rivas-Martínez 1979*Teucro mariani-Cistetum populifolii nova**Halimio ocymoidis-Ericetum australis nova*

5. Conclusions

This study reveals a set of new associations for scrub dominated by species of the genera *Erica* and *Cistus*. All these communities act as dynamic states of cork oak forests *Quercus suber*, *Q. broteroi*, *Q. pyrenaica*, *Q. marianica* and *Q. canariensis*, of which the forests of *Q. pyrenaica* and *Q. canariensis* are the most fragile, both because they are at the edge of the range and are threatened by climate change. It is therefore essential to protect these forest formations and their dynamic stages, constituted firstly by scrub of *Arbutus unedo* and secondly by heaths of *Erica* sp. and *C. populifolius*. Since all these plant communities are located in sub-humid or humid mountain environments, where a high rate of endemic, rare or threatened species thrive, protection measures must be put in place against fire and excessive livestock pressure. The territory dominated by these shrubs contains a high rate of endemisms, which is why their protection is necessary by subjecting these areas to sustainable development, with control of livestock.

Author Contributions: Conceptualization, E.C.; Data curation, J.C.P.F., S.d.R.G., C.J.P.G. and E.C.; Formal analysis, J.C.P.F. and E.C.; Investigation, E.C.; Methodology, E.C.; Project administration, E.C.; Resources, E.C.; Supervision, G.S. and E.C.; Validation, J.C.P.F., S.d.R.G., C.J.P.G. and E.C.; Visualization, M.R. and E.C.; Writing—original draft, E.C.; Writing—review & editing, J.C.P.F., S.d.R.G. and E.C. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: No new data were created or analyzed in this study. Data sharing is not applicable to this article.

Conflicts of Interest: The authors declare no conflict of interest.

Appendix A

Table A1. Geographical and bibliographical data of the Teucrio mariani-Cistetum populifolii association.

Inventories	Localities	Coord. X	Coord. Y	References
TCp1	Sierra de San Andrés.	30S0440046	4259889	
TCp2	Prox. Río Robledillo	30S0412374	4252579	
TCp3	Umbría sierra de Niebla	30S0380443	4266304	
TCp4	Umbría de Juán Rodríguez	30S0458460	4254905	
TCp5	Prox. Barranco de las Palomas	30S0440174	4259923	
TCp6	Monte Rosalejo			Cano 1988
TCp7	Hontanar de Flores a Quintana			Cano 1988
TCp8	Barranco de Valdecañas (30SUH9227)			Melendo 1998
TCp9	Brezorrubios (30SUH9226)			Melendo 1998
TCp10	Barranco del Valle de En medio (30SUH9128)			Melendo 1998
TCp11	Barranco del Valle de En medio (30SUH9128)			Melendo 1998
TCp12	Prox. Cortijos de Fimia (30SUH9126)			Melendo 1998
TCp13	Barranco de Valdeinfierro (30SUH9428)			Melendo 1998
TCp14	Barranco del Cuervo (30SUH9229)			Melendo 1998
TCp15	Barranco de los Gabilanes	30S0459282	4250323	
TCp16	Umbría de los Gabilanes	30S0459360	4250242	
TCp17	Puerta del Telmo	30S0459676	4250349	
TCp18	Collado de los Jardines	30S0457259	4249203	

Table A2. Geographical and bibliographical data of the *Halimio ocymoidis-Ericetum australis* association.

Inventories	Localities	Coord. X	Coord. Y	References
HEau1	Sierra Madrona	30S0405702	4250017	
HEau2	Valle de Valmayor	30S0394329	4250907	
HEau3	Cumbres sierra Niebla	30S0377945	4266753	
HEau4	Abulagoso	30S0387149	4257635	
HEau5	Sierra Navalmanzano	30S0392537	4254795	
HEau6	Sierra Canalizos	30S0362282	4297986	
HEau7	V. Pilones. Sierra de los Canalizos	30S0364121	4306199	
HEau8	Hontanar de Flores			Cano 1988
HEau9	Prox. Risquillo			Cano 1988
HEau10	Burcio del Pino. Sierra Quintana			Cano 1988
HEau11	Monte Rosalejo			Cano 1988
HEau12	Monte Rosalejo			Cano 1988

Table A3. Geographical and bibliographical data of the *Drosophyollo lusitanicae-Ericetum umbellatae* association.

Inventories	Localities	Coord. X	Coord. Y	References
DEu1	Sierra Madrona	30S0405425	4249829	
DEu2	Abulagoso	30S0385725	4258301	
DEu3	Puerto de Madrona	30S0406569	4251696	
DEu4	Subida desde Puerto Madrona a Cumbres de Madrona	30S0406107	4251824	
DEu5	Subida desde Puerto Madrona a Cumbres de Madrona	30S0406072	4251798	
DEu6	Subida desde Puerto Madrona a Cumbres de Madrona	30S0405918	4251663	
DEu7	Subida desde Puerto Madrona a Cumbres de Madrona	30S0405731	4251616	
DEu8	Prox. Cumbres de Madrona	30S044993	4251508	
DEu9	Sierra Canalizos	30S0362282	4297986	
DEu10	Finca la Tapiada. Sierra Canalizos	30S0344784	4305623	
DEu11	Ladera oeste de Sierra Quintana			Cano 1988
DEu12	Majada del Reloj. Monte Limones	30S0439899	4255085	
DEu13	Hontanar de Flores (Jaén)			Cano 1988
DEu14	Hontanar de Flores (Jaén)			Cano 1988
DEu15	Hontanar de Flores a Risquillo			Cano 1988
DEu15	Subida a Burcio del Pino. Sierra Quintana			Cano 1988
DEu17	Subida a Burcio del Pino. Sierra Quintana			Cano 1988
DEu18	Ladera oeste de Sierra Quintana			Cano 1988
DEu19	Brezorrubios. 30SUH9226			Melendo 1998
DEu20	Brezorrubios. 30SUH9226			Melendo 1998
DEu21	Brezorrubios. 30SUH9226			Melendo 1998
DEu22	Brezorrubios. 30SUH9226			Melendo 1998
HEau1	Sierra Madrona	30S0405702	4250017	
HEau2	Valle de Valmayor	30S0394329	4250907	
HEau3	Cumbres sierra Niebla	30S0377945	4266753	
HEau4	Abulagoso	30S0387149	4257635	
HEau5	Sierra Navalmanzano	30S0392537	4254795	
HEau6	Sierra Canalizos	30S0362282	4297986	
HEau7	V. Pilones. Sierra de los Canalizos	30S0364121	4306199	
HEau8	Hontanar de Flores			Cano 1988
HEau9	Prox. Risquillo			Cano 1988
HEau10	Burcio del Pino. Sierra Quintana			Cano 1988
HEau11	Monte Rosalejo			Cano 1988
HEau12	Monte Rosalejo			Cano 1988
TCP1	Sierra de San Andrés.	30S0440046	4259889	
TCP2	Prox. Río Robledillo	30S0412374	4252579	
TCP3	Umbría sierra de Niebla	30S0380443	4266304	
TCP4	Umbría de Juán Rodríguez	30S0458460	4254905	

Table A3. Cont

Inventories	Localities	Coord. X	Coord. Y	References
TCp5	Prox. Barranco de las Palomas	30S0440174	4259923	
TCp6	Monte Rosalejo			Cano 1988
TCp7	Hontanar de Flores a Quintana			Cano 1988
TCp8	Barranco de Valdecañas (30SUH9227)			Melendo 1998
TCp9	Brezorrubios (30SUH9226)			Melendo 1998
TCp10	Barranco del Valle de En medio (30SUH9128)			Melendo 1998
TCp11	Barranco del Valle de En medio (30SUH9128)			Melendo 1998
TCp12	Prox. Cortijos de Fimia (30SUH9126)			Melendo 1998
TCp13	Barranco de Valdeinfierro (30SUH9428)			Melendo 1998
TCp14	Barranco del Cuervo (30SUH9229)			Melendo 1998
TCp15	Barranco de los Gabilanes	30S0459282	4250323	
TCp16	Umbría de los Gabilanes	30S0459360	4250242	
TCp17	Puerta del Telmo	30S0459676	4250349	
TCp18	Collado de los Jardines	30S0457259	4249203	

References

- Vila-Viçosa, C.; Vázquez, F.M.; Mendes, P.; del Rio, S.; Musarella, C.; Cano-Ortiz, A.; Meireles, C. Syntaxonomic update on the relict groves of Mirbeck's oak (*Quercus canariensis* Willd. and *Q. mariatica* C. Vicioso) in southern Iberia. *Plant Biosyst.* **2015**, *149*, 512–526. [[CrossRef](#)]
- Quinto-Canas, R.; Mendes, P.; Cano-Ortiz, A.; Musarella, C.M.; Pinto-Gomes, C. Forest fringe communities of the southwestern Iberian Peninsula. *Rev. Chapingo Ser. Cienc. For. Y Ambiente* **2018**, *24*, 415–434. [[CrossRef](#)]
- Piñar Fuentes, J.C.; Cano-Ortiz, A.; Musarella, C.M.; Quinto Canas, R.; Pinto Gomes, C.J.; Spampinato, G.; del Río, S.; Cano, E. Bioclimatology, Structure, and Conservation Perspectives of *Quercus pyrenaica*, *Acer opalus* subsp. *Granatensis*, and *Corylus avellanea* Deciduous Forests on Mediterranean Bioclimate in the South-Central Part of the Iberian Peninsula. *Sustainability* **2019**, *11*, 6500. [[CrossRef](#)]
- Rivas Goday, S. *Vegetación y Florula de la Cuenca Extremeña del Guadiana*; Excma: Madrid, Spain; Diputación Provincial de Badajoz: Badajoz, Spain, 1964; pp. 1–777.
- Rivas Martínez, S. Brezales y jarales de Europa occidental (Revisión Fitosociológica de las Clases Calluno-Ulicetea y Cisto-Lavanduletea). *Lazaroa* **1979**, *1*, 5–127.
- Ruiz Tellez, T. Flora y Vegetación Vascular del Tramo Medio del valle del Tietar y el Campo Arañuelo. Ph.D. Thesis, Universidad de Salamanca, Salamanca, Spain, 1986; pp. 1–631.
- Cano, E. Estudio Fitosociológico de la Sierra de Quintana (sierra Morena, Jaén). Ph.D. Thesis, Universidad de Granada, Granada, Spain, 1988; pp. 1–465.
- Sánchez Pascual, N. Estudio Fitosociológico y Cartográfico de la Comarca de Despeñaperros (Jaén). Ph.D. Thesis, Universidad de Granada, Granada, Spain, 1994; pp. 1–465.
- Amor, A.; Ladero, M.; Valle, C.J. Flora y vegetación vascular de la comarca de la vera y laderas meridionales de la sierra de Tormantos (Cáceres, España). *Studia Bot.* **1993**, *11*, 11–207.
- Melendo Luque, M. Cartografía y Ordenación Vegetal de Sierra Morena: Parque Natural de las Sierras de Cardeña y Montoro (Córdoba). Ph.D. Thesis, Universidad de Jaén, Jaén, Spain, 1998; pp. 1–616.
- Rivas-Martínez, S.; Lousa, M.; Díaz, T.E.; Fernández González, F.; Costa, J.C. La vegetación del sur de Portugal (Sado, Alentejo y Algarve). *Itinera Geobot.* **1990**, *3*, 5–126.
- Rivas-Martínez, S.; Fernández González, F.; Loidi, J.; Lousa, M.; Penas, A. Syntaxonomical checklist of vascular plant communities of Spain and Portugal to association level. *Itinera Geobot.* **2001**, *14*, 5–341.
- Quinto Canas, R.J. Flora y Vegetación de la Serra de Caldeirao. Ph.D. Thesis, Universidad de Jaén, Jaén, Spain, 2014; pp. 1–692.
- Specht, R.L. *Heathlands and Related Shrublands. Descriptive Studies*. *Ecosystems of the World* 9A; Elsevier: Amsterdam, The Netherlands, 1979.
- Gil-López, M.J.; Copete, F.O. La herriza y otros brezales europeos: ¿sabemos lo que tenemos? *Almoraima* **2011**, *43*, 111–130.
- Loidi, J.; Biurrun, L.; Campos, J.A.; García-Mijangos, I.; Herrera, M. A Sbiogeographical análisis of the European Atlantic lowland heathlands. *J. Veg. Sci.* **2010**, *21*, 832–842. [[CrossRef](#)]
- Marañón, T.; Ajbilou, R.; Ojeda, F.; Arroyo, J. Biodiversity of woody species in oak woodlands of southern Spain and northern Morocco. *For. Ecol. Manag.* **1999**, *115*, 147–156. [[CrossRef](#)]
- Gorissen, I. Die Zwerpstrauch-heiden Europas-vom Atlantik to Caucasus and Ural. (Dwarf shrub heaths of Europe from Atlantic to Caucasus and Ural). In *Umschlaggestaltung Und Druck*, 1st ed.; NHBS Ltd: Totnes, UK, 2004; 182p, ISBN 9783000136856.
- Ojeda, F. 4030 Brezales secos europeos. In *VV.AA, Bases Ecológicas Preliminares para la Conservación de los tipos de Hábitat de Interés Comunitario en España*; Ministerio de Medio Ambiente, y Medio Rural y Marino: Madrid, Spain, 2009; 66p.
- Ojeda, F. *El Brezal Mediterráneo o Herriza; Investigación y Ciencia*; Madrid, Spain, 2003.

21. Quintana, J.R.; Cruz, A.; Fernández-González, F.; Moreno, J.M. Time of germination and establishment success after fire of three obligate seeders in a Mediterranean shrubland of central Spain. *J. Biogeogr.* **2004**, *31*, 241–249. [[CrossRef](#)]
22. Bayer, E. *Erica in Flora Ibérica IV*; Real Jardín Botánico, C.S.I.C.: Madrid, Spain, 1993; pp. 485–506.
23. Coutinho, A.X.P. *Flora de Portugal*; Bertrand: Lisboa, Portugal, 1939; pp. 1–933.
24. Van der Maarel, E. Transformation of cover-abundance values in phytosociology and its effects on community similarity. *Vegetatio* **1979**, *39*, 97–114.
25. Pérez Latorre, A.V. Ericaceae. In *Vascular de Andalucía Oriental IV*; Blanca, G., Cabezudo, B., Cueto, M., López, C.F., Torres, C.M., Eds.; Flora Consejería de Medio Ambiente: Sevilla, Spain, 2009; pp. 258–266.
26. Campos-Vargas, C.; Sanchez-Azofeifa, A.; Laakso, K.; Marzahn, P. Unmanned Aerial System and Machine Learning Techniques Help to Detect Dead Woody Components in a Tropical Dry Forest. *Forests* **2020**, *11*, 827. [[CrossRef](#)]
27. Cano-Ortiz, A.; Pinto Gomes, C.J.; Musarella, C.M.; Cano, E. Expansion of the *Juniperus* genus due to anthropic activity. In *Old-Growth Forest and Coniferous Forests*; Weber, R.P., Ed.; Nova Science Publishers: New York, NY, USA, 2015; pp. 55–65.
28. Musarella, C.M.; Cano-Ortiz, A.; Piñar Fuentes, J.C.; Navas-Urena, J.; Pinto Gomes, C.J.; Quinto-Canas, R.; Cano, E.; Spampinato, G. Similarity analysis between species of the genus *Quercus* L. (Fagaceae) in southern Italy based on the fractal dimension. *PhytoKeys* **2018**, *113*, 79–95. [[CrossRef](#)] [[PubMed](#)]
29. Piñar Fuentes, J.C.; Cano-Ortiz, A.; Musarella, C.M.; Pinto Gomes, C.J.; Spampinato, G.; Cano, E. Rupicolous habitats of interest for conservation in the central-southern Iberian Peninsula. *Plant Sociol.* **2017**, *54* (Suppl. 1), 29–42.
30. Fakhry, A.M.; El-Keblawy, A.; Shabana, H.A.; Gamal, I.E.; Shalouf, A. Microhabitats Affect Population Size and Plant Vigor of Three Critically Endangered Endemic Plants in Southern Sinai Mountains, Egypt. *Land* **2019**, *8*, 86. [[CrossRef](#)]
31. Nunes, L.J.R.; Raposo, M.A.M.; Meireles, C.I.R.; Pinto Gomes, C.J.; Ribeiro, N.M.C.A. Fire as a Selection Agent for the Dissemination of Invasive Species: Case Study on the Evolution of Forest Coverage. *Environments* **2020**, *7*, 57. [[CrossRef](#)]
32. Cano, E.; Musarella, C.M.; Cano-Ortiz, A.; Piñar Fuentes, J.C.; Rodríguez Torres, A.; Del Río González, S.; Pinto Gomes, C.J.; Quinto-Canas, R.; Spampinato, G. Geobotanical Study of the Microforests of *Juniperus oxycedrus* subsp. *badia* in the Central and Southern Iberian Peninsula. *Sustainability* **2019**, *11*, 1111. [[CrossRef](#)]
33. Mendes, P.; Meireles, C.; Vila-Viçosa, C.; Musarella, C.; Pinto-Gomes, C. Best management practices to face degraded territories occupied by *Cistus ladanifer* shrublands—Portugal case study. *Plant Biosyst.* **2015**, *149*, 494–502. [[CrossRef](#)]
34. Cano, E.; Musarella, C.M.; Cano-Ortiz, A.; Piñar, J.C.; Pinto Gomes, C.J.; Rodríguez Torres, A.; Spampinato, G. A phytosociological review of siliceous sedges in C-W Spain and their state of conservation based on diversity indices. *Plant Sociol.* **2017**, *54*, 5–14.