

Phenol Biodegradation and Bioelectricity Generation by a Native Bacterial Consortium Isolated from Petroleum Refinery Wastewater

Sara Shebl ¹, Nourhan N. Hussien ¹, Mohab H. Elsabrouty ¹, Sarah M. Osman ¹, Bassma H. Elwakil ^{2,*}, Doaa A. Ghareeb ³, Safaa M. Ali ⁴, Nevine Bahaa El Din Ghanem ¹, Yehia M. Youssef ⁵, Essam El Din A. Moussad ⁶ and Zakia A. Olama ¹

¹ Botany & Microbiology Department, Faculty of Science, Alexandria University, Alexandria 21500, Egypt

² Medical Laboratory Technology Department, Faculty of Applied Health Sciences Technology, Pharos University in Alexandria, Alexandria 21500, Egypt

³ Biochemistry Department, Faculty of Science, Alexandria University, Alexandria 21500, Egypt


⁴ Nucleic Acid Research Center, City of Scientific Research and Technology Applications, Alexandria 21934, Egypt

⁵ Chemical and Petrochemical Department, Faculty of Engineering and Technology, Arab Academy for Science, Technology and Maritime Transport, Alexandria 1129, Egypt

⁶ Zoology Department, Faculty of Science, Alexandria University, Alexandria 21500, Egypt

* Correspondence: bassma.hassan@pua.edu.eg

Supplementary data






Figure S1. Phenol detection and estimation using HPLC; medium before fermentation (a), after fermentation with bacterial consortium B (b).