

Article

Vermicomposting as a Sustainable Option for Managing Biomass of the Invasive Tree *Acacia dealbata* Link

Celestino Quintela-Sabaris ^{*}, Luís A. Mendes  and Jorge Domínguez 

Grupo de Ecología Animal (GEA), Universidade de Vigo, 36310 Vigo, Spain

* Correspondence: tino.quintela.sabaris@gmail.com

Abstract: The tree *Acacia dealbata* is native to Australia but has become invasive in many parts of the world thanks to its N-fixing capacity and to the allelopathic compounds present in its biomass. We conducted a pilot-scale study to assess the potential conversion of *A. dealbata* biomass by vermicomposting via the earthworm *Eisenia andrei*. The flowering aerial *A. dealbata* biomass was shredded and placed in a vermireactor under greenhouse conditions for 56 days. The vermicomposted material was sampled every two weeks to analyse its biological and chemical parameters. The phytotoxicity of the fresh *A. dealbata* material and vermicompost was assessed via an ecotoxicological test with *Lepidium sativum* seeds. The activity of the earthworms caused strong modifications of the properties of the processed material: the electric conductivity, basal respiration, and organic matter content were reduced, whereas the concentrations of other elements such as N, P, or Zn increased. The earthworm biomass increased steadily until day 42 and then decreased, probably due to the depletion of labile organic matter during the initial stages of vermicomposting. The fresh *A. dealbata* material reduced the germination and radicle elongation of *L. sativum*, whereas vermicompost showed the same values as control. The produced vermicompost was an organic fertiliser rich in N and was not phytotoxic. Vermicomposting provides an opportunity to create a new value chain for the control of the invasive tree *A. dealbata*.

Keywords: *Eisenia andrei*; epigeic earthworms; germination tests; organic fertiliser; phytotoxicity



Citation: Quintela-Sabaris, C.; Mendes, L.A.; Domínguez, J. Vermicomposting as a Sustainable Option for Managing Biomass of the Invasive Tree *Acacia dealbata* Link. *Sustainability* **2022**, *14*, 13828. <https://doi.org/10.3390/su142113828>

Academic Editors: Rajeev Pratap Singh, Mallavarapu Megharaj, Vaibhav Srivastava, Surindra Singh Suthar and Vinod Kumar Garg

Received: 6 September 2022

Accepted: 20 October 2022

Published: 25 October 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Acacia dealbata Link (Fabaceae) is native to Victoria, New South Wales, and Tasmania (Australia), where it can grow as a shrub or tree, reaching heights of up to 30 m [1]. *A. dealbata* is a fast-growing species with showy, scented flowers. It is able to develop in nutrient-poor soils thanks to its ability to fix atmospheric nitrogen [2]. During the nineteenth century, *A. dealbata* was introduced in Europe, India, South Africa, and South and North America for multiple purposes: ornamental use, wood and tannins production, dune stabilization, or for the revegetation of bare land [3–7]. After its introduction in new areas, *A. dealbata* has expanded its range and is currently considered an invasive weed in SW Europe (Spain, Portugal, France, and Italy), South Africa (Cape floristic region), and areas with a Mediterranean climate in Chile [3,8]. The invasive behaviour of *A. dealbata* has been explained by the presence of several traits: a tolerance for changing soil conditions (including a capacity for nitrogen fixation and high water consumption), the ability to take advantage of environmental disturbance, phenotypic plasticity, vegetative reproduction (by the presence of rhizomes), fire tolerance (with the capacity to resprout and germinate profusely after forest fires), and an allelopathic potential [5,6,9]. Several compounds with phytotoxic effect have been found in *A. dealbata*, such as alkaloids, cyanogenic glycosides, terpenes, flavonoids, and tannins [7]. Although these compounds are mainly present during the flowering period [10], some studies have indicated a phytotoxic effect throughout the entire phenological cycle [7,11]. In spite of this recognised phytotoxicity of *A. dealbata*'s biomass, recent studies indicate that allelopathy may be a secondary factor in the success

of this species, whereas modifications in microhabitat induced by *A. dealbata* after its colonisation of a site constitute the main driver affecting seedling establishment in the European non-native range [12].

In Galicia (NW Spain), *A. dealbata* has extensively invaded natural forests, abandoned arable land, and water courses, causing an important impact on the native flora and even impacting protected areas [5,9]. According to these impacts, this species has been included since 2013 in the Spanish Catalog of Invasive Alien Species [13]. The inclusion of a species in this Catalog entails, among other measures, a generic prohibition of the possession, transport, traffic, and trade of live specimens, their remains, or propagules that could survive or reproduce, as well as the preparation of a control and eradication program for those species considered priority risks [14].

Although some biological control measures have been proposed for *A. dealbata* and other *Acacia* species in Europe [3,5] and tested in South Africa [15], the main strategy used is the mechanical removal of the aerial parts, sometimes in combination with the use of herbicides to reduce resprouting [13,16]. However, this process is expensive and generates biomasses with potentially negative effects on the flora and soils due to the presence of allelopathic compounds.

Recently, some authors have proposed the creation of value chains from *A. dealbata*, either by its use as biofuel or as a source of valuable compounds and biochemicals, as a way to improve the control and eradication of this invasive species [17,18]. Here, we propose an alternative: the conversion of *A. dealbata* biomass into a fertiliser by vermicomposting.

Vermicomposting is the managed processing of organic waste by the synergistic action of earthworms and microbial communities [6]. The enzymes involved in the biochemical decomposition of organic matter are produced by microorganisms, and earthworms accelerate this decomposition by comminution and mixing of plant debris, which boosts the activity of bacteria and saprophytic fungi. Vermicomposting is based on the use of epigeic earthworms, which are litter dwellers that live in organic horizons [19]. In spite of the high diversity of terrestrial earthworms, with around 4000 species described to date [20], only four species are routinely used in vermicomposting facilities: *Eisenia andrei* (Bouché), *E. fetida* (Savigny), *Perionyx excavatus* (Perrier), and *Eudrilus eugeniae* (Kinberg) [21]. Previous studies have demonstrated the feasibility of vermicomposting for converting organic waste, such as sewage sludge, paper waste, urban residues, and animal waste (see [19,22] for reviews), into organic fertilisers. Vermicomposting has also been shown to be effective for processing plant wastes rich in phytotoxic compounds, such as grape marc, spent coffee grounds, and Scotch broom biomass [23–25].

In this research, we explored the potential of vermicomposting as a sustainable option for the management of *A. dealbata* biomass and its conversion into an organic fertiliser. We had two objectives: (i) to estimate whether the earthworm *E. andrei* was able to degrade the biomass of *A. dealbata*, rich in secondary compounds, and (ii) to assess the potential phytotoxic risk of the obtained vermicompost. For this purpose, we conducted a pilot-scale vermicomposting trial and combined the analysis of biochemical parameters throughout the process with an ecotoxicological bioassay of the phytotoxicity of the final product.

2. Materials and Methods

2.1. Plant Material

Several young (around 5 m in height) silver wattle trees (*Acacia dealbata* Link) were cut down in a forest near the University of Vigo (NW Spain) in spring, when the trees were flowering. Young branches, leaves, and flowers were manually collected from those fallen trees. The branches were chopped into smaller pieces of 3–6 cm, whereas the flowers and leaves were left intact. The fresh weight of the total sample was 120 kg.

2.2. Vermicomposting Setup

The collected material was processed in a rectangular, metal, pilot-scale vermireactor (length 4 m × width 1.5 m wide × height 1 m) held in a greenhouse without temperature

control. Before the material was added to the vermireactor, it contained a layer of vermicompost (12 cm height), which acted as a bed for the earthworms (*Eisenia andrei*). The initial earthworm density in the vermireactor was 344 ± 10 individuals per m^{-2} , with a mean biomass of $99.4 \pm 4.5 \text{ g m}^{-2}$. The vermicompost layer was covered by a plastic mesh (5 mm mesh size), and the tree material was added to the vermireactor in a layer (12 cm) on top of the plastic mesh. The plastic mesh allowed earthworms to move between the layers but prevented mixing of the processed material and the vermicompost bedding. It also simplified sampling of the processed material during vermicomposting.

2.3. Data Collection

The density and biomass of the earthworm population was determined periodically by collecting 10 samples (five from above and five from below the plastic mesh) of the material in the vermireactor with a core sampler (7.5 cm diameter and 12 cm height) at each sampling time. Sampling was conducted every 14 days during the vermicomposting trial and ended after 56 days. In total, 50 samples were collected. At the end of the experiment, a total of 21 kg (fresh weight) of vermicompost had been produced.

In order to estimate the evolution of the physico-chemical properties of the substrate along the vermicomposting process, we divided the surface of the vermireactor into five sections and removed two subsamples (10 g) at random from each section at the beginning of the experiment (day 0) and after 14, 28, 42, and 56 days of vermicomposting. The two subsamples from each section were bulked and stored in plastic bags at -80°C until analysis. In total, 25 samples of substrate were collected and analysed.

2.4. Physico-Chemical Analyses

The moisture content was assessed after drying the samples for 24 h at 105°C . The organic matter (OM) content was determined from the weight loss after combustion in a Carbolite CWF 1000 muffle furnace at 550°C for 5 h. The pH and the electrical conductivity (EC) were measured in aqueous extracts (1:10 w/v) with a Crison MicropH 2000 pH meter and a Crison CM35 conductivity meter, respectively. Samples were oven-dried at 60°C . Total C and N content were determined in dry samples in a Carlo Erba EA 1108 CHNS-O 1500 C/N analyser. The total P, K, Ca, Mg, S, Fe, Mn, B, and Mo concentrations were measured by inductively coupled plasma optical emission spectrometry (ICP-OES) in dry, ground samples that were subjected to microwave-assisted acid digestion. The cellulose, hemicellulose, and lignin content were assessed by detergent fibre methods [26] using the FibreBag System (Gerhardt, Königswinter, Germany).

2.5. Microbiological Analyses

Basal respiration was estimated from CO_2 production. Subsamples of the plant material and vermicompost (5 g fresh weight) were placed in 100 mL glass vessels, which were then hermetically sealed and incubated at 22°C for 6 h. The CO_2 produced from the sample was trapped in 0.02 M NaOH. CO_2 production was quantified by titration with 0.01 M HCl to a phenolphthalein endpoint, after addition of excess BaCl_2 . Microbial biomass carbon (C_{mic}) and microbial biomass nitrogen (N_{mic}) were determined by the fumigation–extraction method [27,28] with moist subsamples (5 g fresh weight).

2.6. Phytotoxicity Assay

A germination and radicle elongation test was performed with *Lepidium sativum* L. to estimate the potential phytotoxicity of *A. dealbata* biomass (collected at $t = 0$) and the vermicompost (collected at the end of the experiment; $t = 56$). Briefly, 20 *L. sativum* seeds were placed on a filter paper in a sterile 9 mm diameter Petri dish into which 5 mL of one of the following solutions had already been added: distilled water (control, C), 5% w/v of *A. dealbata* fresh biomass (AD5), 10% w/v of *A. dealbata* fresh biomass (AD10), 5% w/v of vermicompost (VC5), or 10% w/v of vermicompost (VC10). To prepare the AD5, AD10, VC5, and VC10 solutions, the relevant material was suspended in water and filtered after

shaking for 1 h. Each treatment was prepared in triplicate, yielding a total of 15 Petri dishes and 300 seeds. After the seeds were sown, the Petri dishes were sealed with parafilm to prevent desiccation. The plates were incubated for 7 days at 20 °C in dark conditions. The plates were then collected and the number of germinated seeds in each plate was recorded. The radicle length of all germinated seeds was also recorded.

2.7. Statistical Analyses

The effect of time on the earthworm density and biomass in the vermireactor was estimated by a one-way ANOVA, followed by post hoc Tukey tests. Vermicompost chemical and microbiological parameters (except the content of fibers) were analysed by Principal Component Analysis (PCA) and a reduced number of Principal Components (PCs) that captured most of the variability in the data were extracted. The PCA was based on a correlation matrix, which is appropriate for data with different measurement scales. Scores of vermicompost samples on PCs with eigenvalues higher than 1 were retained and correlated with earthworm density and biomass by the Pearson correlation index. The phytotoxic effect of *A. dealbata* biomass and vermicompost on germination was assessed by a one-way ANOVA, and the differences in radicle length were assessed by a nested ANOVA, with the replicate nested within the treatment. In case of significant differences, Tukey post hoc tests were used to determine where the differences lay. Where necessary, the data were transformed to meet ANOVA requirements. ANOVAs, PCA, and linear correlations were computed with SPSS (v. 15.0, SPSS Inc., Chicago, IL, USA).

3. Results

3.1. Changes in the Properties of *A. dealbata* Vermicompost during Vermicomposting

The vermicompost produced by processing the *A. dealbata* biomass in the vermireactor for 56 days was slightly acidic (pH = 6.42), with total concentrations of N, P, and K of 39.2, 1.4, and 1.4 g kg^{−1}, respectively. The C/N ratio was 12.01, the concentrations of Cu and Zn were 27.8 and 270 mg kg^{−1}, and the concentrations of lignin, cellulose, and hemicellulose were 530.7, 188.1, and 57.4 g kg^{−1}, respectively. The values of the other chemical and microbiological parameters of *A. dealbata* vermicompost samples (averaged per sampling time) are included in Table A1 (Appendix A).

The application of a PCA analysis of the *A. dealbata* substrate and vermicompost allowed us to identify four Principal Components (PCs) that explained more than 90% of the total variance. The first principal component (PC1) alone explained 61.5% of the variance. The loadings (i.e., the correlations) of each parameter on the four PCs are presented in Table 1. Almost all parameters were correlated with PC1. The variables with positive correlations included the moisture content and the concentrations of several elements such as N, P, S, and others. By contrast, electrical conductivity, organic matter content, basal respiration, and the concentrations of C, B, and Cl were negatively correlated with PC1 (Table 1). Microbial C, pH, and potassium concentration were positively correlated with PC2. Microbial N was negatively correlated with PC3, whereas K was positively correlated with PC3. Only one parameter (Mo concentration) was correlated with PC4 (Table 1).

After plotting the *A. dealbata* and vermicompost samples on the PC1–PC2 axes, we observed that the samples were arranged along PC1 as a function of time (Figure 1): the PC1 scores for t_0 samples were below −1.5, the scores for t_{14} samples were around −1, the scores for t_{28} and t_{42} were around 0.5, and the scores for t_{56} samples were higher than 1. Considering these values and the parameters correlated with PC1, we interpreted that this axis provides information about the “vermicompost maturity”. Regarding PC2, the samples from t_0 and t_{56} had the lowest values, whereas the samples from the intermediate vermicomposting steps had the highest values. Given the correlation between microbial C biomass and PC2, we find that this axis indicated the “size of the bacterial community”. The samples tended towards lower values of PC3 over the sampling time (Figure A1, Appendix A), whereas the values of PC4 are similar for all sampling times except for the t_{28} samples, which yielded lower values (Figure A1, Appendix A).

Table 1. PCA analysis of *A. dealbata* vermicomposting process. The variation explained by four principal components (PC1 to PC4) and their eigenvalues are shown. The loadings of measured parameters on the four PCs are also shown. PCA was computed using a correlation matrix. Parameters strongly correlated with PCs (i.e., loadings > 0.55 or loadings < −0.55) are marked in **bold**.

	PC1	PC2	PC3	PC4
Explained variation (%)	61.5	15.3	8.1	5.4
Eigenvalue	12.9	3.2	1.7	1.1
Variable loadings				
pH	0.47	0.64	−0.45	0.03
Electrical Conductivity	−0.89	−0.25	0.31	−0.00
Moisture content	0.90	0.23	−0.14	−0.14
Organic Matter content	−0.87	0.36	−0.00	−0.00
N total concentration	0.89	0.31	−0.18	−0.20
C total concentration	−0.78	0.54	−0.04	0.14
B total concentration	−0.83	0.14	0.28	−0.13
Ca total concentration	0.83	0.45	0.25	−0.07
Cl total concentration	−0.95	−0.13	0.16	0.12
Cu total concentration	0.90	−0.37	0.06	0.15
Fe total concentration	0.93	−0.34	0.07	0.05
K total concentration	−0.07	0.73	0.60	0.15
Mg total concentration	0.71	0.55	0.35	−0.22
Mn total concentration	0.95	0.12	0.15	−0.10
Mo total concentration	0.42	−0.20	0.34	0.74
P total concentration	0.94	0.13	0.29	−0.00
S total concentration	0.95	0.08	0.25	−0.02
Zn total concentration	0.86	−0.39	0.05	0.23
Basal Respiration	−0.92	0.24	0.12	−0.02
Microbial biomass C	−0.07	0.73	−0.23	0.51
Microbial biomass N	0.24	0.16	−0.58	0.24




Figure 1. PCA results. *A. dealbata* samples collected at different times during vermicomposting are plotted in the space determined by the first and second principal components (PCs). Each empty dot represents one sample. Lines connect each sample with the group centroid. Different colours indicate different sampling times.

3.2. Changes in Earthworm Population

The population of earthworms in the vermireactor varied significantly throughout the experiment (Table 2). The total earthworm density and biomass increased steadily

from the beginning of the experiment. The maximum density (877 ± 34 individuals m^{-2}) was reached on day 28 and remained stationary until day 42. The maximum earthworm biomass (404.2 ± 15.2 g m^{-2}) was recorded on day 42. Both parameters then decreased until day 56 (Table 2).

Table 2. Variation in earthworm population during the vermicomposting of *A. dealbata*. The results of ANOVA (F and p-values) for the effects of the factor time are shown. The average \pm standard deviations of each parameter for each sampling time are also shown. Different superscript letters indicate significant differences between sampling times, according to Tukey post hoc tests.

	F _(4,20)	p-Val	Day 0	Day 14	Day 28	Day 42	Day 56
Earthworm density (individuals m^{-2})	254.9	7.5×10^{-17}	345 ^d \pm 23	434 ^c \pm 40	877 ^a \pm 34	889 ^a \pm 27	665 ^b \pm 46
Earthworm biomass (g m^{-2})	493.3	1.13×10^{-19}	99.4 ^e \pm 10	134.2 ^d \pm 15.6	371.9 ^b \pm 18.9	404.2 ^a \pm 15.2	208.2 ^c \pm 5.8

The earthworm population parameters (density and biomass) and principal component PC1 were highly significantly correlated. These parameters were also negatively correlated with PC4 (Figure 2).




Figure 2. Correlation between earthworm population and vermicompost parameters. (a) Scatter plot of total earthworm density (individuals m^{-2}) vs. PC1 to PC4 scores; (b) scatter plot of total earthworm biomass (g m^{-2} fw) vs. PC1 to PC4 scores. The Pearson's correlation index (r) is shown in those cases of significant linear correlations. * p-value < 0.05, *** p-value < 0.001.

3.3. Phytotoxic Effects before and after Vermicomposting

The results show that the fresh *A. dealbata* material was phytotoxic. Thus, the germination percentage of *L. sativum* seeds in treatments AD5 and AD10 was around 80%, whereas the percentage of germination in the control was around 97% (Table 3). The differences in root elongation between the control and treatments with *A. dealbata* were clearer: in the control, the average radicle length was slightly below 7 cm, whereas in AD5 and AD10 the average length was around 2 cm (Figure 3). By contrast, the treatments with mature vermicompost (VC5 and VC10) yielded similar germination percentages and radicle elongation rates as the control (Table 3 and Figure 3).

Table 3. Variation in the germination percentage (after 7 days) of *L. sativum* seeds treated with water (C), *A. dealbata* extracts (AD5 and AD10), and vermicompost extracts (VC5 and VC10). The results of ANOVA (F and *p*-values) regarding the effects of the factor *treatment* are shown. The average \pm standard deviation of germination percentage in each treatment are also shown. Different superscript letters indicate significant differences between treatments, according to Tukey post hoc tests.

	$F_{(4,10)}$	<i>p</i> -Val	C	AD5	AD10	VC5	VC10
Germination percentage (%)	4.63	0.023	96.7 ^a \pm 2.9	81.7 ^b \pm 5.8	78.3 ^b \pm 5.8	88.3 ^{a,b} \pm 7.6	93.3 ^{a,b} \pm 7.6




Figure 3. Boxplot of data obtained in the radicle elongation test with *L. sativum*. Treatments with extracts of fresh *A. dealbata* biomass are yellow and orange, whereas treatments with extracts of mature vermicompost are light and dark purple. Each box corresponds to the inter-quartile range, and the band inside the box denotes the median for radicle elongation in each treatment. Different letters indicate significant differences between treatments as indicated by Tukey post hoc test.

4. Discussion

We performed a pilot-scale study to assess the feasibility of vermicomposting as a way of managing and valorising the biomass of the invasive tree *A. dealbata*. The earthworm *Eisenia andrei* was able to grow and reproduce in this substrate, with a sustained increase in the number of individuals and earthworm biomass until day 28, followed by a plateau between days 28 and 42. After that point, the number of earthworms and their total biomass decreased until the end of the experiment. This demographic pattern is typical of populations growing in closed systems (i.e., systems without any matter exchange with any part outside the system), where populations grow steadily until the point when the consumption of all the available resources or the accumulation of wastes provokes a decrease in the population (e.g., [29]). Our results show a continuous reduction in hemicellulose (from 13.3% to 5.7%) and cellulose (from 21.7% to 18.8%) and an increase in lignin (from 38.8% to 53.2%) in the substrate during vermicomposting, which indicates the consumption of the more labile organic fractions by the earthworms and an enrichment in recalcitrant, less-palatable materials. Similarly, high concentrations of lignin and polyphenols in the leaf litter of *Acacia auriculiformis* were suggested to be responsible for the high earthworm mortality during the vermicomposting of this litter [30]. Other researchers found that the increase in N compounds and the decrease in the C:N ratio (depending on the residues, changes of C:N ratio from 40 to 20) negatively affected the growth and reproduction of *Eisenia fetida* during the vermicomposting of diverse organic residues [31]. However, it seems that the C:N ratio is not the limiting factor for *E. andrei*, because (i) the population in our experiment grew while the C:N ratio decreased from 19.9 to 12, and (ii) in another

vermicomposting study with *E. andrei*, the C:N ratio of the substrate stayed at values around 12 (11.5 at day 0 and 12.9 at day 42) and the earthworm population and biomass increased steadily with no signs of population stagnation [25].

In addition to the relative changes in the hemicellulose, cellulose, and lignin proportions mentioned above, the main trends observed during the vermicomposting of *A. dealbata* biomass included reductions in the EC; OM content; concentrations of B, C, and Cl; and basal respiration, and increases in moisture content and in the concentrations of N, Ca, Cu, Fe, Mg, Mn, P, S, and Zn. Similar processes have been documented during the vermicomposting of other substrates [23–25] and are mainly related to (i) rapid C mineralisation throughout the vermicomposting process (especially of the most labile fractions of the cell walls), which involves the concentration of other elements, and (ii) the reduction in the concentration of soluble ions due to leaching, precipitation into non-mobile salts, or immobilisation by microorganisms [24,32]. Other observed changes included an increase in pH and, especially, the microbial biomass of C during the intermediate stages of vermicomposting (day 14) and a decrease in these parameters until the final product. The observed variation in pH during vermicomposting (from acid values in the fresh material to more basic values in different vermicomposting stages) contrasts with the most common changes observed in the literature (progressive acidification during vermicomposting; see [33–35] and references therein), but is congruent with the changes observed during the aerobic composting of *A. dealbata* biomass [36]. While the acidification of the substrate during vermicomposting has been explained by the mineralisation of N and P and the formation of organic acids during the bioconversion of organic material [35], other authors have proposed that the increase in the pH in *A. dealbata* after composting was due to NH_3 formation and the degradation of organic acids [36].

The final product was stabilised organic matter, as indicated by the decrease in the microbial biomass of C and the basal respiration, as well as the decrease and stabilisation of the C:N ratio. Moreover, the vermicompost obtained met the EU-criteria for solid, nitrogen-rich organic fertilisers [37], i.e., solid and of organic origin; an N concentration above 2.5%; an organic carbon content higher than 15%; and concentrations of Cu and Zn below 300 and 800 mg kg^{-1} , respectively.

A. dealbata is known to possess secondary compounds such as phenolics, triterpenes, or alkaloids with allelopathic effects [7,10,11]. Thus, the use of *A. dealbata* as an organic fertiliser could be hampered by the presence of these phytotoxic compounds in the vermicompost. We conducted an ecotoxicological test with *Lepidium sativum*, a plant which has been widely used in this kind of study (e.g., [38]). Our results indicate that vermicomposting eliminated the phytotoxicity of the fresh material. Moreover, radicle elongation was a parameter with a higher sensitivity than the seed germination percentage. Similar reductions in phytotoxicity were obtained after the composting of the vermicompost of diverse organic residues such as manure, biosolids, and urban or food wastes [39–42]. Although we did not quantify the secondary compounds present in the fresh *A. dealbata* biomass and the vermicompost, we assume that our results indicated that the phytotoxic compounds have been degraded during the vermicomposting process. Accordingly, previous research found that vermicomposting with *E. andrei* totally degraded the polyphenols present in Scotch broom biomass [25] and reduced the concentrations of these phytotoxic compounds in grape marc [23]. Moreover, a recent study indicated that vermicompost derived from different types of organic waste can contain high amounts of different extracellular catalytic enzymes that are even capable of degrading organophosphorus pesticides [24].

5. Conclusions

We have performed a pilot-scale assay of vermicomposting as an alternative for the management of the invasive tree *Acacia dealbata*. We have found that *E. andrei* is able to grow and process the biomass of this tree and produce a stabilised material after 56 days. The obtained vermicompost meets the EU requirements for solid organic fertilisers. Our results also indicate that vermicomposting eliminated the phytotoxicity of *A. dealbata* biomass.

Some value chains have been proposed as a way to promote the control and eradication of the invasive tree *A. dealbata*. Overall, our findings indicate that vermicomposting with *E. andrei* is a feasible alternative for the conversion of this organic residue into a valuable fertiliser, thereby contributing to decreasing the impact of alien species on native biodiversity.

Author Contributions: Conceptualization, J.D.; formal analysis, C.Q.-S. and L.A.M.; investigation, J.D. and C.Q.-S.; resources, J.D.; data curation, C.Q.-S. and L.A.M.; writing—original draft preparation, C.Q.-S.; writing—review and editing, J.D., L.A.M. and C.Q.-S.; supervision, J.D.; project administration, J.D.; funding acquisition, J.D. All authors have read and agreed to the published version of the manuscript.

Funding: C.Q.-S. contract is funded by the European Union’s Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No 892814.

Data Availability Statement: Dataset of the paper “Vermicomposting as a sustainable option for the management of the biomass of the invasive tree *Acacia dealbata* Link”, openly available online at Zenodo: <https://doi.org/10.5281/zenodo.7024774> (accessed on 19 October 2022).

Acknowledgments: The technical support provided by Hugo Martínez-Cordeiro, Alberto da Silva, and Victor González Sales is gratefully acknowledged.

Conflicts of Interest: The authors declare no conflict of interest. The funders had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript; or in the decision to publish the results.

Appendix A

Table A1. Parameters of *A. dealbata* biomass at different points of the vermicomposting process. The values were averaged per sampling time ($n = 5$; except for fibers, $n = 3$).

Variable (Unit)	Time (Days)				
	0	14	28	42	56
Moisture (%)	64.23	72.08	78.63	80.7	79.56
pH	5.14	7.26	7.15	6.63	6.42
Electrical Conductivity (mS cm ⁻²)	752.4	370.8	259	263.6	211.8
Organic Matter (%)	96.64	96.38	92.89	93.41	89.12
Total C (g kg ⁻¹)	506.2	516.3	489.2	493.6	471.1
Total N (g kg ⁻¹)	25.5	33.3	40.8	41.3	39.2
C:N	19.9	15.49	11.99	11.96	12.01
Total B (mg kg ⁻¹)	62.39	43.56	37.22	28.19	15.71
Total Ca (g kg ⁻¹)	4.03	5.84	8.24	8.29	7.35
Total Cl (g kg ⁻¹)	6.86	4.22	0.89	0.98	0.56
Total Cu (mg kg ⁻¹)	2.7	3.6	12.3	13.6	27.8
Total Fe (g kg ⁻¹)	0.11	0.12	0.57	0.57	1.07
Total K (g kg ⁻¹)	1.82	1.96	1.78	2.23	1.38
Total Mg (g kg ⁻¹)	0.92	1.16	1.74	1.78	1.34
Total Mn (mg kg ⁻¹)	89.3	119.1	216.8	205.3	229.9
Total Mo (mg kg ⁻¹)	0.8	0.9	0.71	0.92	1.14
Total P (g kg ⁻¹)	0.61	0.76	1.23	1.39	1.42
Total S (g kg ⁻¹)	0.26	0.36	0.74	0.95	0.94
Total Zn (mg kg ⁻¹)	11.5	15.0	85.8	122.0	269.6
Basal respiration (μg CO ₂ g ⁻¹ dw h ⁻¹)	701.5	594.5	391.8	320.1	127.6
Microbial biomass C (g kg ⁻¹)	9.55	24.08	12.44	16.12	10.81
Microbial biomass N (g kg ⁻¹)	4.26	5.72	5.60	5.37	5.84
Lignin (g kg ⁻¹)	388.4	519.2	518.5	544.9	530.7
Cellulose (g kg ⁻¹)	217.1	219.9	209.1	192.1	188.1
Hemicellulose (g kg ⁻¹)	133.1	94.7	71.6	71.3	57.4




Figure A1. Plots of *A. dealbata* biomass and vermicompost samples: (a) PC1 vs. PC3 scores; (b) PC1 vs. PC4 scores. Each empty dot represents one sample. Lines connect each sample with the group centroid. Colours indicate different sampling times.

References

1. Flora of Australia. Available online: <https://profiles.ala.org.au/opus/foa/profile/Acaciadealbata> (accessed on 29 July 2022).
2. May, B.W.; Attiwill, P.M. Nitrogen-fixation by *Acacia dealbata* and changes in soil properties 5 years after mechanical disturbance or slash-burning following timber harvest. *For. Ecol. Manag.* **2003**, *181*, 339–355. [\[CrossRef\]](#)
3. Sheppard, A.W.; Shaw, R.H.; Sforza, R. Top 20 environmental weeds for classical biological control in Europe: A review of opportunities, regulations and other barriers to adoption. *Weed Res.* **2006**, *46*, 93–117. [\[CrossRef\]](#)
4. Gouws, A.J.; Shackleton, C.M. Abundance and correlates of the *Acacia dealbata* invasion in the northern Eastern Cape, South Africa. *For. Ecol. Manag.* **2019**, *432*, 455–466. [\[CrossRef\]](#)
5. Vieites-Blanco, C.; González-Prieto, S.J. Invasiveness, ecological impact and control of acacias in southwestern Europe—A review. *Web Ecol.* **2020**, *20*, 33–51. [\[CrossRef\]](#)
6. Lorenzo, P.; González, L.; Reigosa, M.J. The genus *Acacia* as invader: The characteristic case of *Acacia dealbata* Link in Europe. *Ann. For. Sci.* **2010**, *67*, 101. [\[CrossRef\]](#)
7. Aguilera, N.; Becerra, J.; Guedes, L.M.; Villaseñor-Parada, C.; González, L.; Hernández, V. Allelopathic effect of the invasive *Acacia dealbata* Link (Fabaceae) on two native plant species in south-central Chile. *Gayana Bot.* **2015**, *72*, 231–239. [\[CrossRef\]](#)
8. *Acacia dealbata* (*acacia bernier*). CABI Invasive Species Compendium. Species Datasheet. Available online: <https://www.cabi.org/isc/datasheet/2207> (accessed on 19 September 2022).
9. Lorenzo, P.; Pazos-Malvido, E.; Rubido-Bará, M.; Reigosa, M.J.; González, L. Invasion by the leguminous tree *Acacia dealbata* (Mimosaceae) reduces the native understorey plant species in different communities. *Aust. J. Bot.* **2012**, *60*, 669–675. [\[CrossRef\]](#)
10. Reigosa, M.J.; Carballeira, A. Phenological phase influences allelopathic potential of water leachates of *Acacia dealbata* obtained using simulated rain. *J. Allelochem. Interact.* **2016**, *2*, 9–15.
11. Aguilera, N.; Becerra, J.; Villaseñor-Parada, C.; Lorenzo, P.; González, L.; Hernández, V. Effects and identification of chemical compounds released from the invasive *Acacia dealbata* Link. *Chem. Ecol.* **2015**, *31*, 479–493. [\[CrossRef\]](#)
12. Lorenzo, P.; Rodríguez, J.; González, L.; Rodríguez-Echeverría, S. Changes in microhabitat, but not allelopathy, affect plant establishment after *Acacia dealbata* invasion. *J. Plant Ecol.* **2017**, *10*, 610–617.
13. *Acacia dealbata*: Catálogo Español de Especies Exóticas Invasoras. Available online: https://www.miteco.gob.es/es/biodiversidad/temas/conservacion-de-especies/acacia_dealbata_2013_tcm30-69798.pdf (accessed on 20 September 2022).
14. Spanish Government. Ley 42/2007, de 13 de diciembre, del Patrimonio Natural y de la Biodiversidad. Available online: <https://www.boe.es/buscar/pdf/2007/BOE-A-2007-21490-consolidado.pdf> (accessed on 20 September 2022).
15. Moran, V.C.; Hoffmann, J.H. Conservation of the fynbos biome in the Cape Floral Region: The role of biological control in the management of invasive alien trees. *BioControl* **2012**, *57*, 139–149. [\[CrossRef\]](#)
16. Flora Invasora de Galicia, Xunta de Galicia. Available online: https://cmatv.xunta.gal/seccion-feitos-vitais?content=Direccion_Xeral_Conservacion_Natureza/Biodiversidade/seccion.html&sub=Informacion_especies/&ui=Direccion_Xeral_Conservacion_Natureza/Dinamico/Especies/Invasoras/Flora/especie_0003.html (accessed on 25 August 2022).
17. Nunes, L.J.R.; Raposo, M.A.M.; Meireles, C.I.R.; Pinto Gomes, C.J.; Almeida Ribeiro, N.M.C. Control of Invasive Forest Species through the Creation of a Value Chain: *Acacia dealbata* Biomass Recovery. *Environments* **2020**, *7*, 39. [\[CrossRef\]](#)

18. López-Hortas, L.; Rodríguez-González, I.; Díaz-Reinoso, B.; Torres, M.D.; Moure, A.; Domínguez, H. Tools for a multiproduct biorefinery of *Acacia dealbata* biomass. *Ind. Crop. Prod.* **2021**, *169*, 113655. [\[CrossRef\]](#)
19. Domínguez, J.; Aira, M.; Gómez-Brandón, M. Vermicomposting: Earthworms Enhance the Work of Microbes. In *Microbes at Work: From Wastes to Resources*; Insam, H., Franke-Whittle, I., Goberna, M., Eds.; Springer: Berlin/Heidelberg, Germany, 2010; pp. 93–114.
20. Reynolds, J.W. Earthworms of the world. *Global. Divers.* **1994**, *4*, 11–16.
21. Domínguez, J. Earthworms and vermicomposting. In *Earthworms—The Ecological Engineers of Soil*; Ray, S., Ed.; Intech Open Science: Rijeka, Croatia, 2018; pp. 63–77.
22. Domínguez, J. State of the art and new perspectives on vermicomposting research. In *Earthworm Ecology*, 2nd ed.; Edwards, C.A., Ed.; CRC Press LLC: Boca Raton, FL, USA, 2004; pp. 401–424.
23. Domínguez, J.; Martínez-Cordeiro, H.; Álvarez-Casas, M.; Lores, M. Vermicomposting grape marc yields high quality organic biofertiliser and bioactive polyphenols. *Waste Manag. Res.* **2014**, *32*, 1235–1240. [\[CrossRef\]](#) [\[PubMed\]](#)
24. Sanchez-Hernandez, J.C.; Domínguez, J. Vermicompost derived from spent coffee grounds: Assessing the potential for enzymatic bioremediation. In *Handbook of Coffee Processing By-Products Sustainable Applications*; Galanakis, C.M., Ed.; Academic Press: London, UK, 2017; pp. 369–398.
25. Domínguez, J.; Gómez-Brandón, M.; Martínez-Cordeiro, H.; Lores, M. Bioconversion of Scotch broom into a high-quality organic fertiliser: Vermicomposting as a sustainable option. *Waste Manag. Res.* **2018**, *36*, 1092–1099. [\[CrossRef\]](#) [\[PubMed\]](#)
26. Goering, H.K.; van Soest, P.J. *Forage Fiber Analysis. Agricultural Handbook No. 379*; Agricultural Research Service, United States Department of Agriculture: Washington, DC, USA, 1970.
27. Vance, E.D.; Brookes, P.C.; Jenkinson, D.S. An extraction method for measuring soil microbial biomass C. *Soil Biol. Biochem.* **1987**, *19*, 703–707. [\[CrossRef\]](#)
28. Brookes, P.C.; Landman, A.; Pruden, G.; Jenkinson, D.S. Chloroform fumigation and the release of soil nitrogen: A rapid direct extraction method for measuring microbial biomass nitrogen in soil. *Soil Biol. Biochem.* **1985**, *17*, 837–842. [\[CrossRef\]](#)
29. Klein, D.R. The introduction, increase, and crash of reindeer on St. Matthew Island. *J. Wildl. Manag.* **1968**, *32*, 350–367. [\[CrossRef\]](#)
30. Ganesh, P.S.; Gajalakshmi, S.; Abbasi, S.A. Vermicomposting of the leaf litter of acacia (*Acacia auriculiformis*): Possible roles of reactor geometry, polyphenols, and lignin. *Bioresour. Technol.* **2009**, *100*, 1819–1827. [\[CrossRef\]](#)
31. Zhou, B.; Chen, Y.; Zhang, C.; Li, J.; Tang, H.; Liu, J.; Dai, J.; Tang, J. Earthworm biomass and population structure are negatively associated with changes in organic residue nitrogen concentration during vermicomposting. *Pedosphere* **2021**, *31*, 433–439. [\[CrossRef\]](#)
32. Fornes, F.; Mendoza-Hernández, D.; García-de-la-Fuente, R.; Abad, M.; Belda, R.M. Composting versus vermicomposting: A comparative study of organic matter evolution through straight and combined processes. *Bioresour. Technol.* **2012**, *118*, 296–305. [\[CrossRef\]](#) [\[PubMed\]](#)
33. Cai, L.; Gong, X.; Sun, X.; Li, S.; Yu, X. Comparison of chemical and microbiological changes during the aerobic composting and vermicomposting of green waste. *PLoS ONE* **2018**, *13*, e0207494. [\[CrossRef\]](#) [\[PubMed\]](#)
34. Domínguez, J.; Aira, M.; Kolbe, A.R.; Gómez-Brandón, M.; Pérez-Losada, M. Changes in the composition and function of bacterial communities during vermicomposting may explain beneficial properties of vermicompost. *Sci. Rep.* **2019**, *9*, 9657. [\[CrossRef\]](#)
35. Ndegwa, P.M.; Thompson, S.A.; Das, K.C. Effects of stocking density and feeding rate on vermicomposting of biosolids. *Bioresour. Technol.* **2000**, *71*, 5–12. [\[CrossRef\]](#)
36. Brito, L.M.; Mourao, I.; Coutinho, J.; Smith, S. Composting for management and resource recovery of invasive *Acacia* species. *Waste Manag. Res.* **2013**, *31*, 1125–1132. [\[CrossRef\]](#)
37. European Union. Consolidated Text: Regulation (EU) 2019/1009 of the European Parliament and of the Council of 5 June 2019 Laying Down Rules on the Making Available on the Market of EU Fertilising Products and Amending Regulations (EC) No 1069/2009 and (EC) No 1107/2009 and Repealing Regulation (EC) No 2003/2003 (Text with EEA Relevance) Text with EEA Relevance. Available online: <https://eur-lex.europa.eu/legal-content/EN/TXT/?uri=CELEX%3A02019R1009-20220716> (accessed on 29 August 2022).
38. Mañas, P.; de Heras, J.L. Phytotoxicity test applied to sewage sludge using *Lactuca sativa* L. and *Lepidium sativum* L. seeds. *Int. J. Environ. Sci. Technol.* **2018**, *15*, 273–280. [\[CrossRef\]](#)
39. Paré, T.; Gregorich, E.G.; Diné, H. Effects of stockpiled and composted manures on germination and initial growth of cress (*Lepidium sativum*). *Biol. Agric. Hortic.* **1997**, *14*, 1–11. [\[CrossRef\]](#)
40. Contreras-Ramos, S.M.; Escamilla-Silva, E.M.; Dendooven, L. Vermicomposting of biosolids with cow manure and oat straw. *Biol. Fertil. Soils* **2005**, *41*, 190–198. [\[CrossRef\]](#)
41. Majlessi, M.; Eslami, A.; Saleh, H.N.; Mirshafiean, S.; Babaii, S. Vermicomposting of food waste: Assessing the stability and maturity. *J. Environ. Health Sci. Eng.* **2012**, *9*, 25. [\[CrossRef\]](#)
42. Paradelo, R.; Moldes, A.B.; Rodríguez, M.; Barral, M.T. Relationship between heavy metals and phytotoxicity in composts. *Cienc. Tecnol. Aliment.* **2008**, *6*, 143–151. [\[CrossRef\]](#)