

Table S1. Primers used in qPCR.

Gene	Forward	Reverse
β -actin	GGCTGTATCCCCCTCCATCG	CCAGTTGGTAACAATGCCATGT
Hnnpa2b1	CAGGGTAGTTGAGCCAAAACG	TTCCAGACTGCCTATCGGTAA
Tnfa	CCCTCACACTCAGATCATCTTCT	GCTACGACGTGGGCTACAG
Il-6	GGGACTGATGCTGGTGACAA	TGCCATTGCACAACCTCTTTCT
Il-1 β	GAAATGCCACCTTTTGACAGTG	TGGATGCTCTCATCAGGACAG
Pdgfra	ATGAGAGTGAGATCGAAGGCA	CGGCAAGGTATGATGGCAGAG
Pdgfr β	CAAGAAGCGGCCATGAATCAG	CGGCCCTAGTGAGTTGTTGT
Ki67	ATCATTGACCGCTCCTTTAGGT	GCTCGCCTTGATGGTTCCT




Figure S1. The NF-kappaB activation is not altered in hnRNPA2B1-silenced macrophages. hnRNPA2B1, p65 and p-p65 protein levels in NC and A2B1 knockdown (siA2B1) LPS-treated macrophages. Results are shown as mean \pm SEM; ns, not significant; * $p < 0.05$.