

Table S1. Effect of maternal methyl donor or bisphenol A supplementation during gestation on body weight in newborn and weaning pigs.

	Treatment				<i>p</i> -value		
	CON	BPA	MET	BPA + MET	BPA	MET	BPA × MET
No. litters, <i>n</i>	13	12	13	12			
No. piglets, <i>n</i>	12.86 ± 0.89	12.62 ± 0.56	12.25 ± 1.05	14.55 ± 0.90	0.24	0.45	0.15
Birth weight, kg	1.24 ± 0.02 ^b	1.38 ± 0.03 ^a	1.34 ± 0.03 ^a	1.33 ± 0.03 ^a	0.01	0.66	<0.01
Weaning weight, kg	6.51 ± 0.35 ^c	7.06 ± 0.51 ^b	7.74 ± 0.24 ^{a,b}	7.14 ± 0.37 ^{a,b}	0.95	0.09	0.16

CON, control; BPA, bisphenol A; MET, methyl donor; BPA + MET, both bisphenol A and methyl donor supplementation in control diet. No. litters, *n*: the average number of litters in this group; No. piglets, *n*: the average number of piglets per litter in this group; Birth weight: the average weight of all newborn piglets in this group; Weaning weight: the average weight of all weaning piglets in this group. Within a row, means with different superscript letters are significantly different (*p* < 0.05).