

Urinary metabolic signatures detect recurrences in non-muscle invasive bladder cancer

Alba Loras, M. Carmen Martínez-Bisbal, Guillermo Quintás, Salvador Gil, Ramón Martínez-Máñez and José Luis Ruiz-Cerdá

Table S1. Confusion tables obtained from the evaluation of the predictive performance of PLS-DA models between pre-TUR (BC) and post-TUR (CTRL) or MONITOR samples (control) in the calibration (left) and validation (right) sets.

Class	Calibration (CV)		External Validation		
	BC	CTRL	BC	CTRL	CTRL+ MONITOR
Predicted as BC	39	7	19	1	9
Predicted as CTRL	9	14	3	7	37

Table S2. Patients and samples used in the calibration and validation set of PLS-DA model.

Patient	Calibration		Validation		
	BC	CTRL	BC	CTRL	MONITOR
1	3	0	--	--	--
2	1	0	--	--	--
3	3	1	--	--	--
4	3	1	--	--	--
5	1	0	--	--	--
6	2	1	--	--	--
7	2	2	--	--	--
8	1	1	--	--	--
9	1	1	--	--	--
10	1	0	--	--	--
11	1	1	--	--	--
12	2	1	--	--	--
13	2	1	--	--	--
14	1	0	--	--	--
15	1	0	--	--	--
16	1	1	--	--	--
17	3	1	--	--	--
18	2	1	--	--	--
19	2	1	--	--	--
20	3	2	--	--	--
21	2	1	--	--	--
22	--	--	2	0	10
23	--	--	8	5	6
24	6	2	2	0	5
25	2	2	0	0	5
26	--	--	1	2	9
27	2	0	--	--	--
28	--	--	9	1	3

