

Table S3. Genotype distribution of the IL-6 and IL-10 in DLBCL patients.

SNP ID	Model	Genotype	OR [^] (95% CI)	P-value
rs1800795 IL-6	Codominant	G/G	1	0.89
		C/G	0.91 (0.54-1.53)	
		C/C	1.26 (0.24-6.66)	
	Dominant	G/G	1	0.79
		C/G-C/C	0.93 (0.56-1.54)	
	Recessive	G/G-C/G	1	0.76
		C/C	1.30 (0.25-6.78)	
	Overdominant	G/G-C/C	1	0.71
		C/G	0.91 (0.54-1.52)	
rs1800796 IL-6	Codominant	G/G	1	0.66
		G/C	1.02 (0.55-1.89)	
		C/C	NA (0.00-NA)	
	Dominant	G/G	1	0.88
		G/C-C/C	1.05 (0.57-1.94)	
	Recessive	G/G-G/C	1	0.36
		C/C	NA (0.00-NA)	
	Overdominant	G/G-C/C	1	0.96
		G/C	1.02 (0.55-1.88)	
rs1800797 IL-6	Codominant	G/G	1	0.88
		A/G	0.90 (0.54-1.49)	
		A/A	1.24 (0.24-6.54)	
	Dominant	G/G	1	0.74
		A/G-A/A	0.92 (0.56-1.50)	
	Recessive	G/G-A/G	1	0.77
		A/A	1.28 (0.24-6.68)	
	Overdominant	G/G-A/A	1	0.67
		A/G	0.90 (0.54-1.48)	
rs1800871 IL-10	Codominant	G/G	1	0.76
		G/A	0.85 (0.53-1.37)	
		A/A	1.07 (0.44-2.63)	
	Dominant	G/G	1	0.58
		G/A-A/A	0.88 (0.56-1.39)	
	Recessive	G/G-G/A	1	0.75
		A/A	1.15 (0.48-2.75)	
	Overdominant	G/G-A/A	1	0.46
		G/A	0.84 (0.53-1.34)	
rs1800872 IL-10	Codominant	G/G	1	0.76
		G/T	0.93 (0.58-1.47)	
		T/T	0.97 (0.41-2.28))	
	Dominant	G/G	1	0.58
		G/T-T/T	0.93 (0.60-1.45)	
	Recessive	G/G-G/T	1	NA*
		T/T	1.00 (0.43-2.30)	
	Overdominant	G/G-T/T	1	0.76
		G/T	0.93 (0.59-1.46)	

rs1800890 IL-10	Codominant	A/A A/T T/T	1 0.85 (0.52-1.37) 0.78 (0.33-1.81)	0.73
	Dominant	A/A A/T-T/T	1 0.83 (0.53-1.32)	0.44
	Recessive	A/A-A/T T/T	1 0.84 (0.37-1.90)	0.67
	Overdominant	A/A-T/T A/T	1 0.88 (0.55-1.40)	0.58
rs1800896 IL-10	Codominant	T/T C/T C/C	1 0.89 (0.55-1.45) 1.18 (0.61-2.30)	0.67
	Dominant	T/T C/T-C/C	1 0.96 (0.60-1.51)	0.85
	Recessive	T/T-C/T C/C	1 1.26 (0.69-2.30)	0.44
	Overdominant	T/T-C/C C/T	1 0.85 (0.55-1.31)	0.45

^: Odd Ratio

*NA: not available.