

Supplementary information

Effect of surface chemistry and crystallographic parameters of TiO₂ anatase nanocrystals on photocatalytic degradation of bisphenol A

Gregor Žerjav ¹, Albin Pintar ^{2*}, Michael Ferentz ³, Miron Landau ⁴, Anat Haimovich ⁵, Amir Goldbourt ⁶ and Moti Herskowitz⁷

¹ Department for Environmental Sciences and Engineering, National Institute of Chemistry, Hajdrihova 19, SI-1001 Ljubljana, Slovenia; gregor.zerjav@ki.si

² Department for Environmental Sciences and Engineering, National Institute of Chemistry, Hajdrihova 19, SI-1001 Ljubljana, Slovenia; albin.pintar@ki.si

³ Blechner Center for Industrial Catalysis and Process Development, Department of Chemical Engineering, Ben-Gurion University of the Negev, Beer Sheva 84105, Israel; Ferenm@gmail.com

⁴ Blechner Center for Industrial Catalysis and Process Development, Department of Chemical Engineering, Ben-Gurion University of the Negev, Beer Sheva 84105, Israel; mlandau@bgu.ac.il

⁵ School of Chemistry, Raymond and Beverly Sackler Faculty of Exact Sciences, Tel Aviv University, Ramat Aviv 6997801, Israel; anathaim@post.tau.ac.il

⁶ School of Chemistry, Raymond and Beverly Sackler Faculty of Exact Sciences, Tel Aviv University, Ramat Aviv 6997801, Israel; amirgo@tauex.tau.ac.il

⁷ Blechner Center for Industrial Catalysis and Process Development, Department of Chemical Engineering, Ben-Gurion University of the Negev, Beer Sheva 84105, Israel; herskow@bgu.ac.il

* Correspondence: albin.pintar@ki.si; Tel.: +386 1 47 60 237




Fig. S1. PL calibration curve obtained by plotting the PL intensity measured at $\lambda=425$ nm as a function of 2-hydroxyterephthalic acid (TAOH) concentration.