

Electronic Supporting Information

Copper Metallopolymer Catalyst for the Electrocatalytic Hydrogen Evolution Reaction (HER)

Sait Elmas* ¹, Thomas J. Macdonald ², William Skinner ³, Mats Andersson* ¹, and Thomas Nann⁴

¹ Institute for NanoScale Science & Technology, Flinders University, Bedford Park SA 5042, Australia; sait.elmas@flinders.edu.au, mats.andersson@flinders.edu.au.

² Department of Chemistry, University College London, WC1H 0AJ, United Kingdom; tom.macdonald@ucl.ac.uk.

³ Future Industries Institute, University of South Australia Mawson Lakes Campus, Mawson Lakes SA 595, Australia; william.skinner@unisa.edu.au.

⁴ School of Mathematical and Physical Sciences, University of Newcastle, Callaghan NSW 2308, Australia; thomas.nann@newcastle.edu.au.

* Correspondence: sait.elmas@flinders.edu.au; Tel.: +61-8820-12684

Table of content

Figure S1: XPS survey spectrum of the polymer sample P_{OS}.....S2

Figure S2: XPS survey spectrum of the metallopolymer sample P_{OS}[Cu].....S2

Figure S3: (a) Gas chromatogram of the reference gas (200 ppm H₂) and b) gas chromatogram obtained from the head-space during the HER.....S3




Figure S 1: XPS survey spectrum of the polymer sample P_{OS}.




Figure S 2: XPS survey spectrum of the metallopolymer sample P_{OS}[Cu].




Figure S 3: (a) Gas chromatogram of the reference gas (200 ppm H_2) and (b) gas chromatogram obtained from the head-space during the HER.

Pictorial:

