

Supporting Information for

Emission Properties of Diblock Copolymers Composed of Poly(ethylene glycol) and Dense 1,2,3-Triazole Blocks

Yanqiong Yang¹, Asami Mori¹ and Akihito Hashidzume^{1,*}

¹ Department of Macromolecular Science, Graduate School of Science, Osaka University;

hashidzume@chem.sci.osaka-u.ac.jp

* Correspondence: hashidzume@chem.sci.osaka-u.ac.jp; Tel.: +81-6-6850-8174




Figure S1. SEC analysis of EG_m-P and EG_m-b-AP_n samples.




Figure S2. ^1H NMR spectra of EG18-*b*-AP n in $\text{DMSO-}d_6$ (500 MHz, 25 $^\circ\text{C}$).




Figure S3. ^1H NMR spectra of EG45-*b*-AP n in DMSO- d_6 (500 MHz, 25 °C).




Figure S4. Steady-state fluorescence spectra for 1.0 g L⁻¹ solutions of EG18-*b*-AP4 in acetonitrile (a) and water (b).




Figure S5. Steady-state fluorescence spectra for 1.0 g L⁻¹ solutions of EG45-*b*-AP6 in acetonitrile (a) and water (b).




Figure S6. DOSY spectrum for a 10.0 g L^{-1} solution of EG45-*b*-AP22 in $\text{DMF-}d_7$ at 298 K.