

Online Supplementary for

Anthropogenic CH₄ Emissions in the Yangtze River Delta Based on A “Top-Down” Method

Wenjing Huang ^{1,2}, Wei Xiao ^{1,2}, Mi Zhang ¹, Wei Wang ¹, Jingzheng Xu ³, Yongbo Hu ¹, Cheng Hu ^{1,4}, Shoudong Liu ¹ and Xuhui Lee ^{1,2,5,*}

* Correspondence: Xuhui.lee@yale.edu (X.L.)




Figure S1. Prevailing wind direction measured at the Taihu Lake Ecosystem Observatory of the Chinese Academy of Sciences (site id MLW) in the winter.




Figure S2. CH₄/CO₂ emissions ratio (ppm/ppm) estimated with different methods.

Table S1. Summary of CH₄:CO₂ emissions ratio (ppb:ppm) estimated with the atmospheric method found in the literature.

Location	Latitude and Longitude	Type	Time	CH ₄ /CO ₂	Reference
Boulder, Colorado		Surface, in situ	1985.12	7.6 ± 0.7	
Barrow, Alaska	71.32° N, 156.60° W	Surface, in situ	1986.03	21.2 ± 0.4	
Barrow, Alaska	71.32° N, 156.62° W	Surface, in situ	1986.04	13.5 ± 0.9	[1]
Arctic troposphere, lower stratosphere	-	Aircraft, flasks	1986.03–1986.04	17.5 ± 0.6	
Alert, North-West Territories, Canada	82.47° N, 62.50° W	Mountain, in situ	1986.04	20	[2]
Ocean Station "M"	66° N, 2° E	Surface, flasks	1983–1989, winter	9.3 ± 1.7 *	
Barrow, Alaska	71.32° N, 156.62° W	Surface, flasks	1989.03	11.0	[3]
Arctic troposphere, lower stratosphere	-	Aircraft, flasks	1989.03	13.5	
NOAA-Climate Monitoring and Diagnostics Lab Barrow Observatory	71.28° N, 156.78° W	Surface, in situ	1989–1990	9.14 ± 3.27	[4]
Alert, NWT, Canada	82.47° N, 62.50° W	Mountain, in situ	1992.01–1992.04, Black carbon > 100 ng/m ³	13.0 ± 2.2 *	[5]
Black Forest, Germany	48° N, 8° E	Mountain, in situ	1991–1995, winter (hourly & daily)	7.8 ± 1.0	[6]
Barrow, Alaska	71.32° N, 156.62° W	Surface, in situ	1986–1997, winter, midnight	12.8 ± 0.3	[7]
Tatra Mountains, southern Poland	49.17° N, 20.13° E	Mountain, in situ	1997, winter	10.7 ± 0.3	[8]
Pasadena, Los Angeles	34.2° N, 118.2° W	Aircraft, in situ	2007.082008.06	7.8 ± 0.8	[9]
Los Angeles	-	Aircraft, in situ	2008.06	6.74 ± 0.58	[10]
Los Angeles	-	Aircraft, in situ	2010.05–2010.06	6.55–6.7	[10,11]
Pasadena, Los Angeles	-	Surface, in situ	2012.02–2012.08	6.30 ± 0.01	
Pasadena, Los Angeles	34.14° N, 118.04° W	Surface, in situ	2011.09–2013.06	6.10 ± 0.10	[12]
Los Angeles	34.22° N, 118.06° W	Column (FTS)	2011.09–2013.10	6.40 ± 0.50	
Nanjing University of Information Science and Technology, Nanjing, China	32.20° N, 118.72° E	Building, in situ	2010, winter, midday	4.3 ± 0.7	[13]
Taihu Lake Ecosystem Observatory of the Chinese Academy of Sciences (site id MLW), Wuxi, China	31.42° N, 120.21° E	Surface, in situ	2012–2015, winter, midday	6.1 ± 0.5 *	This study

* The average among the period.

References

1. Conway, T.J.; Steele, L.P. Carbon dioxide and methane in the Arctic atmosphere. *J. Atmos. Chem.* **1989**, *9*, 81–99.
2. Trivett, N.B.A.; Worthy, D.E.J.; Brice, K.A. Surface measurements of carbon dioxide and methane at Alert during an Arctic haze event in April, 1986. *J. Atmos. Chem.* **1989**, *9*, 383–397, doi:10.1007/BF00052844.
3. Conway, T.J.; Steele, L.P.; Novelli, P.C. Correlations among atmospheric CO₂, CH₄, and CO in the Arctic, March 1989. *Atmos. Environ. Part A Gen. Top.* **1993**, *27*, 2881–2894, doi:10.1016/0960-1686(93)90319-T.
4. Jaffe, D.A.; Honrath, R.E.; Furness, D.; Conway, T.J.; Dlugokencky, E.; Steele, L.P. A determination of the CH₄, NO_x, and CO₂, emissions from the Prudhoe Bay, Alaska oil development. *J. Atmos. Chem.* **1995**, *20*, 213–227, doi:10.1007/BF00694494.
5. Worthy, D.E.J.; Trivett, N.B.A.; Hopper, J.F.; Bottenheim, J.W. Analysis of long-range transport events at Alert, Northwest Territories, during the Polar Sunrise Experiment. *J. Geophys. Res.* **1994**, *99*, 25329–25344, doi:10.1029/94JD01209.
6. Schmidt, M.; Graul, R.; Sartorius, H.; Levin, I. Carbon dioxide and methane in continental Europe: A climatology, and ²²²Radon-based emission estimates. *Tellus B* **1996**, *48*, 457–473, doi:10.1034/j.1600-0889.1994.t01-2-00002.x-i1.
7. Harris, J.M.; Dlugokencky, E.J.; Oltmans, S.J.; Tans, P.P.; Conway, T.J.; Novelli, P.C.; Thoning, K.W. An interpretation of trace gas correlations during Barrow, Alaska, winter dark periods, 1986–1997. *J. Geophys. Res.* **2000**, *105*, 17267–17278, doi:10.1029/2000JD900167.
8. Necki, J.; Schmidt, M.; Rozanski, K.; Zimnoch, M.; Korus, A.; Lasa, J.; Graul, R.; Levin, I. Six-year record of atmospheric carbon dioxide and methane at a high-altitude mountain site in Poland. *Tellus* **2010**, *55*, 94–104, doi:10.3402/tellusb.v55i2.16763.
9. Wunch, D.; Wennberg, P.O.; Toon, G.C.; Keppel-Aleks, G.; Yavin, Y.G. Emissions of greenhouse gases from a North American megacity. *Geophys. Res. Lett.* **2009**, *36*, 139–156, doi:10.1029/2009GL039825.
10. Wennberg, P.O.; Mui, W.; Wunch, D.; Kort, E.A.; Blake, D.R.; Atlas, E.L.; Santoni, G.W.; Wofsy, S.C.; Diskin, G.S.; Jeong, S.; et al. On the sources of methane to the Los Angeles atmosphere. *Environ. Sci. Technol.* **2012**, *46*, 9282–9289, doi:10.1021/es301138y.
11. Peischl, J.; Ryerson, T.B.; Brioude, J.; Aikin, K.C.; Andrews, A.E.; Atlas, E.; Blake, D.; Daube, B.C.; Gouw, J.A.de.; Dlugokencky, E.; et al. Quantifying sources of methane using light alkanes in the Los Angeles basin, California. *J. Geophys. Res.* **2013**, *118*, 4974–4990, doi:10.1002/jgrd.50413.ptional.
12. Wong, K.W.; Fu, D.; Pongetti, T.J.; Newman, S.; Kort, E.A.; Duren, R.; Hsu, Y.K.; Miller, C.E.; Yung, Y.L.; Sander, S.P. Mapping CH₄: CO₂ ratios in Los Angeles with CLARS-FTS from Mount Wilson, California. *Atmos. Chem. Phys.* **2015**, *15*, 241–252, doi:10.5194/acp-15-241-2015.
13. Shen, S.; Dong, Y.; Wei, X.; Liu, S.; Lee, X. Constraining anthropogenic CH₄ emissions in Nanjing and the Yangtze River Delta, China, using atmospheric CO₂ and CH₄ mixing ratios. *Adv. Atmos. Sci.* **2014**, *31*, 1343–1352, doi:10.1007/s00376-014-3231-3.



© 2019 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).