

Article

Monochrome Symmetric Subsets in Colorings of Finite Abelian Groups

Yuliya Zelenyuk

School of Mathematics, University of the Witwatersrand, Private Bag 3, Wits 2050, South Africa;
E-Mail: yuliya.zelenyuk@wits.ac.za

Received: 13 January 2011; in revised form: 17 March 2011 / Accepted: 21 March 2011 /

Published: 24 March 2011

Abstract: A subset S of a group G is symmetric if there is an element $g \in G$ such that $gS^{-1}g = S$. We study some Ramsey type functions for symmetric subsets in finite Abelian groups.

Keywords: finite Abelian group; symmetric subset; Ramsey functions

Classification: MSC 2010. Primary 05A20, 05C55, 20K01; Secondary 05D10.

1. Introduction

Let G be a finite group. Given an element $g \in G$, the *symmetry* on G with the *centre* g is the mapping

$$\eta_g : G \ni x \mapsto gx^{-1}g \in G$$

This is an old notion, which can be found in the book [1]. And it is a very natural one, since

$$\eta_g = \lambda_g \circ \iota \circ \lambda_g^{-1} = \rho_g \circ \iota \circ \rho_g^{-1}$$

where

$$\lambda_g : G \ni x \mapsto gx \in G, \rho_g : G \ni x \mapsto xg \in G, \text{ and } \iota : G \ni x \mapsto x^{-1} \in G$$

are the left translation, the right translation, and the inversion, respectively. Indeed, it follows from $\lambda_g(x) = gx$ that $\lambda_g^{-1}(gx) = x$, so $\lambda_g^{-1}(x) = g^{-1}x$. Consequently, $\lambda_g^{-1} = \lambda_{g^{-1}}$. Similarly, $\rho_g^{-1} = \rho_{g^{-1}}$. Then

$$\begin{aligned} \lambda_g \circ \iota \circ \lambda_g^{-1}(x) &= \lambda_g \circ \iota \circ \lambda_{g^{-1}}(x) = g(g^{-1}x)^{-1} = gx^{-1}g \text{ and} \\ \rho_g \circ \iota \circ \rho_g^{-1}(x) &= \rho_g \circ \iota \circ \rho_{g^{-1}}(x) = (xg^{-1})^{-1}g = gx^{-1}g \end{aligned}$$

A subset $S \subseteq G$ is *symmetric* if it is invariant with respect to some symmetry on G . Equivalently, S is *symmetric* if there exists an element $g \in G$ (centre of symmetry) such that $gS^{-1}g = S$.

Given $r \in \mathbb{N}$, an r -coloring of G is any mapping $\chi : G \rightarrow \{1, \dots, r\}$.

Definition 1.1 For every finite group and $r \in \mathbb{N}$, define the numbers $s_r(G)$ and $\sigma_r(G)$ as follows.

$s_r(G)$ is the greatest number of the form $\frac{k}{|G|}$, where $k \in \mathbb{N}$ such that for every r -coloring of G there exists a monochrome symmetric subset of cardinality k .

$\sigma_r(G)$ is the greatest number of the form $\frac{k}{|G|}$, where $k \in \mathbb{N}$ such that for every r -coloring χ of G there exists a subset $X \subseteq G$ of cardinality k and element g such that $\chi(x) = \chi(gx^{-1}g)$ for all $x \in X$.

It is easy to see that

$$s_r(G) \leq \frac{1}{r} + \frac{1}{|G|}, \quad \sigma_r(G) \leq 1, \quad s_r(G) \geq \frac{\sigma_r(G)}{r}$$

For every finite Abelian group G , $\sigma_r(G) \geq \frac{1}{r}$, and consequently, $s_r(G) \geq \frac{1}{r^2}$ [2]. In the non-Abelian case this inequality fails [3]. In this note we describe groups with $\sigma_r(G) = \frac{1}{r}$, $\sigma_r(G) = 1$, and $s_2(G) = \frac{1}{4}$. Since the journal [2] is not easy to access and it is in Ukrainian, we give here also a short proof of the inequality from [2].

2. The Inequality

In this section we prove the following theorem.

Theorem 2.1 Let G be a finite group of odd order or any finite Abelian group, and let $r \in \mathbb{N}$. Then

$$\sigma_r(G) \geq \frac{1}{r}$$

and consequently

$$s_r(G) \geq \frac{1}{r^2}$$

Let G be a finite group. For every r -coloring $\chi : G \rightarrow \{1, \dots, r\}$ and $g \in G$, let

$$S(\chi, g) = |\{x \in G : \chi(x) = \chi(gx^{-1}g)\}|$$

and let

$$\sigma(\chi) = \frac{1}{|G|} \max_{g \in G} S(\chi, g)$$

Then

$$\sigma_r(G) = \min_{\chi: G \rightarrow \{1, \dots, r\}} \sigma(\chi)$$

For every $a \in G$, let

$$\nu(a) = |\{x \in G : x^2 = a\}|$$

Lemma 2.2 For every $\chi : G \rightarrow \{1, \dots, r\}$,

$$\sum_{g \in G} S(\chi, g) = \sum_{i=1}^r \sum_{(x,y) \in A_i^2} \nu(yx^{-1})$$

where $A_i = \chi^{-1}(i)$.

Proof Computing in two ways the number of all triples $(g, x, y) \in G \times G \times G$ such that $gx^{-1}g = y$, we obtain

$$\sum_{g \in G} S(\chi, g) = \sum_{i=1}^r \sum_{(x,y) \in A_i^2} |\{g \in G : gx^{-1}g = y\}|$$

It remains to notice that

$$|\{g \in G : gx^{-1}g = y\}| = |\{g \in G : gx^{-1}gx^{-1} = yx^{-1}\}| = \nu(yx^{-1})$$

Proof of Theorem 2.1 Let $\chi : G \rightarrow \{1, \dots, r\}$ and let $A_i = \chi^{-1}(i)$. By Lemma 2.2

$$\sum_{g \in G} S(\chi, g) = \sum_{i=1}^r \sum_{(x,y) \in A_i^2} \nu(yx^{-1})$$

If G has odd order, then $\nu(yx^{-1}) = 1$ for any $x, y \in G$. Since the function $x_1^2 + \dots + x_r^2$, where $x_1 + \dots + x_r = C$, attains minimum when $x_1 = \dots = x_r = \frac{C}{r}$,

$$\sum_{g \in G} S(\chi, g) = \sum_{i=1}^r |A_i|^2 \geq \underbrace{\left(\frac{|G|}{r}\right)^2 + \dots + \left(\frac{|G|}{r}\right)^2}_r = \frac{|G|^2}{r}$$

If G is Abelian, then $\nu(yx^{-1}) > 0$ if and only if $yx^{-1} \in G^2 = \{g^2 : g \in G\}$ and in this case $\nu(yx^{-1}) = [G : G^2]$. Let C_j ($1 \leq j \leq k$) be cosets of G modulo G^2 , $C_{j,i} = C_j \cap A_i$. Then

$$\sum_{g \in G} S(\chi, g) = \sum_{i=1}^r \sum_{j=1}^k |C_{j,i}|^2 \cdot k \geq rk \left(\frac{|G|}{rk}\right)^2 \cdot k = \frac{|G|^2}{r}$$

Therefore, in each case, there exists an element $g \in G$ such that $S(\chi, g) \geq \frac{|G|}{r}$ and so $\sigma(\chi) \geq \frac{1}{r}$.

3. Finite Abelian Groups with $\sigma_r(G) = \frac{1}{r}$ and $\sigma_r(G) = 1$

In this section we describe finite Abelian groups with $\sigma_r(G) = \frac{1}{r}$ and $\sigma_r(G) = 1$.

Theorem 3.1 $\sigma_r(G) = \frac{1}{r}$ if and only if r divides $|2G|$.

Proof Define the subgroups $2G = \{2x : x \in G\}$ and $B(G) = \{x \in G : 2x = 0\}$. Denote $|2G| = m$ and $|B(G)| = k$. Obviously, $|G| = mk$.

Consider first the case when r does not divide m . Fix any r -coloring χ of a group G . Let C_j ($1 \leq j \leq k$) be cosets of G modulo $2G$, $C_{j,i} = C_j \cap \chi^{-1}(i)$. Then

$$\sum_{i=1}^r |C_{j,i}|^2 > r \left(\frac{m}{r}\right)^2 = \frac{m^2}{r}$$

Hence,

$$\sum_{g \in G} S(\chi, g) = k \sum_{j=1}^k \sum_{i=1}^r |C_{j,i}|^2 > k^2 \frac{m^2}{r} = \frac{|G|^2}{r}$$

Therefore, there exists an element $g \in G$ such that $S(\chi, g) > \frac{|G|}{r}$ and so $\sigma(\chi) > \frac{1}{r}$.

Now consider the case where r divides m . By Theorem 2.1, $\sigma_r(G) \geq \frac{1}{r}$, so it suffices to construct a coloring χ with $\sigma(\chi) = \frac{1}{r}$. Pick subgroup H of a group G such that $B(G) \subseteq H$ and $[G : H] = r$. Then $[2G : 2H] = r$. Define r -coloring χ of G as follows:

(1) every coset of G modulo $2H$ is monochrome;

(2) every r cosets of G modulo $2H$ which form a coset of G modulo $2G$ are colored in r different colors.

Then

$$\begin{aligned}\chi(x) = \chi(2g - x) &\Leftrightarrow x - (2g - x) \in 2H \\ &\Leftrightarrow 2(x - g) \in 2H \\ &\Leftrightarrow \exists h \in H : 2(x - g - h) = 0 \\ &\Leftrightarrow \exists h \in H : x - g - h \in B(G) \\ &\Leftrightarrow x - g \in H + B(G) = H \\ &\Leftrightarrow x \in g + H.\end{aligned}$$

So $S(\chi, g) = |H|$ for every $g \in G$. Therefore $\sigma(\chi) = \frac{|H|}{|G|} = \frac{1}{[G:H]} = \frac{1}{r}$.

Theorem 3.2 $\sigma_r(G) = 1$ if and only if one of the following cases holds:

- (1) $r = 1$;
- (2) $r = 2$ and G is a cyclic group of order either 3 or 5;
- (3) G is a Boolean group.

Proof Sufficiency is obvious. We need to prove Necessity. Assume on the contrary that neither of cases (1)–(3) holds.

Suppose first that $|G|$ is even. Then both subgroups $2G$ and the elementary Abelian 2-group $B(G)$ are different from G . Pick $a, b \in G$ such that $a + b \notin 2G$ and $a - b \notin B(G)$. Define $\chi : G \rightarrow \{1, 2\}$ by

$$\chi(x) = \begin{cases} 1 & \text{if } x \in \{a, b\} \\ 2 & \text{otherwise} \end{cases}$$

Let $g \in G$. Since $a + b \notin 2G$, $2g - a \neq b$. If $2g - a = a$, then $2g - b \neq b$, because $a - b \notin B(G)$. It follows that either $\chi(a) \neq \chi(2g - a)$ or $\chi(b) \neq \chi(2g - b)$, a contradiction.

Now suppose that $|G|$ is odd. Then $2G = G$. Since $|G| \geq 7$, we can choose distinct $a, b, c \in G$ such that for any distinct $g, x \in \{a, b, c\}$, $2g - x \notin \{a, b, c\}$.

To see this, pick any distinct $a, b \in G$. There is a unique $g \in G$ such that $b = 2g - a$. We then pick $c \in G \setminus \{a, b, g, 2a - b, 2b - a\}$.

Define $\chi : G \rightarrow \{1, 2\}$ by

$$\chi(x) = \begin{cases} 1 & \text{if } x \in \{a, b, c\} \\ 2 & \text{otherwise} \end{cases}$$

Let $g \in G$. If $g \notin \{a, b, c\}$ and $2g - a = b$, then $2g - c \notin \{a, b, c\}$. If $g \in \{a, b, c\}$, say $g = a$, then $2g - b \notin \{a, b, c\}$. It follows that there is $x \in \{a, b, c\}$ such that $\chi(x) \neq \chi(2g - x)$, again a contradiction.

Remark Theorem 3.2 describes finite Abelian groups where each r -coloring is symmetric. A coloring χ of G is *symmetric* if there exists $g \in G$ such that

$$\chi(gx^{-1}g) = \chi(x) \text{ for all } x \in G$$

Obviously, the number of all r -colorings of a group G of order n equals r^n . To find the number of all symmetric r -colorings of a group G is a quite complicated exercise involving Möbius inversion on the lattice of subgroups. Precise formula for the number of all symmetric r -colorings of finite Abelian group was established in [4], and the corresponding formula for every finite group has been found only recently [5] (for the quaternion group, see [6]).

4. Finite Abelian Groups with $s_2(G) = \frac{1}{4}$

In this section we prove the following.

Theorem 4.1 Let G be a finite Abelian group and $n \in \mathbb{N}$.

- (1) If G contains subgroup $\bigoplus_n \mathbb{Z}_4$, then $s_{2^n}(G) = \frac{1}{4^n}$;
- (2) If G does not contain subgroup $\mathbb{Z}_4$, then $s_2(G) > \frac{1}{4}$.

We first prove some auxiliary statements.

Lemma 4.2 $s_{2^n}(\bigoplus_n \mathbb{Z}_4) = \frac{1}{4^n}$.

Proof Define the coloring $\chi : \bigoplus_n \mathbb{Z}_4 \rightarrow \bigoplus_n \mathbb{Z}_2$ by

$$(\chi(x))_i = \begin{cases} 0 & \text{if } (x)_i \in \{0, 1\} \\ 1 & \text{if } (x)_i \in \{2, 3\} \end{cases}$$

Fix $g \in \bigoplus_n \mathbb{Z}_4$. If $\chi(x) = \chi(2g - x)$, then $(\chi(x))_i = (\chi(2g - x))_i$. It remains to notice that $s_2(\mathbb{Z}_4) = \frac{1}{4}$.

For every group G and a coloring χ , let $s(\chi)$ denote the cardinality of the largest monochrome symmetric subset of G divided by $|G|$.

Lemma 4.3 Let G be a finite group, let $f : G \rightarrow H$ be a surjective homomorphism and let χ be a coloring of H . Define coloring φ of G by $\varphi = \chi \circ f$. Then $s(\varphi) = s(\chi)$.

Proof Let S be a monochrome subset of G symmetric with respect to $g \in G$. By definition of φ it follows that $\varphi(x) = \varphi(gx^{-1}g)$ if and only if $\chi(f(x)) = \chi(f(g)f(x)^{-1}f(g))$. So, $f(S)$ is a monochrome subset of H symmetric with respect to $f(g)$. Since $|S| \leq |\ker f| \cdot |f(S)|$,

$$\frac{|S|}{|G|} \leq \frac{|\ker f| \cdot |f(S)|}{|G|} = \frac{|f(S)|}{|H|}$$

Thus $s(\varphi) \leq s(\chi)$.

Conversely, let S be a monochrome subset of H symmetric with respect to $h \in H$. Then $f^{-1}(S)$ is a monochrome subset of G symmetric with respect to any $g \in f^{-1}(h)$. Since $|f^{-1}(S)| = |\ker f| \cdot |f(S)|$,

$$\frac{|f^{-1}(S)|}{|G|} = \frac{|S|}{|H|}$$

Thus $s(\varphi) \geq s(\chi)$.

Corollary 4.4 Let G be a finite group and let H be a homomorphic image of G . Then $s_r(G) \leq s_r(H)$.

Proof of Theorem 4.1 (1) By Theorem 2.1, we have that $s_{2^n}(G) \geq \frac{1}{4^n}$. If G contains subgroup $\bigoplus_n \mathbb{Z}_4$, then there exists a homomorphism from G onto $\bigoplus_n \mathbb{Z}_4$. Thus, by Corollary 4.4, $s_{2^n}(G) \leq s_{2^n}(\bigoplus_n \mathbb{Z}_4)$. By Lemma 4.2, $s_{2^n}(\bigoplus_n \mathbb{Z}_4) = \frac{1}{4^n}$. Thus $s_{2^n}(G) = \frac{1}{4^n}$.

(2) Suppose that G does not contain $\mathbb{Z}_4$. Then $G = H \times B$ for some subgroup H of odd order and Boolean group B (which can be trivial). Let χ be an arbitrary 2-coloring of G . For every $b \in B$ define the coloring χ_b on H by $\chi_b(x) = \chi(x, b)$. Then

$$\sum_{h \in H} S(\chi, h) = \sum_{h \in H} \sum_{b \in B} S(\chi_b, h) = \sum_{b \in B} \sum_{h \in H} S(\chi_b, h)$$

Since H has odd order, we obtain that

$$\sum_{h \in H} S(\chi_b, h) > \frac{|H|^2}{2}$$

Then

$$\sum_{h \in H} S(\chi, h) = \sum_{h \in H} \sum_{b \in B} S(\chi_b, h) > |B| \cdot \frac{|H|^2}{2}$$

It follows that there exists $h \in H$ such that

$$S(\chi, h) > \frac{|B| \cdot |H|}{2} = \frac{|G|}{2}$$

Consequently,

$$\sigma(\chi) > \frac{1}{2}$$

and hence


$$s_2(G) > \frac{1}{4}$$

Theorem 4.1 implies the following two criteria.

Corollary 4.5 For every finite Abelian group G , $s_2(G) = \frac{1}{4}$ if and only if G contains subgroup $\mathbb{Z}_4$.

Corollary 4.6 $s_2(\mathbb{Z}_n) = \frac{1}{4}$ if and only if $4|n$.

Below we give the corresponding coloring.



We conclude the paper with the following table (Table 1).

Table 1. Ramsey functions $s_2(\mathbb{Z}_n)$ and $\sigma_2(\mathbb{Z}_n)$ for $n \leq 8$.					
n	k	$s_2(\mathbb{Z}_n) = \frac{k}{n}$	m	$\sigma_2(\mathbb{Z}_n) = \frac{m}{n}$	χ
1	1	1	1	1	
2	1	0.5	2	1	
3	2	0.66666666	3	1	
4	1	0.25	2	0.5	
5	3	0.6	5	1	
6	2	0.33333333	4	0.66666666	
7	3	0.42857142	5	0.71428571	
8	2	0.25	4	0.5	

Acknowledgements

The author is grateful to the referees for a careful reading of the paper and useful suggestions.

This work was supported by NRF grant IFR2008041600015, the John Knopfmacher Centre for Applicable Analysis and Number Theory, and the Friedel Sellschop Award.

References

1. Loos, O. *Symmetric Spaces*; Benjamin: New York, NY, USA, 1969.
2. Gryshko, Yu. Symmetric subsets and colorings of finite Abelian groups. *Visnyk Kyiv Univ., Ser. Fiz.-Mat.* **1999**, 3, 200–202. (Ukrainian)
3. Zelenyuk, Ye.; Zelenyuk, Yu. Transformations and colorings of groups. *Can. Math. Bull.* **2007**, 50, 632–636.
4. Gryshko, Yu. Symmetric colorings of regular polygons. *Ars Comb.* **2006**, 78, 277–281.
5. Zelenyuk, Yu. Symmetric colorings of finite groups. In *Proceedings of Groups St Andrews 2009*, Bath, UK, 1–15 August 2009.
6. Zelenyuk, Yu. The Number of Symmetric Colorings of the Quaternion Group. *Symmetry* **2010**, 2, 69–75.

© 2011 by the author; licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution license (<http://creativecommons.org/licenses/by/3.0/>.)