

Nestin Expression Is Associated with Relapses in Head and Neck Lesions

Mario Pérez-Sayáns ^{1,2*}, Cintia M Chamorro-Petronacci ², Fátima Baltazar ³, Fabio Ramoa Pires ⁴, Ángel Insua ⁵, Juan A Suárez-Quintanilla ⁶ and José M Suárez-Peñaranda ⁷

- ¹ Oral Medicine, Oral Surgery and Implantology Unit (MedOralRes), Faculty of Medicine and Dentistry, Universidade de Santiago de Compostela, 15782 Santiago de Compostela, Spain
 - ² Research Institute of Santiago de Compostela (IDIS), 15706 Santiago de Compostela, Spain; cintiamica.chamo@yahoo.es
 - ³ Life and Health Science Research Institute (ICVS), School of Medicine University of Minho, 4710-057 Braga, Portugal; fbaltazar@med.uminho.pt
 - ⁴ Post-graduation Program in Dentistry, Department of Oral Pathology, Estácio de Sá University, State University of Rio de Janeiro, Rio de Janeiro 22631-000, Brazil; ramoafop@yahoo.com
 - ⁵ Department of Periodontics and Oral Medicine, University of Michigan, Ann Arbor, MI 48109-1078, USA; angel_insua@yahoo.es
 - ⁶ Area of Human Anatomy and Embryology, Faculty of Medicine and Dentistry. University of Santiago de Compostela, 15782 Santiago de Compostela, Spain; juanantonisuarez.suarez@usc.es
 - ⁷ Pathological Anatomy Service, University Hospital Complex of Santiago (CHUS), Santiago de Compostela, 15706 Santiago de Compostela, Spain; jm.suarez.penaranda@gmail.com.
- * Correspondence: perezsayans@gmail.com; Tel.: +34 626233504

Citation: Pérez-Sayáns, M.; Chamorro-Petronacci, C.; Baltazar, F.; Pires, F.R.; Insua, Á.; Suárez-Quintanilla, J.A.; Suárez-Peñaranda, J.M. Nestin Expression Is Associated with Relapses in Head and Neck Lesions. *Diagnostics* **2021**, *11*, 583. <https://doi.org/10.3390/diagnostics11040583>

Academic Editor: Maciej Misiolek

Received: 10 February 2021

Accepted: 21 March 2021

Published: 24 March 2021

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).

Tumor subtypes

Mucoepidermoid carcinoma: 8 (9.08%)
Adenoid cystic carcinoma: 6 (6.81%)
Warthin tumor: 4 (4.54%)
Squamous cell carcinoma: 15 (17.04%)
Polyp: 13 (14.77%)
Inverted papilloma: 13 (14.77%)
Inverted papilloma with atypia: 1 (1.13%)
Pleomorphic adenoma: 7 (7.95%)
Parotid squamous cell carcinoma: 1 (1.13%)
Mucosecretory adenoma: 1 (1.13%)
Moderately differentiated adenocarcinoma: 3 (3.4%)
Well-differentiated adenocarcinoma: 1 (1.13%)
Well-differentiated mucinous adenocarcinoma, enteric type: 1 (1.13%)
Moderately differentiated mucinous adenocarcinoma: 2 (2.27%)
Undifferentiated carcinoma trabecular pattern: 1 (1.13%)
Undifferentiated sinonasal carcinoma: 1 (1.13%)
Poorly differentiated adeno squamous carcinoma: 2 (2.27%)
Epithelial-myoepithelial carcinoma: 1 (1.13%)
Large B-cell lymphoma: 1 (1.13%)
Hodgkin lymphoma: 2 (2.27%)
Salivary duct carcinoma: 2 (2.27%)
Small cell carcinoma: 1 (1.13%)
Acinic cell carcinoma: 1 (1.13%)