

Table S1. Mortality percentage (mean  $\pm$  SE) at day 14 of cockroach nymphs exposed for 15 min or 1 h to fresh or 1-month aged insecticide-based paints on various substrates.

<b>Insecticide</b>	<b>Treatment</b>	<b>Concrete</b>	<b>Metal</b>	<b>PVC</b>
0.7% alphacypermethrin	Control	8 $\pm$ 6	6 $\pm$ 2	8 $\pm$ 6
	Fresh-15 min	12 $\pm$ 7	10 $\pm$ 4	10 $\pm$ 5
	Fresh-1h	34 $\pm$ 7	20 $\pm$ 5	20 $\pm$ 4
	1 month aged-1h	54 $\pm$ 17	82 $\pm$ 6	50 $\pm$ 10
0.25% deltamethrin	Control	8 $\pm$ 6	6 $\pm$ 2	8 $\pm$ 6
	Fresh-15 min	8 $\pm$ 6	10 $\pm$ 5	14 $\pm$ 4
	Fresh-1h	16 $\pm$ 5	10 $\pm$ 4	18 $\pm$ 9
	1 month aged-1h	40 $\pm$ 14	54 $\pm$ 9	44 $\pm$ 11
0.5% transfluthrin	Control	8 $\pm$ 6	6 $\pm$ 2	8 $\pm$ 6
	Fresh-15 min	8 $\pm$ 6	10 $\pm$ 6	6 $\pm$ 4
	Fresh-1h	14 $\pm$ 9	12 $\pm$ 8	14 $\pm$ 9
	1 month aged-1h	14 $\pm$ 2	12 $\pm$ 8	2 $\pm$ 2