

Table S2: Determination of the SARS-CoV-2 replicative activity in the lungs of monoinfected and coinfecting with HAdV-5 hamsters by the formazan-based MTT assay.

Dilutions of lung homogenates	SARS-CoV-2	SARS-CoV-2/HAdV-5	HAdV-5/3 days/SARS-CoV-2	Mock-infected
Optical Density (OD <sub>570</sub> )				
10 <sup>0</sup>	0.123±0.024	0.137±0.029	0.144±0.035	1.230±0.215
10 <sup>-1</sup>	0.143±0.035	0.153±0.044	0.147±0.041	1.244±0.207
10 <sup>-2</sup>	0.135±0.041	0.129±0.037	0.157±0.045	1.233±0.217
10 <sup>-3</sup>	0.128±0.037	0.125±0.041	0.121±0.029	1.250±0.220
10 <sup>-4</sup>	0.155±0.046	0.167±0.044	0.141±0.035	1.227±0.195
10 <sup>-5</sup>	0.513±0.094	0.453±0.087	0.537±0.115	1.242±0.211
10 <sup>-6</sup>	0.836±0.144	0.723±0.127	0.870±0.163	1.239±0.223
10 <sup>-7</sup>	1.248±0.234	1.237±0.222	1.229±0.192	1.243±0.227
Virus Titer (lg TCID <sub>50</sub> /g)				
	6.43±0.36	6.60±0.44	6.27±0.32	<1.00

Data are the mean ±SEM of 2 independent MTT tests (3 replicates per point). lgTCID<sub>50</sub> values were determined according to the method of Kärber as modified [1].

1. Ashmarin I.P., Vorob'ev A.A. *Statistical Methods in the Microbiological Research*. State Press of Medical Literature; Leningrad, USSR: 1962. pp. 85–104.