

Supporting Information

Analysing Mouse Skin Cell Behaviour under a Non-Thermal kHz Plasma Jet

Andrea Jurov ^{1,2}, **Špela Kos** ³, **Nataša Hojnik** ¹, **Ivana Sremački** ⁴, **Anton Nikiforov** ⁴, **Christophe Leys** ⁴, **Gregor Serša** ³, and **Uroš Cvelbar** ^{1,*}

¹ Jožef Stefan Institute, Jamova cesta 39, 1000 Ljubljana, Slovenia; andrea.jurov@ijs.si, natasa.hojnik@ijs.si, uros.cvelbar@ijs.si

² Jožef Stefan International Postgraduate School, Jamova cesta 39, 1000 Ljubljana, Slovenia; andrea.jurov@ijs.si

³ Department of Experimental Oncology, Institute of Oncology, Zaloška cesta 2, 1000 Ljubljana, Slovenia; skos@onko-i.si, gserša@onko-i.si

⁴ Department of Applied Physics, Ghent University, Sint-Petersnieuwstraat 41, 9000 Gent, Belgium; ivana.sremacki@ugent.be, anton.nikiforov@ugent.be, christophe.leys@ugent.be

* Correspondence: uros.cvelbar@ijs.si

Supplementary Materials:




Figure 1. Spatially-resolved OES with intensity normalised to 1. ‘1. position’ is the exit of the glass tube and ‘2. position’ is the end of the plasma plume




Figure 2. Conductivity of PAM measured right after treatment

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



© 2021 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).