



Review

Sesquiterpenes and Sesquiterpene Derivatives from *Ferula*: Their Chemical Structures, Biosynthetic Pathways, and Biological Properties

Junchi Wang ^{1,†} , Qi Zheng ^{1,†}, Huaxiang Wang ¹, Leiling Shi ², Guoping Wang ², Yaqin Zhao ², Congzhao Fan ^{2,*} and Jianyong Si ^{1,*}

- ¹ The Key Laboratory of Bioactive Substances and Resources Utilization of Chinese Herbal Medicine, Ministry of Education, Institute of Medicinal Plant Development, Chinese Academy of Medical Sciences & Peking Union Medical College, Beijing 100193, China; jcwang@implad.ac.cn (J.W.); zhengqi@implad.ac.cn (Q.Z.); wanghuaxiang@implad.ac.cn (H.W.)
- ² Xinjiang Institute of Chinese Materia Medica and Ethnodrug, Urumqi 830002, China; shileiling@sina.com (L.S.); ping112_003@163.com (G.W.); xjzyq123@126.com (Y.Z.)
- * Correspondence: fcz_840701@163.com (C.F.); jysi@implad.ac.cn (J.S.)
- † These authors contributed equally to this work.

Abstract: *Ferula* is a genus of flowering plants known for its edible and medicinal properties. Since ancient times, many species of *Ferula* have been used in traditional medicine to treat various health issues across countries, such as digestive disorders, respiratory problems, and even as a remedy for headaches and toothaches. In addition, they are also used as a flavoring agent in various cuisines. As the main active ingredients in *Ferula*, sesquiterpenes and their derivatives, especially sesquiterpene coumarins, sesquiterpene phenylpropanoids, and sesquiterpene chromones, have attracted the attention of scientists due to the diversity of their chemical structures, as well as their extensive and promising biological properties, such as antioxidative, anti-inflammatory, antibacterial properties. However, there has not been a comprehensive review of sesquiterpenes and their derivatives from this plant. This review aims to provide an overview of the chemical structures, biosynthetic pathways, and biological properties of sesquiterpenes and sesquiterpene derivatives from *Ferula*, which may help guide future research directions and possible application methods for this valuable edible and medicinal plant.

Keywords: *Ferula*; sesquiterpene; biosynthetic pathway; antioxidative; antibacterial; anti-inflammatory



Citation: Wang, J.; Zheng, Q.; Wang, H.; Shi, L.; Wang, G.; Zhao, Y.; Fan, C.; Si, J. Sesquiterpenes and Sesquiterpene Derivatives from *Ferula*: Their Chemical Structures, Biosynthetic Pathways, and Biological Properties. *Antioxidants* **2024**, *13*, 7. <https://doi.org/10.3390/antiox13010007>

Academic Editors: Raquel Bridi and Christian Espinosa-Bustos

Received: 9 November 2023

Revised: 13 December 2023

Accepted: 15 December 2023

Published: 19 December 2023



Copyright: © 2023 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Ferula is a diverse genus of flowering plants belonging to the Apiaceae family, which primarily grow in dry and temperate regions of the Euro-Asian continent, surrounded by India and China in the east, the Canary Islands in the west, Central Europe in the north, and North Africa in the south [1]. This genus comprises about 180 recognized species [2], and it is renowned for its distinctive and often aromatic plants due to the presence of volatile essential oils and sulfide compounds. There are 94 species distributed in the erstwhile USSR, 32 species in Iran, 25 species in China, 19 species in the western Himalayas, 18 species in Turkey, 15 species in Pakistan, 4 species in Saudi Arabia, and 3 species in India [3]. Among these, 15 species are endemic to Iran, 9 species to Turkey, 7 species to China, and 1 species to Italy [4].

Since ancient times, different species of *Ferula* have been used in traditional medicine to treat various diseases across countries [5]. Asafoetida is an oleo–gum–resin obtained from the stems of *Ferula* plants, and in many parts of the world, it is used as a traditional medicine and as a flavoring agent in various cuisines [6]; its dual role in cuisine and traditional medicine is notable in several cultures. Asafoetida is commonly used in Indian,

Iranian, and some Middle Eastern dishes, and it is known for enhancing the flavor of dishes, especially in vegetarian recipes. It is a common ingredient in spice blends and seasoning for lentils, vegetables, and rice dishes [7,8]. In the folk medicine of Russia, Iran, China, Turkey, Pakistan, and India, asafoetida is often called “Asafetida”, “Rechina fena (Zaz)”, “A-wei”, “Setan bokosu (Seytan tersi)”, “Anjadana (Kama, Anguza)”, and “Hengu (Hing, Hingu, Ingu, etc.)”, respectively [9]. It is traditionally used to treat various health issues, such as digestive disorders, respiratory problems, and even as a remedy for headaches and toothaches [10–12]. The digestive-stimulating effect of Asafoetida is the most common beneficial physiological effect. In addition, other parts of some *Ferula* species also have edible and medicinal values. For instance, some nomadic peoples in central Iran use fried aerial parts of *F. assafoetida* and some seasonings as carminative foods. In Brazil, a hot-water extract from the dried stems and leaves of *F. assafoetida* is used as an aphrodisiac that is orally taken for the treatment of erectile dysfunction [13]. People in Pakistan extensively use the *F. narthex* Boiss herb for the treatment of coughs, fever, scorpion stings, hysteria, gastric dysfunction, constipation, habitual miscarriage, and toothache [14]. In Saudi Arabia, the rhizomes of *F. communis* are called alkalakh, which are used locally as a traditional medicine to treat skin infections, while its roasted flower buds are used to treat fever and dysentery [15]. In Lebanon and Syria, the roots of *F. hermonis* Boiss are used in folk medicine to reduce plasma cholesterol levels and total weight, as well as to treat skin infections, stomach diseases, erectile dysfunction, fever, dysentery, frostbite, and hysteria [16].

Phytochemical studies of the oleo–gum–resin, roots, seeds, and aerial parts of more than 70 species have revealed coumarins [17], phenylpropanoids [18], lignans [19], steroidal esters [20], organic acid glycosides [20], aromatic acids [21], sesquiterpenes [22], monoterpenes [23], benzofurans [24], and sulfur-containing derivatives [25] of the *Ferula* genus. Among these constituents, sesquiterpenes and their derivatives, especially sesquiterpene coumarins, sesquiterpene phenylpropanoids, and sesquiterpene chromones, have attracted the attention of scientists due to the diversity of their chemical structures, as well as their extensive and promising biological properties, such as antioxidative [26], anti-inflammatory [27], antibacterial [28], antitumor [29], and antiviral [30] properties.

While they are the main active ingredients in *Ferula*, there has not, however, been a comprehensive review of sesquiterpenes and their derivatives from this plant. In this review, we aim to report the chemical structures, biosynthetic pathways, and biological properties of sesquiterpenes and sesquiterpene derivatives from *Ferula*. Overall, the purpose of this work is to provide a comprehensive introduction to the bioactive sesquiterpenes of *Ferula*, which may help guide future research directions and possible application methods for this valuable edible and medicinal plant.

2. Chemical Structures

2.1. Sesquiterpenes

Ferula species are known for their production of various secondary metabolites, including sesquiterpenes. Sesquiterpenes are a class of terpenes composed of three isoprene units. The structural types of sesquiterpenes in *Ferula* are dominated by monocyclic and bicyclic sesquiterpenes, such as the daucane-type (I), guaiane-type (II), humulane-type (III), eudesmane-type (IV), germacrane-type (V), and elemene-type (VI) sesquiterpenes (Figure 1). Among them, the daucane-type sesquiterpene is the most common skeleton type.

Except for a few compounds, the sesquiterpenes in *Ferula* mostly exist in the form of esters, with substituents including fatty acids, aromatic acids, etc. Due to the presence of multiple substituent sites and substituents, the structures of sesquiterpenes are diverse. Here, sesquiterpenes with medicinal or potential medicinal prospects are summarized, including 88 daucane-type sesquiterpenes (1–88), 27 guaiane-type sesquiterpenes (89–115), 13 humulane-type sesquiterpenes (116–128), 11 eudesmane-type sesquiterpenes (129–139), 3 germacrane-type sesquiterpenes (140–142), 1 elemene-type sesquiterpene (143), and 6 other types of sesquiterpenes (144–149). The names and sources of compounds 1–149 are listed in Table 1, and their chemical structures are shown in Figure 2.




Figure 1. The main structural types of sesquiterpenes in *Ferula* plants (the daucane-type (I), guaiane-type (II), humulane-type (III), eudesmane-type (IV), germacrane-type (V), and elemene-type (VI) sesquiterpenes).

Table 1. Sesquiterpenes in *Ferula* plants.

Names	No.	Types	Species	References
2 α -Acetoxy-6 α - <i>p</i> -methoxybenzoyl-10 α -acetoxy-jaeschkeanadiol	6	I	<i>F. communis</i> subsp. <i>communis</i>	[31]
2 α -Acetoxy-6 α - <i>p</i> -methoxybenzoyl-10 α -hydroxy-jaeschkeanadiol	3	I	<i>F. communis</i> subsp. <i>communis</i>	[31]
2 α -Acetoxy-6 α - <i>p</i> -methoxybenzoyl-10 β -acetoxy-jaeschkeanadiol	5	I	<i>F. communis</i> subsp. <i>communis</i>	[31]
2 α -Acetoxy-6 α - <i>p</i> -methoxybenzoyl-10 β -hydroxy-jaeschkeanadiol	4	I	<i>F. communis</i> subsp. <i>communis</i>	[31]
2 α -Acetyl ferutinin	19	I	<i>F. campestris</i> (Besser) Grecescu	[32]
9- <i>O</i> -Acetyl-8- <i>O</i> -tigloyltovarol	140	V	<i>F. persica</i> Willd	[33]
Aeschkeanadiol-2-methylbutyrate	13	I	<i>F. linkii</i> Webb & Berthel.	[34]
8 α -Angeloyloxy-10 β -hydroxyslov-3-en-6,12-olide	113	II	<i>F. varia</i> (Schrenk) Trautv.	[35]
10-Angeloyloxy-6- <i>p</i> -hydroxybenzoyl-jaeschkeandiol	20	I	<i>F. communis</i> var. <i>brevifolia</i> <i>F. sinaica</i> L.	[36] [37]
2 α -Angeloyloxy-6-veratrate-jaeschkeanadiol	28	I	<i>F. communis</i> subsp. <i>communis</i> <i>F. licentiana</i> var. <i>tunshanica</i> (Su) Shan et Q.X.Liu	[38] [39]
14-(4'-Anisoyloxy)dauc-4,8-diene	38	I	<i>F. communis</i> subsp. <i>communis</i>	[40]
Badrakemonin	143	VI	<i>F. badrakema</i> Kos.-Pol.	[41]
Benzoylfervanol	124	III	<i>F. lycia</i> Boiss.	[22]
Carotdiol acetate	41	I	<i>F. linkii</i> Webb & Berthel.	[42]
Carotdiol veratrate	42	I	<i>F. linkii</i> Webb & Berthel.	[42]
Daucol	73	I	<i>F. linkii</i> Webb & Berthel.	[42]
10-Deangeloylpallinin	31	I	<i>F. arrigonii</i> Bocchieri	[43]
Decipenin D	132	IV	<i>F. penninervis</i> Regel et Schmalh.	[44]
7,11-Dehydrogrilactone	114	II	<i>F. arrigonii</i> Bocchieri	[43]

Table 1. Cont.

Names	No.	Types	Species	References
Dehydrooopin	138	IV	<i>F. varia</i> (Schrenk) Trautv.	[35]
2,10-Diacetyl-8-hydroxyferutrinol-6-anisate	49	I	<i>F. vesceritensis</i> Coss. & Durieu ex Trab.	[45]
3 α ,4 β -Dihydroxy-5 β H,11 α H-eudesman-6,12-olide	130	IV	<i>F. sinaica</i> L.	[46]
4 β ,8 β -Dihydroxy-6 α -(4-hydroxy-3-methoxybenzoyl)-dauc-9-ene	52	I	<i>F. hermonis</i> Boiss	[47]
			<i>F. kuhistanica</i> Korovin	[48]
4 β ,8 α -Dihydroxy-6 α -vanilloy-loxydauc-9-ene	54	I	<i>F. kuhistanica</i> Korovin	[48]
Diversolide A	103	II	<i>F. diversivittata</i> Regel & Schmalh.-Rech.	[49]
Diversolide B	104	II	<i>F. diversivittata</i> Regel & Schmalh.-Rech.	[49]
Diversolide C	105	II	<i>F. diversivittata</i> Regel & Schmalh.-Rech.	[49]
Diversolide D	106	II	<i>F. diversivittata</i> Regel & Schmalh.-Rech.	[49]
Diversolide E	107	II	<i>F. diversivittata</i> Regel & Schmalh.-Rech.	[49]
Diversolide F	108	II	<i>F. diversivittata</i> Regel & Schmalh.-Rech.	[49]
Diversolide G	109	II	<i>F. diversivittata</i> Regel & Schmalh.-Rech.	[49]
Elaeochytrin A	45	I	<i>F. elaeochytris</i> Korovin	[50]
Elaeochytrin B	50	I	<i>F. elaeochytris</i> Korovin	[50]
Epoxy ferutinin benzoate	80	I	<i>F. hermonis</i> Boiss	[51]
8,9-Epoxy-ferutinin	81	I	<i>F. kuhistanica</i> Korovin	[48]
Epoxyjaeschkeanadiol	78	I	<i>F. hermonis</i> Boiss	[52]
2,3-Epoxy-jaeschkeanadiol- <i>p</i> -methoxybenzoate	82	I	<i>F. kuhistanica</i> Korovin	[48]
			<i>F. communis</i> L.	[53]
			<i>F. lancerottensis</i> Parl.	[54]
			<i>F. linkii</i> Webb & Berthel.	[55]
Epoxyvesceritenol	84	I	<i>F. vesceritensis</i> Coss. & Durieu ex Trab.	[45]
(1 <i>R</i> ,3 <i>S</i> ,8 <i>S</i>)-3-Ethoxy-8-angeloyloxydauca-4-en-9-one	70	I	<i>F. hermonis</i> Boiss	[56]
Fercolide	47	I	<i>F. communis</i> subsp. <i>communis</i>	[40]
Fercomin	39	I	<i>F. communis</i> subsp. <i>communis</i>	[40]
			<i>F. licentiana</i> var. <i>tunshanica</i> (Su) Shan et Q.X.Liu	[57]
Ferugin	71	I	<i>F. sinaica</i> L.	[58]
Feruginin	48	I	<i>F. jaeschkeana</i> Vatke	[59,60]
Ferhermonin A	57	I	<i>F. hermonis</i> Boiss	[47]
Ferhermonin B	51	I	<i>F. hermonis</i> Boiss	[47]
Ferhermonins C	62	I	<i>F. hermonis</i> Boiss	[47]
Ferulactone A	147	others	<i>F. ferulaeoides</i> (Steud.) Korov	[61]
Ferulactone B	148	others	<i>F. ferulaeoides</i> (Steud.) Korov	[61]
Ferulinkiol-1-hydroxy-5(2-methylbutyrate)	11	I	<i>F. linkii</i> Webb & Berthel.	[34]
Ferupennin A	89	II	<i>F. penninervis</i> Regel et Schmalh.	[44]
Ferupennin B	90	II	<i>F. penninervis</i> Regel et Schmalh.	[44]
Ferupennin C	91	II	<i>F. penninervis</i> Regel et Schmalh.	[44]
Ferupennin D	92	II	<i>F. penninervis</i> Regel et Schmalh.	[44]
Ferupennin E	93	II	<i>F. penninervis</i> Regel et Schmalh.	[44]
Ferupennin F	94	II	<i>F. penninervis</i> Regel et Schmalh.	[44]
Ferupennin G	95	II	<i>F. penninervis</i> Regel et Schmalh.	[44]

Table 1. Cont.

Names	No.	Types	Species	References
Ferupennin H	96	II	<i>F. penninervis</i> Regel et Schmalh.	[44]
Ferupennin I	97	II	<i>F. penninervis</i> Regel et Schmalh.	[44]
Ferupennin J	98	II	<i>F. penninervis</i> Regel et Schmalh.	[44]
Ferupennin K	99	II	<i>F. penninervis</i> Regel et Schmalh.	[44]
Ferupennin L	100	II	<i>F. penninervis</i> Regel et Schmalh.	[44]
Ferupennin L	111	II	<i>F. varia</i> (Schrenk) Trautv.	[35]
Ferupennin M	101	II	<i>F. penninervis</i> Regel et Schmalh.	[44]
Ferupennin N	102	II	<i>F. penninervis</i> Regel et Schmalh.	[44]
Ferupennin O	112	II	<i>F. penninervis</i> Regel et Schmalh.	[44]
Ferutidin	14	I	<i>F. elaeochoytris</i> Korovin	[62]
			<i>F. lancerottensis</i> Parl.	[54]
			<i>F. licentiana</i> var. <i>tunshanica</i> (Su) Shan et Q.X.Liu	[57]
			<i>F. arrigonii</i> Bocchieri	[34]
			<i>F. communis</i> L.	[43]
			<i>F. communis</i> subsp. <i>communis</i>	[31]
			<i>F. glauca</i> subsp. <i>glauca</i>	[31,38]
Ferutinin	18	I	<i>F. kuhistanica</i> Korovin	[48]
			<i>F. hermonis</i> Boiss	[52]
			<i>F. jaeschkeana</i> Vatke	[59,60,63]
			<i>F. kuhistanica</i> Korovin	[64]
			<i>F. elaeochoytris</i> Korovin	[62]
			<i>F. sinaica</i> L.	[37]
			<i>F. lancerottensis</i> Parl.	[54]
			<i>F. licentiana</i> var. <i>tunshanica</i> (Su) Shan et Q.X.Liu	[57]
			<i>F. kingdom-wardii</i> Wolff	[57]
Ferutionone	33	I	<i>F. communis</i> L.	[53,65]
			<i>F. jaeschkeana</i> Vatke	[63]
Fetidone A	145	others	<i>F. assa-foetida</i> L.	[66]
Fetidone B	146	others	<i>F. assa-foetida</i> L.	[66]
2 α -Hydroxy ferutinin	1	I	<i>F. glauca</i> subsp. <i>glauca</i>	[31]
1 α -Hydroxy-2-oxo-5 α ,7 β -11 β H-eudesm-3-en-6 α ,12-olide	129	IV	<i>F. penninervis</i> Regel et Schmalh.	[44]
2 β -Hydroxy-3,4-epoxyjaes-chkeanadiol	74	I	<i>F. jaeschkeana</i> Vatke	[67]
14-(4'-Hydroxy-3'-methoxy-benzoyloxy)dauc-4,8-diene	37	I	<i>F. hermonis</i> Boiss	[52]
2 α -Hydroxy-6 α -p-methoxybenzoyl-10 β -acetoxo-jaeschkeanadiol	2	I	<i>F. communis</i> subsp. <i>communis</i>	[31]
5-p-Hydroxybenzoyl ester of ferutiol	55	I	<i>F. sinaica</i> L.	[37]
14-(4'-Hydroxybenzoyloxy)-dauc-4,8-diene	36	I	<i>F. hermonis</i> Boiss	[52]
5 α -p-Hydroxybenzoyloxydauc-2-ene-1-one	23	I	<i>F. kuhistanica</i> Korovin	[48]
6- β -p-Hydroxybenzoyloxygermacra-1(10),4-diene	141	V	<i>F. lycia</i> Boiss.	[22]
14-Hydroxy-dauc-4-ene	68	I	<i>F. sinaica</i> L.	[68]
(1R,4R)-4-Hydroxydauc-7-ene-6,9-dione	64	I	<i>F. hermonis</i> Boiss	[56]
(1R,4R)-4-Hydroxydauc-7-ene-6-one	63	I	<i>F. hermonis</i> Boiss	[56]
10 α -Hydroxyfertidin	30	I	<i>F. arrigonii</i> Bocchieri	[43]
10-Hydroxylancerodiol-6-anisate	59	I	<i>F. vesceritensis</i> Coss. & Durieu ex Trab.	[45]
10-Hydroxylancerodiol-6-benzoate	60	I	<i>F. vesceritensis</i> Coss. & Durieu ex Trab.	[45]
14-Hydroxyvaginatin	44	I	<i>F. sinaica</i> L.	[37]
14-Hydroxyvaginatin	86	I	<i>F. sinaica</i> L.	[37]
Isolancerotriol	72	I	<i>F. sinaica</i> L.	[37,58]

Table 1. Cont.

Names	No.	Types	Species	References
5-Isovalerate of lapiferol	83	I	<i>F. communis</i> L. <i>F. linkii</i> Webb & Berthel.	[53] [69]
Jaeschkeanadiol	16	I	<i>F. hermonis</i> Boiss <i>F. jaeschkeana</i> Vatke <i>F. kuhistanica</i> Korovin <i>F. elaeochytris</i> Korovin <i>F. sinaica</i> L. <i>F. communis</i> L. <i>F. lancerottensis</i> Parl.	[52] [59,60,63] [48,64] [62] [37] [53] [54]
Jaeschkeanadiol isovalerate	12	I	<i>F. linkii</i> Webb & Berthel.	[34]
Jaeskeanadiol angelate	26	I	<i>F. jaeschkeana</i> Vatke <i>F. lancerottensis</i> Parl. <i>F. elaeochytris</i> Korovin	[63] [54] [62]
Jaeskeanadiol salicylate	27	I	<i>F. elaeochytris</i> Korovin	[62]
Jaeskeanadiol veratrate	29	I	<i>F. arrigonii</i> Bocchieri <i>F. licentiana</i> var. <i>tunshanica</i> (Su) Shan et Q.X.Liu	[43] [57]
Juniferdin	121	III	<i>F. lycia</i> Boiss.	[22]
Juniferin	123	III	<i>F. lycia</i> Boiss.	[22]
Juniferinin	122	III	<i>F. lycia</i> Boiss.	[22]
Kuhistaferone	149	others	<i>F. kuhistanica</i> Korovin	[70]
Kuhistanicaol A	76	I	<i>F. kuhistanica</i> Korovin	[48]
Kuhistanicaol D	24	I	<i>F. kuhistanica</i> Korovin	[48]
Kuhistanicaol E	65	I	<i>F. kuhistanica</i> Korovin	[48]
Kuhistanicaol F	66	I	<i>F. kuhistanica</i> Korovin	[48]
Kuhistanicaol G	53	I	<i>F. kuhistanica</i> Korovin	[48]
Kuhistanicaol H	75	I	<i>F. kuhistanica</i> Korovin	[64]
Kuhistanicaol I	25	I	<i>F. kuhistanica</i> Korovin	[64]
Kuhistanicaol J	88	I	<i>F. kuhistanica</i> Korovin	[64]
Lancerodiol- <i>p</i> -hydroxybenzoate	58	I	<i>F. lancerottensis</i> Parl. <i>F. jaeschkeana</i> Vatke <i>F. sinaica</i> L. <i>F. linkii</i> Webb & Berthel.	[54] [67] [58] [71]
Lancerodiol- <i>p</i> -methoxybenzoate	67	I	<i>F. glauca</i> subsp. <i>glauca</i>	[31]
Lancerotriol-9-acetate-6- <i>p</i> -hydrooxybenzoate	56	I	<i>F. sinaica</i> L.	[46]
Lanerotriol- <i>p</i> -hydroxy-benzoate	61	I	<i>F. kuhistanica</i> Korovin	[48]
Lapidin	35	I	<i>F. lapidosa</i> Korov.	[72]
Lapido	34	I	<i>F. jaeschkeana</i> Vatke	[67]
Lapido isobutyrate	9	I	<i>F. linkii</i> Webb & Berthel.	[34]
Lapido vanillate	22	I	<i>F. kuhistanica</i> Korovin	[48]
Lapido-2-methybutyrate	8	I	<i>F. linkii</i> Webb & Berthel.	[34]
Lapido- <i>p</i> -anisate	10	I	<i>F. linkii</i> Webb & Berthel.	[34]
Lapiferin	77	I	<i>F. vesceritensis</i> Coss. & Durieu ex Trab. <i>F. arrigonii</i> Bocchieri	[73] [43]
Lasidiol-10-anisate	43	I	<i>F. vesceritensis</i> Coss. & Durieu ex Trab.	[45]
Lyciferin A	116	III	<i>F. lycia</i> Boiss.	[22]
Lyciferin B	117	III	<i>F. lycia</i> Boiss.	[22]
Lyciferin C	118	III	<i>F. lycia</i> Boiss.	[22]
Lyciferin D	119	III	<i>F. lycia</i> Boiss.	[22]
Lyciferin E	120	III	<i>F. lycia</i> Boiss.	[22]
1 α ,10 β -Epoxy-4-humden-6 β - <i>p</i> -anisate	127	III	<i>F. linkii</i> Webb & Berthel.	[34]
1 α ,10 β -Epoxy-4-humden-6 β - <i>p</i> -veratrate	128	III	<i>F. linkii</i> Webb & Berthel.	[34]
14- <i>p</i> -Methoxybenzoyl-4,5-epoxy-dauc-8-ene	46	I	<i>F. communis</i> subsp. <i>communis</i>	[31]
Nerolidol	144	others	<i>F. fukanensis</i> K. M. Shen	[74]
Oopodin	139	IV	<i>F. varia</i> (Schrenk) Trautv.	[35]

Table 1. Cont.

Names	No.	Types	Species	References
2-Oxoferutidin	32	I	<i>F. arrigonii</i> Bocchieri	[43]
Pallinin or 6 α ,10 α -diangeloyl-jaeschkeanadiol	7	I	<i>F. communis</i> subsp. <i>communis</i>	[31]
Penninnervin	131	IV	<i>F. penninervis</i> Regel et Schmalh.	[44]
<i>p</i> -Hydroxybenzoylfervanol	125	III	<i>F. lycia</i> Boiss.	[22]
Spathulenol	115	II	<i>F. varia</i> (Schrenk) Trautv.	[35]
Teferidin	17	I	<i>F. hermonis</i> Boiss	[52]
			<i>F. elaeochytris</i> Korovin	[62]
			<i>F. jaeschkeana</i> Vatke	[63]
			<i>F. sinaica</i> L.	[37]
Teferin	21	I	<i>F. hermonis</i> Boiss	[52]
			<i>F. jaeschkeana</i> Vatke	[59,60,63]
			<i>F. kuhistanica</i> Korovin	[48,64]
			<i>F. kingdom-ardii</i> Wolff	[57]
			<i>F. elaeochytris</i> Korovin	[62]
4 β , 8 β , 9 α -Trihydroxy-6 α - <i>p</i> -hydroxybenzoyoxydaucane	87	I	<i>F. sinaica</i> L.	[37,58]
Tunetanin A	15	I	<i>F. tunetana</i> Pom	[75]
Vaginatn	40	I	<i>F. communis</i> subsp. <i>communis</i>	[40]
5 α -Vanillate-2,3-epoxy-jaeschkeanadiol	79	I	<i>F. kuhistanica</i> Korovin	[48]
			<i>F. jaeschkeana</i> Vatke	[67]
Vanilloylfervanol	126	III	<i>F. lycia</i> Boiss.	[22]
6- β -Vanilloxyoxygermacra-1(10),4-diene	142	V	<i>F. lycia</i> Boiss.	[22]
Vesceritenone	85	I	<i>F. vesceritensis</i> Coss. & Durieu ex Trab.	[45]
Webbiol angelate	69	I	<i>F. linkii</i> Webb & Berthel.	[71]
-	110	II	<i>F. varia</i> (Schrenk) Trautv.	[35]
-	133	IV	<i>F. varia</i> (Schrenk) Trautv.	[35]
-	134	IV	<i>F. varia</i> (Schrenk) Trautv.	[35]
-	135	IV	<i>F. varia</i> (Schrenk) Trautv.	[35]
-	136	IV	<i>F. varia</i> (Schrenk) Trautv.	[35]
-	137	IV	<i>F. varia</i> (Schrenk) Trautv.	[35]

2.2. Sesquiterpene Coumarins

Sesquiterpene coumarins are often found in *Ferula* plants and are known for their unique chemical structures and potential bioactivity. According to the connection site between the sesquiterpene unit and the coumarin skeleton, sesquiterpene coumarins can be classified into those connected by a 7-position C-O-C bridge (I) (150–345), those connected by a 4-position C-O-C bridge (II) (346–348), and those connected by a 3-position C-C bond (III) (349–407). According to the structural types of sesquiterpenes, type I compounds can be further classified into a straight-chain type (Ia) (150–175), monocyclic type (Ib) (176–219), and bicyclic type (Ic) (220–345). Type III compounds can be classified into straight-chain coumarin type (IIIa) (349–361), furanocoumarin type (IIIb) (362–399), and pyranocoumarin type (IIIc) (400–407) compounds depending on whether the hydroxyl group in the sesquiterpene moiety forms a five- or six-membered heterocyclic ring with the coumarin moiety.

The names and sources of sesquiterpene coumarins (150–407) are listed in Table 2, and their chemical structures are shown in Figure 3.




Figure 2. Cont.


Figure 2. Sesquiterpenes in *Ferula* plants.

Table 2. Sesquiterpene coumarins in *Ferula* plants.

Names	No.	Types	Species	References
ϵ -Acetoxy,4-acetylferulenol	357	IIIa	<i>F. communis</i> var. <i>genuina</i>	[76]
10' <i>R</i> -Acetoxy-11'-hydroxyumbelliprenin	167	Ia	<i>F. assa-foetida</i> L.	[10]
5'-Acetoxy-8'-hydroxyumbelliprenin	156	Ia	<i>F. assa-foetida</i> L.	[10]
8'-Acetoxy-5'-hydroxyumbelliprenin	155	Ia	<i>F. assa-foetida</i> L.	[77]
(<i>Z</i>)- ω -Acetoxyferprenin	401	IIIc	<i>F. communis</i> L.	[78,79]
(<i>E</i>)- ω -Acetoxyferprenin	402	IIIc	<i>F. communis</i> L.	[78,79]
(<i>E</i>)- ω -Acetoxyferulenol	352	IIIa	<i>F. communis</i> L.	[80]
(<i>Z</i>)- ω -Acetoxyferulenol	353	IIIa	<i>F. communis</i> L.	[80]
8- <i>O</i> -Acetyl-sinkiangenorin F	209	Ib	<i>F. sinkiangensis</i> K. M. Shen	[81]
Ammoresinal	356	IIIa	<i>F. vesceritensis</i> Coss. & Durieu ex Trab.	[45]
3-Angeloxycoladin	242	Ic	<i>F. vesceritensis</i> Coss. & Durieu ex Trab.	[5]
Asacoumarin A	154	Ia	<i>F. assa-foetida</i> L. <i>F. foetida</i> (Bunge) Regel	[82] [83]
Asacoumarin B	217	Ib	<i>F. assa-foetida</i> L.	[82]
Asimafoetida	189	Ib	<i>F. assafoetida</i> Linn.	[84]
Asimafoetidol	203	Ib	<i>F. assa-foetida</i> L.	[85]
Assafoetidin	190	Ib	<i>F. fukanensis</i> K. M. Shen <i>F. lehmannii</i> Boss. <i>F. assafoetida</i> Linn.	[86] [87] [88]
Assafoetidol A	230	Ic	<i>F. assa-foetida</i> L.	[89]
Assafoetidol B	231	Ic	<i>F. assa-foetida</i> L.	[89]
Badrakemin	238	Ic	<i>F. teterrima</i> Kar. et Kir. <i>F. badrakema</i> Kos.-Pol.	[90] [91]
Badrakemin acetate	239	Ic	<i>F. teterrima</i> Kar. et Kir. <i>F. badrakema</i> Kos.-Pol.	[90] [41]
Badrakemone	237	Ic	<i>F. fukanensis</i> K. M. Shen <i>F. teterrima</i> Kar. et Kir. <i>F. persica</i> Willd <i>F. nevskii</i> Korov.	[86] [90] [92] [93]
Cauferidin	245	Ic	<i>F. conocaula</i> Korov.	[94]
Cauferin	249	Ic	<i>F. conocaula</i> Korov.	[94]
Cauferinin	259	Ic	<i>F. conocaula</i> korov. <i>F. samarkandica</i> Korovin	[95] [96]
Cauferoside	235	Ic	<i>F. gumosa</i> Boiss. <i>F. conocaula</i> Korov.	[97] [98]
Cauloside	252	Ic	<i>F. conocaula</i> Korov.	[99]
Cocanicin	160	Ia	<i>F. cocanica</i>	[99]
Coladin	232	Ic	<i>F. sinkiangensis</i> K. M. Shen <i>F. tunetana</i> Pom <i>F. campestris</i> (Besser) Grecescu	[5] [5,75] [100]
Colladonin	240	Ic	<i>F. teterrima</i> Kar. et Kir. <i>F. foetida</i> (Bunge) Regel <i>F. sinkiangensis</i> K. M. Shen <i>F. sinaica</i> L. <i>F. tunetana</i> POM <i>F. campestris</i> (Besser) Grecescu	[90] [83] [101] [5] [75] [100]
Colladonin isovalerate	241	Ic	<i>F. loscossi</i> (Lge) Wk	[102]
Communiferulin A	366	IIIb	<i>F. communis</i> L.	[103]
Communiferulin B	367	IIIb	<i>F. communis</i> L.	[103]
Communiferulin C	368	IIIb	<i>F. communis</i> L.	[103]
Conferdione	283	Ic	<i>F. flabelliloba</i> Rech. f. & Aell	[104,105]
Conferin	286	Ic	<i>F. conocaula</i> Korov.	[106]
Conferol	267	Ic	<i>F. assa-foetida</i> L. <i>F. gumosa</i> Boiss. <i>F. conocaula</i> Korov.	[10] [97] [107]
Conferol acetate	264	Ic	<i>F. badrakema</i> Kos.-Pol.	[41]

Table 2. Cont.

Names	No.	Types	Species	References
Conferone	265	Ic	<i>F. flabelliloba</i> Rech. f. & Aell <i>F. badrakema</i> Kos.-Pol.	[104] [41,108]
Conferoside	236	Ic	<i>F. gumosa</i> Boiss. <i>F. conocaula</i> Korov.	[97] [98]
Conferoside	285	Ic	<i>F. conocaula</i> Korov.	[98]
Deacetylkellerin	323	Ic	<i>F. kelleri</i> K.-Pol. <i>F. kokanica</i> Regel & Schmalh.	[109] [110]
Deacetyltdshikorin	158	Ia	<i>F. tadshikorum</i> M. Pimen	[111]
Diastereomer-samarcandin	289	Ic	<i>F. sinaica</i> L.	[112]
2,3-Dihydro-7-hydroxy-2 <i>R</i> *,3 <i>R</i> *-dimethyl-2-[4,8-dimethyl-3(<i>E</i>),7-nonadien-6-onyl]-furo[3,2- <i>c</i>]coumarin	399	IIIb	<i>F. fukanensis</i> K. M. Shen	[113]
2,3-Dihydro-7-hydroxy-2 <i>R</i> *,3 <i>R</i> *-dimethyl-2-[4,8-dimethyl-3(<i>E</i>),7-nonadienyl]-furo[3,2- <i>c</i>]coumarin	386	IIIb	<i>F. ferulaeoides</i> (Steud.) Korov	[114]
2,3-Dihydro-7-hydroxy-2 <i>R</i> *,3 <i>R</i> *-dimethyl-2-[4-methyl-5-(4-methyl-2-furyl)-3(<i>E</i>)-pentenyl]-furo[3,2- <i>c</i>]coumarin	394	IIIb	<i>F. ferulaeoides</i> (Steud.) Korov	[114]
2,3-Dihydro-7-hydroxy-2 <i>S</i> *,3 <i>R</i> *-dimethyl-2-[4,8-dimethyl-3(<i>E</i>),7-nonadien-6-onyl]-furo[3,2- <i>c</i>]coumarin	390	IIIb	<i>F. ferulaeoides</i> (Steud.) Korov	[114]
2,3-Dihydro-7-hydroxy-2 <i>S</i> *,3 <i>R</i> *-dimethyl-2-[4,8-dimethyl-3(<i>E</i>),7-nonadienyl]-furo[3,2- <i>c</i>]coumarin	383	IIIb	<i>F. ferulaeoides</i> (Steud.) Korov	[114]
2,3-Dihydro-7-hydroxy-2 <i>S</i> *,3 <i>R</i> *-dimethyl-2-[4-methyl-5-(4-methyl-2-furyl)-3(<i>E</i>)-pentenyl]-furo[3,2- <i>c</i>]coumarin	392	IIIb	<i>F. ferulaeoides</i> (Steud.) Korov	[114]
2,3-Dihydro-7-hydroxy-2 <i>S</i> *,3 <i>R</i> *-dimethyl-3-[4,8-dimethyl-3(<i>E</i>),7-nonadienyl]-furo[3,2- <i>c</i>]coumarin	370	IIIb	<i>F. ferulaeoides</i> (Steud.) Korov	[114]
2,3-Dihydro-7-hydroxy-2 <i>S</i> *,3 <i>R</i> *-dimethyl-3-[4-methyl-5-(4-methyl-2-furyl)-3(<i>E</i>)-pentenyl]-furo[3,2- <i>c</i>]coumarin	373	IIIb	<i>F. ferulaeoides</i> (Steud.) Korov	[114]
2,3-Dihydro-7-hydroxy-2 <i>S</i> *,3 <i>S</i> *-dimethyl-2-[4,8-dimethyl-3(<i>E</i>),7-nonadienyl]-furo[3,2- <i>c</i>]coumarin	389	IIIb	<i>F. ferulaeoides</i> (Steud.) Korov	[115]
2,3-Dihydro-7-hydroxy-2 <i>S</i> *,3 <i>S</i> *-dimethyl-3-[4,8-dimethyl-3(<i>E</i>),7-nonadienyl]-furo[3,2- <i>c</i>]coumarin	372	IIIb	<i>F. ferulaeoides</i> (Steud.) Korov	[114]
(2 <i>S</i> *,3 <i>R</i> *)-2,3-Dihydro-7-hydroxy-2-[(3 <i>E</i>)-8-hydroxy-4,8-dimethylnon-3-en-1-yl]-2,3-dimethyl-4 <i>H</i> -furo[3,2- <i>c</i>][1]benzopyran-4-one	364	IIIb	<i>F. ferulaeoides</i> (Steud.) Korov	[116]
(2 <i>S</i> *,3 <i>R</i> *)-2,3-Dihydro-2-[(3 <i>E</i>)-8-hydroxy-4,8-dimethylnon-3-en-1-yl]-7-methoxy-2,3-dimethyl-4 <i>H</i> -furo[3,2- <i>c</i>][1]benzopyran-4-one	365	IIIb	<i>F. ferulaeoides</i> (Steud.) Korov	[116]
2,3-Dihydro-7-methoxy-2 <i>R</i> *,3 <i>R</i> *-dimethyl-2-[4,8-dimethyl-3(<i>E</i>),7-nonadienyl]-furo[3,2- <i>c</i>]coumarin	387	IIIb	<i>F. ferulaeoides</i> (Steud.) Korov	[114]
2,3-Dihydro-7-methoxy-2 <i>S</i> *,3 <i>R</i> *-dimethyl-2-[4,8-dimethyl-3(<i>E</i>),7-nonadien-6-onyl]-furo[3,2- <i>c</i>]coumarin	391	IIIb	<i>F. ferulaeoides</i> (Steud.) Korov	[114]
2,3-Dihydro-7-methoxy-2 <i>S</i> *,3 <i>R</i> *-dimethyl-2-[4,8-dimethyl-3(<i>E</i>),7-nonadienyl]-furo[3,2- <i>c</i>]coumarin	384	IIIb	<i>F. ferulaeoides</i> (Steud.) Korov	[114]
2,3-Dihydro-7-methoxy-2 <i>S</i> *,3 <i>R</i> *-dimethyl-2-[4-methyl-5-(4-methyl-2-furyl)-3(<i>E</i>)-pentenyl]-furo[3,2- <i>c</i>]coumarin	393	IIIb	<i>F. ferulaeoides</i> (Steud.) Korov	[114]
2,3-Dihydro-7-methoxy-2 <i>S</i> *,3 <i>R</i> *-dimethyl-3-[4,8-dimethyl-3(<i>E</i>),7-nonadienyl]-furo[3,2- <i>c</i>]coumarin	371	IIIb	<i>F. ferulaeoides</i> (Steud.) Korov	[114]
6',7'-Dihydroxy-karatavicinol	169	Ia	<i>F. sinaica</i> L.	[37]
(2 <i>S</i> *,3 <i>S</i> *)-3-[(3 <i>E</i>)-4,8-Dimethylnona-3,7-dien-1-yl]-2,3-dihydro-7-methoxy-2,3-dimethyl-4 <i>H</i> -furo[3,2- <i>c</i>][1]benzopyran-4-one	363	IIIb	<i>F. ferulaeoides</i> (Steud.) Korov	[116]
Drimatol B	345	Ic	<i>F. jaeschkeana</i> Vatke	[63]
Ent-Colladonin	248	Ic	<i>F. sinkiangensis</i> K. M. Shen	[117]
Epi-Conferdione	272	Ic	<i>F. assa-foetida</i> L. <i>F. foetida</i> (Bunge) Regel	[10] [83]
Episamarcandin	305	Ic	<i>F. sinkiangensis</i> K. M. Shen <i>F. sinaica</i> L.	[118] [119]
Episamarcandin acetate	295	Ic	<i>F. assa-foetida</i> L.	[120]
Epoxyfarnochrol	175	Ia	<i>F. jaeschkeana</i> Vatke	[63]
Ethyl galbanate	215	Ib	<i>F. pseudalliacea</i> Rech.f.	[121]

Table 2. *Cont.*

Names	No.	Types	Species	References
Farnesiferol A	228	Ic	<i>F. assafoetida</i> Linn.	[84]
			<i>F. assa-foetida</i> L.	[10]
			<i>F. vesceritensis</i> Coss. & Durieu ex Trab.	[45]
			<i>F. persica</i> Willd	[92]
Farnesiferol B	186	Ib	<i>F. sinkiangensis</i> K. M. Shen	[118]
			<i>F. szowitsiana</i> DC.	[122]
			<i>F. flabelliloba</i> Rech. f. & Aell	[104]
			<i>F. lehmannii</i> Boss.	[123]
			<i>F. asafoetida</i> L.	[124]
			<i>F. assa-foetida</i> L.	[10]
Farnesiferol C	202	Ib	<i>F. persica</i> Willd	[92]
			<i>F. lehmannii</i> Boss.	[87]
			<i>F. sinkiangensis</i> K. M. Shen	[118]
			<i>F. szowitsiana</i> DC.	[122,125]
Farnesiferone B	185	Ib	<i>F. asafoetida</i> L.	[124]
			<i>F. assafoetida</i> Linn.	[84]
			<i>F. flabelliloba</i> Rech. f. & Aell	[104]
Fecarpin	330	Ic	<i>F. microcarpa</i> Korovin	[126]
Fekolin	193	Ib	<i>F. kopetdagensis</i> Eug. Kor.	[99,127]
Fekolone	187	Ib	<i>F. fukanensis</i> K. M. Shen	[86]
			<i>F. sinkiangensis</i> K. M. Shen	[118,128]
			<i>F. kopetdagensis</i> Eug. Kor.	[127]
Fekolone	195	Ib	<i>F. kopetdagensis</i> Eug. Kor.	[99,127]
Fekrol	196	Ib	<i>F. krylovii</i> Korov.	[129]
Fekrynol	211	Ib	<i>F. sinkiangensis</i> K. M. Shen	[118]
			<i>F. lehmannii</i> Boss.	[87]
			<i>F. krylovii</i> Korov.	[130]
Fekrynol acetate	212	Ib	<i>F. lehmannii</i> Boss.	[123]
			<i>F. krylovii</i> Korov.	[130]
Fepaldlin	322	Ic	<i>F. pallida</i> Korovin	[99]
Fercoprenol	369	IIIb	<i>F. communis</i> subsp. <i>communis</i>	[131]
Ferocaulicin	288	Ic	<i>F. conocaula</i> Korov.	[132]
Ferocaulidin	287	Ic	<i>F. gumosa</i> Boiss.	[97]
			<i>F. badrakema</i> Kos.-Pol	[41,108]
			<i>F. conocaula</i> Korov.	[132]
Ferocaulin	282	Ic	<i>F. conocaula</i> Korov.	[132]
Ferocaulinin	284	Ic	<i>F. conocaula</i> Korov.	[132]
Feropolidin	263	Ic	<i>F. polyantha</i> Korovin	[133,134]
			<i>F. vicaria</i> Korovin	[135]
Feropolin	207	Ib	<i>F. polyantha</i> Korovin	[133,134]
Feropolol	204	Ib	<i>F. polyantha</i> Korovin	[133,134]
			<i>F. vicaria</i> Korovin	[135]
Feropolone	206	Ib	<i>F. polyantha</i> Korovin	[133,134]
			<i>F. vicaria</i> Korovin	[135]
Feroside	164	Ia	<i>F. korshinskyi</i> Eug. Korov	[136]
Ferprenin	405	IIIc	<i>F. communis</i> L.	[78,79]
Ferubungeanol a	307	Ic	<i>F. bungeana</i> Kitag.	[137]
Ferubungeanol b	308	Ic	<i>F. bungeana</i> Kitag.	[137]
Ferubungeanol c	309	Ic	<i>F. bungeana</i> Kitag.	[137]
Ferubungeanol d	310	Ic	<i>F. bungeana</i> Kitag.	[137]
Ferubungeanol e	311	Ic	<i>F. bungeana</i> Kitag.	[137]
Ferubungeanol f	312	Ic	<i>F. bungeana</i> Kitag.	[137]
Ferubungeanol g	313	Ic	<i>F. bungeana</i> Kitag.	[137]
Ferubungeanol h	314	Ic	<i>F. bungeana</i> Kitag.	[137]
Ferucrin isobutyrate	324	Ic	<i>F. foetidissima</i> Regel & Schmalh.	[138]
Ferucrinone	325	Ic	<i>F. foetidissima</i> Regel & Schmalh.	[138]

Table 2. Cont.

Names	No.	Types	Species	References
Ferukrin	319	Ic	<i>F. kopetdagensis</i> Eug. Kor. <i>F. krylovii</i> Korov.	[139] [140]
Ferukrin acetate	320	Ic	<i>F. kopetdagensis</i> Eug. Kor.	[139]
Ferulenol	349	IIIa	<i>F. communis</i> var. <i>genuina</i>	[141,142]
Ferulenoloxyl ferulenol	361	IIIa	<i>F. communis</i> var. <i>genuina</i>	[76]
Ferulin A	395	IIIb	<i>F. ferulaeoides</i> (Steud.) Korov	[143]
Ferulin B	374	IIIb	<i>F. ferulaeoides</i> (Steud.) Korov	[143]
Ferulin C	375	IIIb	<i>F. ferulaeoides</i> (Steud.) Korov	[143]
Ferulsinic acid	218	Ib	<i>F. sinaica</i> L.	[144,145]
Ferusingensine A	171	Ia	<i>F. sinkiangensis</i> K. M. Shen	[146]
Ferusingensine B	172	Ia	<i>F. sinkiangensis</i> K. M. Shen	[146]
Ferusingensine C	173	Ia	<i>F. sinkiangensis</i> K. M. Shen	[146]
Ferusingensine D	174	Ia	<i>F. sinkiangensis</i> K. M. Shen	[146]
Ferusingensine E	168	Ia	<i>F. sinkiangensis</i> K. M. Shen	[146]
Ferusingensine F	183	Ib	<i>F. sinkiangensis</i> K. M. Shen	[146]
Ferusingensine G	177	Ib	<i>F. sinkiangensis</i> K. M. Shen	[146]
Ferusingensine H	326	Ic	<i>F. sinkiangensis</i> K. M. Shen	[146]
Ferusinol	338	Ic	<i>F. sinaica</i> L.	[112]
Feselol	268	Ic	<i>F. assa-foetida</i> L.	[10]
			<i>F. flabelliloba</i> Rech. f. & Aell	[104]
			<i>F. badrakema</i> Kos.-Pol	[41,108]
			<i>F. vesceritensis</i> Coss. & Durieu ex Trab.	[45]
Feselol angelate	270	Ic	<i>F. diversivittata</i> Regel & Schmalh.-Rech.	[147]
Feshurin	301	Ic	<i>F. teterrima</i> Kar. et Kir.	[90,148]
Feshurin acetate	302	Ic	<i>F. teterrima</i> Kar. et Kir.	[90]
Fesinkin A	184	Ib	<i>F. sinkiangensis</i> K. M. Shen	[149]
Fesinkin B	178	Ib	<i>F. sinkiangensis</i> K. M. Shen	[149]
Fesinkin C	179	Ib	<i>F. sinkiangensis</i> K. M. Shen	[149]
4'E-Fesinkin D	180	Ib	<i>F. sinkiangensis</i> K. M. Shen	[149]
4'Z-Fesinkin D	181	Ib	<i>F. sinkiangensis</i> K. M. Shen	[149]
Fesinkin E	339	Ic	<i>F. sinkiangensis</i> K. M. Shen	[149]
Fesinkin F	340	Ic	<i>F. sinkiangensis</i> K. M. Shen	[149]
Fesinkin G	341	Ic	<i>F. sinkiangensis</i> K. M. Shen	[149]
Feterin	250	Ic	<i>F. teterrima</i> Kar. et Kir.	[150]
Feterin acetate	251	Ic	<i>F. teterrima</i> Kar. et Kir.	[150]
Flabellilobin A	188	Ib	<i>F. flabelliloba</i> Rech. f. & Aell	[104]
Flabellilobin B	192	Ib	<i>F. flabelliloba</i> Rech. f. & Aell	[104]
Fnarthexol	279	Ic	<i>F. narthex</i> Boiss	[151]
Fnarthexone	246	Ic	<i>F. narthex</i> Boiss	[151]
Foetidin	346	II	<i>F. assa-foetida</i> L.	[152]
Foetidin acetate	347	II	<i>F. marmarica</i> Asch. & Taub.	[153]
Foetidone	348	II	<i>F. marmarica</i> Asch. & Taub.	[153]
Foliferin	205	Ib	<i>F. folioca</i> Lipsky	[154]
			<i>F. schtschurowskiana</i> Regel & Schmalh.	[99]
Fukanefuomarin A	376	IIIb	<i>F. fukanensis</i> K. M. Shen	[113]
Fukanefuomarin B	377	IIIb	<i>F. fukanensis</i> K. M. Shen	[113]
Fukanefuomarin C	378	IIIb	<i>F. fukanensis</i> K. M. Shen	[113]
Fukanefuomarin D	379	IIIb	<i>F. fukanensis</i> K. M. Shen	[113]
Fukanefuomarin E	380	IIIb	<i>F. fukanensis</i> K. M. Shen	[155]
Fukanefuomarin F	381	IIIb	<i>F. fukanensis</i> K. M. Shen	[155]

Table 2. Cont.

Names	No.	Types	Species	References
Fukanefuromarin G	382	IIIb	<i>F. fukanensis</i> K. M. Shen	[155]
Fukanefuromarin H	385	IIIb	<i>F. fukanensis</i> K. M. Shen	[156]
Fukanefuromarin I	388	IIIb	<i>F. fukanensis</i> K. M. Shen	[156]
Fukanefuromarin J	396	IIIb	<i>F. fukanensis</i> K. M. Shen	[156]
Fukanefuromarin K	397	IIIb	<i>F. fukanensis</i> K. M. Shen	[156]
Fukanefuromarin L	398	IIIb	<i>F. fukanensis</i> K. M. Shen	[156]
Fukanefuromarin M	407	IIIc	<i>F. fukanensis</i> K. M. Shen	[156]
Fukanemarin A	358	IIIa	<i>F. fukanensis</i> K. M. Shen	[113]
Fukanemarin B	360	IIIa	<i>F. fukanensis</i> K. M. Shen	[155]
Galbanic acid	213	Ib	<i>F. szowitsiana</i> DC. <i>F. asafoetida</i> L. <i>F. assa-foetida</i> L.	[122,157] [124] [10]
Gummosin	225	Ic	<i>F. persica</i> Willd	[92]
Gumoside A	233	Ic	<i>F. gumosa</i> Boiss.	[97]
Gumoside B	234	Ic	<i>F. gumosa</i> Boiss.	[97]
Gumosin	315	Ic	<i>F. gumosa</i> Boiss.	[97]
4'-Hydroxy kamolonol acetate	277	Ic	<i>F. pseudalliacea</i> Rech.f.	[158]
(Z)- ω -Hydroxyferprenin	403	IIIc	<i>F. communis</i> L.	[78,79]
(E)- ω -Hydroxyferprenin	404	IIIc	<i>F. communis</i> L.	[78,79]
(E)- ω -Hydroxyferulenol	350	IIIa	<i>F. communis</i> L.	[80]
(Z)- ω -Hydroxyferulenol	351	IIIa	<i>F. communis</i> L.	[80]
ϵ -Hydroxyferulenol	355	IIIa	<i>F. communis</i> L.	[76]
13-Hydroxyfeselol	271	Ic	<i>F. vesceritensis</i> Coss. & Durieu ex Trab. <i>F. tunetana</i> POM	[5] [75]
5'-Hydroxyumbelliprenin	151	Ia	<i>F. assa-foetida</i> L. <i>F. assa-foetida</i> L.	[77] [10]
8'-Hydroxyumbelliprenin	152	Ia	<i>F. assa-foetida</i> L.	[77]
Isoferprenin	406	IIIc	<i>F. communis</i> var. <i>genuina</i>	[159]
Isofeterin	243	Ic	<i>F. teterrima</i> Kar. et Kir.	[160]
Isosamarcandin	303	Ic	<i>F. sinaica</i> L. <i>F. sinkiangensis</i> K. M. Shen <i>F. microloba</i> Boiss.	[68] [118] [161]
Isosamarkandin	293	Ic	<i>F. sinkiangensis</i> K. M. Shen	[117]
Isosamarkandin angelate	304	Ic	<i>F. arrigonii</i> Bocchieri	[43]
Isosmarcandin	297	Ic	<i>F. tunetana</i> POM	[75]
Kamolol	331	Ic	<i>F. penninervis</i> Regel et Schmalh.	[162]
(3'S, 4'S, 5'R, 8'S, 9'S, 10'S)-Kamolol acetate	327	Ic	<i>F. sinkiangensis</i> K. M. Shen	[146]
Kamolone	332	Ic	<i>F. penninervis</i> Regel et Schmalh.	[162]
Kamololol	334	Ic	<i>F. assa-foetida</i> L. <i>F. pseudalliacea</i> Rech.f.	[10] [158]
Kamololol acetate	299	Ic	<i>F. pseudooreoselinum</i> Koso-Pol.	[121]
Kamololol acetate	278	Ic	<i>F. pseudalliacea</i> Rech.f.	[163]
Karatavic acid	216	Ib	<i>F. karatavica</i> Regel & Schmalh.	[164–166]
Karatavicin	163	Ia	<i>F. karatavica</i> Regel & Schmalh.	[148]
Karatavicinol	161	Ia	<i>F. foetida</i> (Bunge) Regel <i>F. asafoetida</i> L. <i>F. assa-foetida</i> L. <i>F. karatavica</i> Rgl. et Schmalh.	[83] [124] [10] [167]
Kellerin	321	Ic	<i>F. kelleri</i> K.-Pol. <i>F. kokanica</i> Regel & Schmalh.	[109] [110]
Kokanidin	306	Ic	<i>F. kokanica</i> Regel & Schmalh.	[110]
Kopeolin	197	Ib	<i>F. gummosa</i> Boiss. <i>F. kopetdagensis</i> Eug. Kor.	[168] [169,170]
Kopeolone	199	Ib	<i>F. kopetdagensis</i> Eug. Kor.	[169]

Table 2. Cont.

Names	No.	Types	Species	References
Kopeoside	198	Ib	<i>F. gummosa</i> Boiss. <i>F. kopetdagensis</i> Eug. Kor.	[168] [169,170]
Kopetdaghin	194	Ib	<i>F. gummosa</i> Boiss. <i>F. kopetdagensis</i> Eug. Kor.	[168] [169,171]
Lehmannolol	300	Ic	<i>F. sinkiangensis</i> K. M. Shen <i>F. assa-foetida</i> L.	[160] [10]
Lehmannolone	298	Ic	<i>F. fukanensis</i> K. M. Shen <i>F. sinkiangensis</i> K. M. Shen <i>F. lehmannii</i> Boiss.	[86] [160] [172]
Lehmannolone A	200	Ib	<i>F. lehmannii</i> Boss.	[87]
Lehmferidin	244	Ic	<i>F. lehmannii</i> Boss.	[173]
Lehmferin	191	Ib	<i>F. flabelliloba</i> Rech. f. & Aell <i>F. assa-foetida</i> L. <i>F. lehmannii</i> Boss.	[104] [10] [173]
Ligupersin A	273	Ic	<i>F. assa-foetida</i> L. <i>F. flabelliloba</i> Rech. f. & Aell <i>F. gummosa</i> Boiss. <i>F. badrakema</i> Kos.-Pol.	[10] [104] [97] [108]
Methyl galbanate	214	Ib	<i>F. szowitsiana</i> DC. <i>F. assa-foetida</i> L. <i>F. microloba</i> Boiss.	[122] [10] [161]
Microlobidene	335	Ic	<i>F. microloba</i> Boiss.	[174]
Microlobin	333	Ic	<i>F. assa-foetida</i> L.	[10,161]
Mogoltacin	266	Ic	<i>F. badrakema</i> Kos.-Pol.	[41,108]
Mogoltadone	227	Ic	<i>F. mogoltavica</i> Lipsky ex Korovin	[175]
Mogoltavicin	329	Ic	<i>F. mogoltavica</i> Lipsky ex Korovin	[176]
Mogoltavidin	328	Ic	<i>F. mogoltavica</i> Lipsky ex Korovin	[176]
Mogoltavin	280	Ic	<i>F. mogoltavica</i> Lipsky ex Korovin	[177]
Mogoltavinin	281	Ic	<i>F. mogoltavica</i> Lipsky ex Korovin	[177]
Moschatyl acetate	269	Ic	<i>F. incisoserrata</i> Pimenov & J.V.Baranova	[178]
Nevskone	296	Ic	<i>F. neveskii</i> Korovin	[99]
(E)- ω -Oxoferprenin	400	IIIc	<i>F. communis</i> L.	[78,79]
(E)- ω -Oxoferulenol	354	IIIa	<i>F. communis</i> L.	[80]
Pallidone A	359	IIIa	<i>F. pallida</i> Korovin	[179]
Pallidone B	362	IIIb	<i>F. pallida</i> Korovin	[179]
Persicaoside A	342	Ic	<i>F. persica</i> Willd	[180]
Persicaoside B	247	Ic	<i>F. persica</i> Willd	[180]
Persicaoside C	165	Ia	<i>F. persica</i> Willd	[180]
Persicaoside D	166	Ia	<i>F. persica</i> Willd	[180]
Polyanthin	229	Ic	<i>F. assa-foetida</i> L. <i>F. polyanthum</i> Eug. Korov.	[10] [181]
Polyanthinin	226	Ic	<i>F. polyantha</i> Korovin <i>F. polyanthum</i> Eug. Korov.	[99] [181]
(8'S,9'S,10'S)-Propionyl-fekrynol	182	Ib	<i>F. sinkiangensis</i> K. M. Shen	[146]
Reoselin	162	Ia	<i>F. kirialovii</i> Pimenov <i>F. korshinskyi</i> Eug. Korov <i>F. pseudooreoselinum</i> Koso-Pol.	[182] [136] [183,184]
Samarcandicin A	253	Ic	<i>F. samarkandica</i> Korovin	[96]
Samarcandicin B	254	Ic	<i>F. samarkandica</i> Korovin	[96]
Samarcandicin C	255	Ic	<i>F. samarkandica</i> Korovin	[96]
Samarcandicin D	256	Ic	<i>F. samarkandica</i> Korovin	[96]
Samarcandicin E	223	Ic	<i>F. samarkandica</i> Korovin	[96]
Samarcandicin F	257	Ic	<i>F. samarkandica</i> Korovin	[96]

Table 2. Cont.

Names	No.	Types	Species	References
Samarcandicin G	224	Ic	<i>F. samarkandica</i> Korovin	[96]
Samarcandicin H	258	Ic	<i>F. samarkandica</i> Korovin	[96]
Samarcandin	291	Ic	<i>F. samarcandica</i> kor. <i>F. teterrima</i> Kar. et Kir. <i>F. huber-morathii</i> Peşmen	[185] [117] [246]
Samarcandin acetate	290	Ic	<i>F. teterrima</i> Kar. et Kir. <i>F. pseudooreoselinum</i> Koso-Pol. <i>F. huber-morathii</i> Peşmen	[117] [186] [246]
Samarcandone	292	Ic	<i>F. samarcandica</i> kor. <i>F. sinaica</i> L. <i>F. huber-morathii</i> Peşmen	[185] [5] [246]
Saradaferin	261	Ic	<i>F. assafoetida</i> Linn.	[187]
Seravschanin A	274	Ic	<i>F. seravschanica</i> Pimenov & J.V.Baranova	[188]
Seravschanin B	275	Ic	<i>F. seravschanica</i> Pimenov & J.V.Baranova	[188]
Seravschanin C	276	Ic	<i>F. seravschanica</i> Pimenov & J.V.Baranova	[188]
Seravschanin D	170	Ia	<i>F. seravschanica</i> Pimenov & J.V.Baranova	[188]
Seravschanin E	157	Ia	<i>F. seravschanica</i> Pimenov & J.V.Baranova	[188]
Sinkiangenol A	220	Ic	<i>F. sinkiangensis</i> K. M. Shen	[189]
Sinkiangenol B	221	Ic	<i>F. sinkiangensis</i> K. M. Shen	[189]
Sinkiangenol C	176	Ib	<i>F. sinkiangensis</i> K. M. Shen	[189]
Sinkiangenol D	294	Ic	<i>F. sinkiangensis</i> K. M. Shen	[189]
Sinkiangenol E	222	Ic	<i>F. sinkiangensis</i> K. M. Shen	[189]
Sinkiangenorin D	219	Ib	<i>F. sinkiangensis</i> K. M. Shen	[128]
Sinkiangenorin E	344	Ic	<i>F. sinkiangensis</i> K. M. Shen	[30]
Sinkiangenorin F	208	Ib	<i>F. sinkiangensis</i> K. M. Shen	[81]
(3'S,8'R,9'S,10'R)-Sinkianol A	343	Ic	<i>F. sinkiangensis</i> K. M. Shen	[190]
(3'R,5'R,10'R)-Sinkianol B	210	Ib	<i>F. sinkiangensis</i> K. M. Shen	[190]
Sinkianone	201	Ib	<i>F. sinkiangensis</i> K. M. Shen <i>F. lehmannii</i> Boss.	[160] [87]
Sumferin	262	Ic	<i>F. sumbul</i> Hook	[191]
Szowitsiacoumarin A	316	Ic	<i>F. szowitsiana</i> DC.	[122]
Szowitsiacoumarin B	317	Ic	<i>F. szowitsiana</i> DC.	[122]
Tadzhiferin	153	Ia	<i>F. assa-foetida</i> L. <i>F. tadshikorum</i> M. Pimen	[77] [192]
Tadzhikorin	159	Ia	<i>F. tadshikorum</i> M. Pimen	[192]
Tavicone	337	Ic	<i>F. aitchisonii</i> K.-Pol.	[166]
Tunetacoumarin A	318	Ic	<i>F. tunetana</i> POM	[75]
Umbelliprenin	150	Ia	<i>F. aitchisonii</i> K.-Pol. <i>F. arrigonii</i> Bocchieri <i>F. assafoetida</i> Linn. <i>F. campestris</i> (Besser) Grecescu <i>F. fukanensis</i> K. M. Shen <i>F. sinkiangensis</i> K. M. Shen <i>F. flabelliloba</i> Rech. f. & Aell <i>F. tunetana</i> POM <i>F. persica</i> Willd <i>F. szowitsiana</i> DC.	[166] [43] [77,124] [100] [86] [193] [104] [75] [33] [122]
-	260	Ic	<i>F. sinaica</i> L.	[194]
-	336	Ic	<i>F. sinaica</i> L.	[68]




Figure 3. Cont.




Figure 3. Cont.




Figure 3. Cont.




Figure 3. Sesquiterpene coumarins in *Ferula* plants.

2.3. Sesquiterpene Chromones

Sesquiterpene chromones are present relatively rarely in the genus *Ferula*, with only 16 compounds (**408–423**) found in *F. communis* subsp. *communis*, *F. fukanensis*, *F. ferulaeoides*, *F. pallida*, and *F. sinkiangensis*, which can be classified into the furanochromone type (**408–418**) and pyranochromone type (**419–421**). The exceptions are ($\pm$)-ferulasin (**422** and **423**), which are uncommon sesquiterpene chromones with an oxygen-containing macrocyclic framework.


The names and sources of sesquiterpene chromones (**408–423**) are listed in Table 3, and their chemical structures are shown in Figure 4.

Table 3. Sesquiterpene chromones in *Ferula* plants.

Names	No.	Species	References
2,3-Dihydro-7-hydroxy-2 <i>S</i> *,3 <i>R</i> *-dimethyl-2-[4,8-dimethyl-3(<i>E</i>),7-nonadienyl]-furo[2,3- <i>b</i>]chromone	408	<i>F. ferulaeoides</i> (Steud.) Korov	[115]
2,3-Dihydro-7-hydroxy-2 <i>S</i> *,3 <i>R</i> *-dimethyl-2-[4-methyl-5-(4-methyl-2-furyl)-3(<i>E</i>),7-pentenyl]-furo[2,3- <i>b</i>]chromone	409	<i>F. ferulaeoides</i> (Steud.) Korov	[115]
2,3-Dihydro-7-hydroxy-2 <i>R</i> *,3 <i>R</i> *-dimethyl-2-[4-methyl-5-(4-methyl-2-furyl)-3(<i>E</i>),7-pentenyl]-furo[2,3- <i>b</i>]chromone	410	<i>F. ferulaeoides</i> (Steud.) Korov	[115]
Ferchromone	420	<i>F. communis</i> subsp. <i>communis</i>	[131]

Table 3. Cont.

Names	No.	Species	References
Ferchromonol	421	<i>F. communis</i> subsp. <i>communis</i>	[131]
(+)-Ferulasin	422	<i>F. sinkiangensis</i> K. M. Shen	[195]
(-)-Ferulasin	423	<i>F. sinkiangensis</i> K. M. Shen	[195]
Ferulin D	418	<i>F. ferulaeoides</i> (Steud.) Korov	[143]
Ferulin E	419	<i>F. ferulaeoides</i> (Steud.) Korov	[143]
Fukanefurochromone A	411	<i>F. fukanensis</i> K. M. Shen	[196]
Fukanefurochromone B	412	<i>F. fukanensis</i> K. M. Shen	[196]
Fukanefurochromone C	413	<i>F. fukanensis</i> K. M. Shen	[196]
Fukanefurochromone D	414	<i>F. fukanensis</i> K. M. Shen	[196]
Fukanefurochromone E	415	<i>F. fukanensis</i> K. M. Shen	[196]
Pallidone I	416	<i>F. pallida</i> Korovin	[197]
Pallidone J	417	<i>F. pallida</i> Korovin	[197]

Figure 4. Sesquiterpene chromones from *Ferula* plants.

2.4. Sesquiterpene Phenylpropanoids

Sesquiterpene phenylpropanoids (**424–458**) are also present only in a few *Ferula* species, such as *F. fukanensis*, *F. ferulaeoides*, *F. pallida*, *F. sinkiangensis*, and *F. seravschanica*. Their structures vary mainly in the sesquiterpene moiety, including the types and positions of substituents, stereoisomerism, etc., whereas the phenylpropanoid moiety often loses one carbon. The names and sources of compounds **424–458** are listed in Table 4, and their chemical structures are shown in Figure 5.

Table 4. Sesquiterpene phenylpropanoids in *Ferula* plants.

Names	No.	Species	References
3-(2,4-dihydroxybenzoyl)-4 <i>R</i> *,5 <i>R</i> *-dimethyl-5-[4,8-dimethyl-3(<i>E</i>),7(<i>E</i>)-nonadien-1-yl]tetra-hydro-2-furanone	443	<i>F. ferulaeoides</i> (Steud.) Korov.	[198,199]
3 <i>S</i> *-(2,4-dihydroxybenzoyl)-4 <i>R</i> *,5 <i>R</i> *-dimethyl-5-[4-methyl-5-(4-methyl-2-furyl)-3(<i>E</i>)-penten-1-yl]tetrahydro-2-furanone	450	<i>F. ferulaeoides</i> (Steud.) Korov.	[199]
3 <i>S</i> *-(2,4-dihydroxybenzoyl)-4 <i>R</i> *,5 <i>S</i> *-dimethyl-5-[4-methyl-5-(4-methyl-2-furyl)-3(<i>E</i>)-penten-1-yl]tetrahydro-2-furanone	451	<i>F. ferulaeoides</i> (Steud.) Korov.	[199]
8,9-Dihydroxydshamirone	455	<i>F. ferulaeoides</i> (Steud.) Korov.	[200]
(4 <i>E</i> ,8 <i>E</i>)-1-(2,4-dihydroxyphenyl)-2-hydroxy-5,9,13-trimethyltetradeca-4,8,12-trien-1-one	438	<i>F. ferulaeoides</i> (Steud.) Korov.	[198,201]
1-(2,4-dihydroxyphenyl)-3,7,11-trimethyl-3-vinyl-6(<i>E</i>),10-dodecadiene-1,9-dione	440	<i>F. ferulaeoides</i> (Steud.) Korov.	[201]
(6 <i>E</i>)-1-(2,4-dihydroxyphenyl)-3,7,11-trimethyl-3-vinyl-6,10-dodecadien-1-one	439	<i>F. ferulaeoides</i> (Steud.) Korov.	[198,201]
(6 <i>E</i>)-1-(2,4-dihydroxyphenyl)-3,7-dimethyl-3-vinyl-8-(4-methyl-2-furyl)-6-octen-1-one	441	<i>F. ferulaeoides</i> (Steud.) Korov.	[198,201]
Dshamirone	437	<i>F. ferulaeoides</i> (Steud.) Korov.	[198,201]
Ferulaeolactone A	452	<i>F. ferulaeoides</i> (Steud.) Korov.	[200]
Ferulaeone A	429	<i>F. ferulaeoides</i> (Steud.) Korov.	[198]
Ferulaeone B	430	<i>F. ferulaeoides</i> (Steud.) Korov.	[198]
Ferulaeone C	431	<i>F. ferulaeoides</i> (Steud.) Korov.	[198]
Ferulaeone D	432	<i>F. ferulaeoides</i> (Steud.) Korov.	[198]
Ferulaeone E	433	<i>F. ferulaeoides</i> (Steud.) Korov.	[198]
Ferulaeone F	434	<i>F. ferulaeoides</i> (Steud.) Korov.	[198]
Ferulaeone G	435	<i>F. ferulaeoides</i> (Steud.) Korov. <i>F. sinkiangensis</i> K. M. Shen	[198] [21]
Ferulaeone H	436	<i>F. ferulaeoides</i> (Steud.) Korov.	[198]
Fukanedone A	424	<i>F. fukanensis</i> K. M. Shen	[18]
Fukanedone B	425	<i>F. fukanensis</i> K. M. Shen <i>F. ferulaeoides</i> (Steud.) Korov.	[18] [198,200]
Fukanedone C	426	<i>F. fukanensis</i> K. M. Shen	[18]
Fukanedone D	427	<i>F. fukanensis</i> K. M. Shen	[18]
Fukanedone E	428	<i>F. fukanensis</i> K. M. Shen	[18]
3-(2-hydroxyl-4-methoxybenzoyl)-4 <i>S</i> *,5 <i>R</i> *-dimethyl-5-[4,8-dimethyl-3(<i>E</i>),7(<i>E</i>)-nonadien-1-yl]tetrahydro-2-furanone	442	<i>F. ferulaeoides</i> (Steud.) Korov.	[198]
8,9-Oxoisopropanyldshamirone	454	<i>F. ferulaeoides</i> (Steud.) Korov.	[200]
Pallidone C	444	<i>F. pallida</i> Korovin	[179]
Pallidone D	445	<i>F. pallida</i> Korovin	[179]
Pallidone E	446	<i>F. pallida</i> Korovin	[179]
Pallidone F	447	<i>F. pallida</i> Korovin	[179]
Sinkiangenone A	448	<i>F. sinkiangensis</i> K. M. Shen	[21]
Sinkiangenone B	449	<i>F. sinkiangensis</i> K. M. Shen	[21]




Figure 5. Sesquiterpene phenylpropanoids in *Ferula* plants.

3. Biosynthetic Pathways

Sesquiterpenes are synthesized in plants through complex biosynthetic pathways, which involve several enzymatic reactions and intermediates. Sesquiterpene biosynthesis typically begins with the isopentenyl diphosphate (IPP) and dimethylallyl diphosphate (DMAPP) precursors, which are common to all terpenoids. These precursors are generated through the mevalonate (MVA) pathway or the 2-C-methyl-D-erythritol 4-phosphate (MEP) pathway. Once IPP and DMAPP are synthesized, they serve as the building blocks for sesquiterpene biosynthesis. IPP and DMAPP are condensed to form geranyl diphosphate

(GPP), which contains ten carbon atoms. Then farnesyl diphosphate (FPP), which contains fifteen carbon atoms, is formed by the condensation of two molecules of IPP and GPP. FPP is a precursor to various sesquiterpenes, and it undergoes further modifications and cyclization reactions. These cyclization reactions create diverse sesquiterpene skeletons with different ring structures. After the initial cyclization step, the sesquiterpene skeleton may undergo rearrangement or further modification by various enzymes. This step introduces functional groups and structural diversity into the sesquiterpenes. After the sesquiterpenes are synthesized, they may undergo additional enzymatic modifications, such as glycosylation, acylation, or oxidation. These modifications can alter their solubility, stability, and biological activities. The biosynthesis pathways of the typical sesquiterpene skeletons in *Ferula* are shown in Figure 6.




Figure 6. Biosynthesis pathways of the typical sesquiterpene skeletons in *Ferula* [202,203].

The scaffold of sesquiterpene coumarins is formed through the dehydration condensation of the coumarin unit and the sesquiterpene moiety via an ether bond. The biosynthesis pathways of type Ib sesquiterpene coumarins, e.g., **219** [128], type Ic sesquiterpene coumarins, e.g., **225**, **319**, **323**, and **338** [112,204]; type IIIa sesquiterpene coumarins, e.g., **355** and **359** [143,179]; and type IIIb sesquiterpene coumarins, e.g., **362**, **370**, **372**, **374**, **375**, **383**, **384**, **386**, **387**, **392**, **394**, and **395** [143,179], are shown in Figure 7.




Figure 7. Biosynthesis pathways of different types of sesquiterpene coumarins. **(A)** The biosynthesis pathways of type I sesquiterpene coumarins [112,128,204]. **(B)** The biosynthesis pathways of type III sesquiterpene coumarins [143,179]. The biosynthesis pathways of sesquiterpene chromones [115,143,195] and sesquiterpene phenylpropanoids [179,198,199,201] are shown in Figures 8 and 9, respectively.




Figure 8. Proposed biosynthetic pathways for sesquiterpene chromones [115,143,195].

4. Bioactive Properties

4.1. Antibacterial Effects

Antibiotics, such as penicillin, tetracycline, ciprofloxacin, are a common class of antibacterial agents, which are specifically designed to target and kill or inhibit the growth of bacteria. Natural compounds can be used alongside antibiotics as complementary treatments, potentially enhancing the overall effectiveness of the treatment.

The antibacterial effects of sesquiterpenes and sesquiterpene derivatives from *Ferula* have been extensively studied. In 2001, Tamemoto et al. [64] isolated the daucane sesquiterpene ferutinin (**18**) from *F. kuhistanica*, which showed strong activity against Gram-positive bacteria, including *Staphylococcus aureus*, *S. epidermidis*, *Enterococcus faecalis*, and *Bacillus subtilis*, but did not exhibit activity against Gram-negative bacteria. Among them, the MIC values of ferutinin (**18**) were similar to those of the standard antibiotics ampicillin and

chloramphenicol. In another study, Ibraheim et al. [205] reported that ferutinin (18), teferin (21), and teferidin (17) from *F. hermonis* showed strong activity against methicillin-resistant *S. aureus* (MRSA) (MIC, <0.39, 1.56, and 0.78 µg/mL, respectively), *B. subtilis* (MIC: <0.39, 1.56, and <0.39 µg/mL, respectively), *Mycobacterium tuberculosis* (MIC: 2, 8, and 0.69 µg/mL, respectively), and BCG (MIC: 1.56, 6.25, and 3.125 µg/mL, respectively). In addition, the enhancing effect of ferutinin (18) on four antitubercular drugs, i.e., rifampin, isoniazid, streptomycin, and ethionamide was tested against *M. smegmatis*. Ferutinin (18) showed higher activity (MIC, 10 µg/mL) than rifampin and streptomycin (MIC: each at 20 µg/mL) in the agar dilution assay, and the combination of ferutinin (18) with the anti-tuberculosis drugs isoniazid and ethionamide resulted in enhancing the effect of the antimycobacterial activity in the checkerboard method, while the combination with rifampicin or streptomycin did not exhibit this effect [206].

As another structural type of sesquiterpenes, the new guaianolides diversolide A (103), diversolide D (106), and diversolide F (108) from *F. diversivittata* showed moderate antibacterial activities against *S. aureus* (ATCC 29737) and *Escherichia coli* (ATCC 8739), with MIC values ranging from 40 to 80 µg/mL, while the MIC values of the gentamycin positive control were 18 µg/mL and 8 µg/mL, respectively [49].

In 2004, the sesquiterpene coumarin ferulenol (349) and its three derivatives were investigated for their antimycobacterial activity against four strains of fast-growing *Mycobacterium* species: *M. fortuitum*, *M. phlei*, *M. aurum*, and *M. smegmatis*. Ferulenol (349) showed the most promising activity with IC₅₀ values of 0.5–2 µg/mL, which was superior to the therapeutically used antimycobacterials isoniazid and ethambutol (IC₅₀: 0.5–8 µg/mL and 0.5–4 µg/mL, respectively) [207]. Ferulenol (349) showed stronger activity (MIC: each at 0.63 µg/mL) against the above Gram-positive bacteria, and it exhibited potent activity against *Mycobacterium* organisms (MIC: each at 1.25 µg/mL). In contrast, ferchromone (420) proved to be less active [208].

By the disk diffusion method, umbelliprenin (150) was found to have the highest activity against *B. subtilis*, *B. cereus*, *E. coli*, *Salmonella typhi*, *Klebsiella pneumoniae*, *S. aureus*, and *S. epidermidis* at a concentration of 500 µg/mL [209]. Liu et al. [116,200] discovered a series of sesquiterpene phenylpropanoids and sesquiterpene coumarins from *F. ferulioides* with strong antibacterial properties against multidrug-resistant (MDR) *S. aureus* strains, including ATCC25923, RN4220, SA1199B, XU212, EMRSA15, and EMRSA16. Among them, 8,9-oxoisopropanyldshamirone (454) and ferulaeolactone A (452) displayed better antibacterial properties than some of the controls in the above strains, i.e., norfloxacin, tetracycline, erythromycin, and oxacillin. The sesquiterpene phenylpropanoid (6E)-1-(2,4-dihydroxyphenyl)-3,7,11-trimethyl-3-vinyl-6,10-dodecadien-1-one (439) and the sesquiterpene coumarin 2,3-Dihydro-7-hydroxy-2S*,3R*-dimethyl-2-[4,8-dimethyl-3(E),7-nonadienyl]-furo[3,2-c]coumarin (383) from *F. heuffelii* could significantly inhibit *S. epidermidis* (MIC: 11.2 and 5.2 µM, respectively) and *Micrococcus luteus* (MIC: 22.5 and 10.5 µM, respectively) growth. Moreover, (6E)-1-(2,4-dihydroxyphenyl)-3,7,11-trimethyl-3-vinyl-6,10-dodecadien-1-one (439) could also inhibit *B. subtilis* growth, with an MIC value of 11.2 µM [210]. Sun et al. [211] discovered two sesquiterpene phenylpropanoids from *F. ferulioides* using TLC-bioautography-directed isolation. Both compounds showed significant antibacterial activities against five tested strains, especially the MDR strains XU212 and SA1199B, and the MIC values of the two compounds (16 µg/mL and 1 µg/mL, 8 µg/mL and 2 µg/mL, respectively) were lower than the standard antibiotic norfloxacin (32 µg/mL and 8 µg/mL, respectively). Ferulsinaic acid (218) was found to have strong antibacterial activity against Gram-positive strains (*B. cereus* and *S. aureus*) and Gram-negative strains (*Serratia* sp., *Pseudomonas* sp., and *E. coli*), which was comparable to the reference antibiotics ampicillin and amoxillin [145].




Figure 9. Proposed biosynthetic pathways for sesquiterpene phenylpropanoids [179,198,199,201].

In 2007, Shahverdi et al. [212] proved that galbanic acid (**213**) could enhance the antibacterial activity of penicillin G and cephalexin against *S. aureus*. The MIC of penicillin G alone was 64 µg/mL, while the MIC of a combination of penicillin and galbanic acid (**213**) was reduced to 1 µg/mL. In the meanwhile, the MIC of cephalexin decreased from 64 µg/mL to 1 µg/mL when used in combination with galbanic acid (**213**). In 2009, Bazzaz et al. [213] proved that galbanic acid (**213**) could enhance the activity of methicillin, tetracycline, and ciprofloxacin against isolates of *S. aureus*. The MIC of methicillin, tetracycline, and ciprofloxacin decreased from 10–80 µg/mL, 40–80 µg/mL, and 10–20 µg/mL to less than 1.25 µg/mL when used in combination with galbanic acid (**213**). The class A β -lactamase is one of the main causes of β -lactam antibiotic resistance. Umbelliprenin (**150**) and galbanic acid (**213**) showed potent inhibitory activity (IC_{50} : 54 ± 2.9 µM and 47 ± 3.1 µM, respectively) against class A β -lactamase, and the IC_{50} of the positive control, clavulanic acid, was 24.1 ± 2.1 µM. Moreover, the average MIC of penicillin G alone was 244.2 ± 12.3 µM, while the average MIC of penicillin–umbelliprenin and penicillin–galbanic acid were 21.3 ± 4.3 µM and 18.2 ± 5.6 µM, respectively, which was a significant decrease from the MIC of penicillin G. The results indicate that umbelliprenin (**150**) and galbanic acid (**213**) may be good substitutes for clavulanic acid to combat infections caused by *S. aureus* resistance [214]. Galbanic acid (**213**) appears to exert its antibacterial activity by the regulation of drug resistance.

In 2014, Dastan et al. [158] reported that 4'-hydroxy kamolanol acetate (**277**) and kamolanol (**334**) from *F. pseudalliacea* displayed antibacterial activity against *Helicobacter pylori* and *S. aureus* (MIC: each at 64 µg/mL). Later in 2016, they investigated the antibacterial effect of another six sesquiterpene coumarins from *F. pseudalliacea* against seven bacterial strains, including *S. aureus*, *B. cereus*, *E. faecium* (vancomycin-resistant clinical strain), *K. pneumonia* (clinical strain), *P. aeruginosa* (clinical strain), *Helicobacter pylori*, and *E. coli*. All compounds were effective (MIC: 64–128 µg/mL) against *S. aureus*, except for farnesiferol B (**186**). Fekrynol acetate (**212**) and methyl galbanate (**214**) showed significant activity against *E. faecium*, with MIC values of 128 and 64 µg/mL, respectively, while the MIC value

of the chloramphenicol control was 32 µg/mL. Ethyl galbanate (**215**) and kamolonol acetate (**278**) showed strong activity against *H. pylori*, with MIC values of 64 and 128 µg/mL, respectively [215]. The MIC values of these compounds against different bacterial strains are listed in Table 5.

Table 5. The MIC values of sesquiterpenes and sesquiterpene derivatives against different bacterial strains.

Names	No.	Bacterial Strains	MIC	References
2,3-Dihydro-7-hydroxy-2S*,3R*-dimethyl-2-[4,8-dimethyl-3(E),7-nonadienyl]-furo[3,2-c]coumarin	383	<i>S. epidermidis</i>	5.2 µM	[210]
		<i>M. luteus</i>	22.5 µM	[210]
(6E)-1-(2,4-dihydroxyphenyl)-3,7,11-trimethyl-3-vinyl-6,10-dodecadien-1-one	439	<i>S. epidermidis</i>	11.2 µM	[210]
		<i>M. luteus</i>	22.5 µM	[210]
		<i>B. subtilis</i>	11.2 µM	[210]
Diversolide A	103	<i>S. aureus</i>	>160 µg/mL	[49]
		<i>E. coli</i>	80 µg/mL	[49]
Diversolide D	106	<i>S. aureus</i>	40 µg/mL	[49]
		<i>E. coli</i>	>160 µg/mL	[49]
Diversolide F	108	<i>S. aureus</i>	80 µg/mL	[49]
		<i>E. coli</i>	80 µg/mL	[49]
Ethyl galbanate	215	<i>H. pylori</i>	64 µg/mL	[215]
Fekryinol acetate	212	<i>E. faecium</i>	128 µg/mL	[215]
Ferulenol	349	<i>M. fortuitum</i>	2 µg/mL	[207]
		<i>M. phlei</i>	2 µg/mL	[207]
		<i>M. aurum</i>	2 µg/mL	[207]
		<i>M. smegmatis</i>	0.5 µg/mL	[207]
		<i>B. subtilis</i>	0.63 µg/mL	[208]
		<i>S. aureus</i>	0.63 µg/mL	[208]
		<i>S. durans</i>	0.63 µg/mL	[208]
		<i>S. faecalis</i>	0.63 µg/mL	[208]
Ferutinin	18	<i>Mycobacterium</i> organisms	1.25 µg/mL	[208]
		MRSA	<0.39 µg/mL	[205]
		<i>B. subtilis</i>	<0.39 µg/mL	[205]
		MTB	2 µg/mL	[205]
		BCG	1.56 µg/mL	[205]
Galbanic acid	213	<i>M. smegmatis</i>	10 µg/mL	[206]
		class A β-lactamase	47 ± 3.1 µM	[214]
4'-Hydroxy kamolonol acetate	277	<i>H. pylori</i>	64 µg/mL	[158]
		<i>S. aureus</i>	64 µg/mL	[132]
Kamolonol	334	<i>H. pylori</i>	64 µg/mL	[158]
		<i>S. aureus</i>	64 µg/mL	[132]
Kamolonol acetate	278	<i>H. pylori</i>	128 µg/mL	[215]
Methyl galbanate	214	<i>E. faecium</i>	64 µg/mL	[215]
Teferidin	17	<i>S. aureus</i>	0.78 µg/mL	[205]
		<i>B. subtilis</i>	<0.39 µg/mL	[205]
		MTB	0.69 µg/mL	[205]
		BCG	3.125 µg/mL	[205]
Teferin	21	MRSA	1.56 µg/mL	[205]
		<i>B. subtilis</i>	1.56 µg/mL	[205]
		MTB	8 µg/mL	[205]
		BCG	6.25 µg/mL	[205]
Umbelliprenin	150	<i>B. subtilis</i>	500 µg/mL	[209]
		<i>B. cereus</i>	500 µg/mL	[209]
		<i>E. coli</i>	500 µg/mL	[209]
		<i>S. typhi</i>	500 µg/mL	[209]
		<i>K. pneumoniae</i>	500 µg/mL	[209]
		<i>S. aureus</i>	500 µg/mL	[209]
		<i>S. epidermidis</i>	500 µg/mL	[209]
		Class A β-lactamase	54 ± 2.9 µM	[214]

4.2. Antifungal Effects

Over the past several decades, there has been a significant rise in the number of human fungal infections, particularly those affecting the skin and mucosal surfaces. These infections are most common in tropical and subtropical regions and are mostly caused by *Candida* sp. and dermatophytes [216]. According to research conducted by Al-Ja'fari et al. [12], ferutin (18) and teferidin (17) from the rhizome and roots of *F. hermonis* displayed antifungal activity in vitro. The results of determining the minimal fungicidal concentration (MFC) and MIC of both substances showed that ferutin (18) had greater antifungal activity than teferidin (17). Especially in *Tricophyton mentagrophytes*, their MIC and MFC values ranged from 8 to 256 mg/mL.

4.3. Antiparasitic Effects

Iranshahi et al. [122] and Bashir et al. [151] evaluated the inhibitory activity of sesquiterpene coumarins extracted from *F. szowitsiana* and *F. narthex* Boiss against *Leishmania major*. The results showed that umbelliprenin (150) and conferol (267) displayed potent antileishmanial activity (IC₅₀: 4.9 and 11.5 µg/mL, respectively). Dastan et al. [121] determined the in vitro antiparasitic activity of compounds extracted from *F. pseudalliacea* against the *Plasmodium falciparum* strain, K1. The results indicated that kamolonol acetate (299) and methyl galbanate (214) showed moderate antiparasitic activity, with IC₅₀ values of 16.1 and 7.1 µM, respectively, whereas the IC₅₀ value of the positive control, artemisin, was 0.004 µM.

4.4. Antiviral Effects

H1N1, also known as swine flu, is a subtype of the influenza A virus. In 2009, it attracted worldwide attention when a new H1N1 strain emerged and caused a global pandemic. It was found that some sesquiterpene coumarins of *Ferula* species were active against H1N1. Lee et al. [10] discovered that 5'-hydroxyumbelliprenin (151), 8'-acetoxy-5'-hydroxyumbelliprenin (155), methyl galbanate (214), galbanic acid (213), farnesiferol C (202), farnesiferol A (228), conferol (267), ligupersin A (273), and *epi*-conferdione (272) isolated from *F. assa-foetida* displayed significant antiviral activity against H1N1 (IC₅₀: 0.26–0.86 µg/mL), which was more effective than amantadine (IC₅₀: 0.92 µg/mL). Li et al. [30] found that Sinkiangenorin E also had a significant inhibitory effect on H1N1. These findings indicate that sesquiterpene coumarins might be potential lead compounds for new drugs to treat H1N1 viral infection. In addition, the sesquiterpene coumarin kellerin (321) showed an antiviral effect against herpes virus type 1 (HSV-1) by the plaque-reduction assay. It could dramatically reduce the viral titre of the HSV-1 DNA viral strain KOS at concentrations of 10, 5, and 2.5 µg/mL and considerably lessen its cytopathic effects [217].

4.5. Antioxidative Effects

Oxidative stress refers to the imbalance between the antioxidative defense system and the production of oxidants (free radicals). The accumulation of oxidized lipids plays an important role in the incidence of many diseases such as diabetes, cancer, aging, etc. Therefore, compounds that reduce or prevent the production of oxidative products can be used to treat these diseases [218].

In a study conducted by Raafat and El-Lakani [219], it was observed that the administration of ferutin (18), a daucane-type sesquiterpene ester, significantly reversed the decreasing trend of the expression of the antioxidant enzyme catalase observed in diabetic mice. In addition, for the first time, it described the antioxidant property of ferutin (18) on diabetes-related neuropathic pain, indicating that 1.6 mg/kg of ferutin (18) could reduce thermal hyperalgesia and tactile allodynia. At 500 and 1000 µg/kg mice body weight, ferutin (18) could significantly upregulate the gene expression of superoxide dismutase (SOD), catalase (CAT), and glutathione peroxidase (GPx) in liver and kidney tissues, which are known to resist cellular oxidative stress. At the same concentration, it could also significantly decrease the lipid peroxidation in mice liver tissues [220]. An analog of ferutin (18), 2 α -acetyl ferutin (19) could rapidly reduce the mRNA levels of several intracellular antioxidative enzymes, such as catalase, Mn-superoxide dismutase (SOD2),

nuclear factor erythroid 2-related factor 2 (NRF2), peroxiredoxin (PRDX1), and thioredoxin (TRX) between 6 and 12 h, and it could also significantly induce intracellular glutathione (GSH) depletion in a time- and concentration-dependent manner [32]. In addition, the daucane esters teferidin (17), ferutinin (18), and teferin (21) from *F. hermonis* showed strong 2,2-diphenyl-1-(2,4,6-trinitrophenyl)hydrazyl (DPPH) radical scavenging activity, with IC₅₀ values of 17.3, 13.2, and 11.5 µM, respectively, comparable to the positive control of ascorbic acid (12.5 µM). The significant increase in free radical scavenging activity is associated with an increase in the number of hydroxyl groups [205].

Umbelliprenin (150) is the first synthesized sesquiterpene coumarin in *Ferula*. In a study on the antigenotoxicity effects of umbelliprenin (150) on human peripheral lymphocytes exposed to oxidative stress [221], although umbelliprenin (150) showed no scavenging activity (4%), the protective activity of umbelliprenin (150) (10–400 mM) against DNA damage induced by 25 mM H₂O₂ increased in a concentration-dependent manner. There was no significant difference between umbelliprenin (150) and ascorbic acid (positive standard) when the concentration exceeded 50 mM. Kamolonol acetate (299) is also a sesquiterpene coumarin extracted from *F. pseudalliacea* with potent antioxidant activity. It displays strong DPPH radical scavenging activity, with an EC₅₀ value of 65.29 ± 5.6 µM, which is similar to that of the positive control, butylatedhydroxyanisole (BHA), at 59.85 ± 3.7 µM [222].

Kogure et al. [223] evaluated the antioxidative activities of several compounds isolated from *F. penninervis* and *F. pallida*, with the sesquiterpene coumarin KT23 (Pallidone A) (359) having moderate antioxidative properties. Compared with α-tocopherol (43.2%, 200 µM) as a control, KT23 (359) (100 µM) inhibited 16.4% of egg-yolk phosphatidylcholine liposome (EyPC liposome) peroxidation. Ferulsinaic acid (218) is a sesquiterpene coumarin from *F. sinaica* with a rare carbon skeleton. It was found to significantly reduce malondialdehyde (MDA) levels in *Caenorhabditis elegans*, thus attenuating lipid peroxidation. In addition, it could significantly decrease the formation of N-ε-carboxymethyllysine (CML), one of the advanced glycation end-products (AGEs) that is correlated with oxidative stress. These indicate the antioxidative power of ferulsinaic acid (218) [224]. Galbanic acid (213) is also a natural sesquiterpene coumarin abundantly distributed in *Ferula* species; it exhibited antioxidative activity by inhibiting DPPH and ABTS free radicals, with IC₅₀ values of 180 and 60 µg/mL, respectively. In addition, galbanic acid (213) (62.5 µg/mL) and vitamin C (5 µg/mL), as a positive control, could significantly upregulate the expression of SOD, CAT, and GPx. The upregulation of these antioxidative genes enhances the redox state of cells; however, the potential of galbanic acid (213) to upregulate antioxidative enzymes is lower than that of vitamin C [225].

4.6. Anti-Inflammatory Effects

Inflammation is a complex biological response triggered by the immune system in response to harmful stimuli such as infections, injuries, or diseases. Many sesquiterpenes and their derivatives from *Ferula* have anti-inflammatory properties, making them valuable for promoting overall health and potentially reducing the risk of chronic diseases associated with inflammation.

Ferutinin (18) and teferin (21) exhibit anti-inflammatory effects at a dose of 100 mg/kg using the in vivo carrageenan-induced edema model, which may be caused by the antagonistic effects of histamine and/or serotonin actions, and their anti-inflammatory effects may be directly related to the degree of oxidation of the benzene ring [226].

The sesquiterpene coumarins methyl galbanate (214) and umbelliprenin (150) were reported to exert their anti-inflammatory effects by significantly inhibiting the LPS-induced production of nitric oxide (NO) and prostaglandin E₂ (PGE₂), leading to a decrease in the expression of inducible nitric oxide synthase (iNOS) and cyclooxygenase-2 (COX-2) [227]. In another study, the anti-inflammatory effect of umbelliprenin (150) was evaluated in vitro and in vivo. It displayed a significant inhibitory effect on soybean lipoxygenase (a key enzyme in the process of inflammation), with an IC₅₀ value of 0.0725 µM, whereas the IC₅₀ value of the positive control, caffeic acid, was 600 µM. Furthermore, it showed a significant

anti-inflammatory effect (39%) in vivo, which was comparable to the positive control, indomethacin (47%), using the carrageenin mouse-paw edema model [228]. In RAW264.7 cells stimulated by lipopolysaccharide (LPS)/interferon- γ (IFN- γ), Kohno et al. [229] found that methyl galbanate (**214**) significantly reduced NO production. In the presence of methyl galbanate (**214**), the mRNA expression of iNOS stimulated by LPS/IFN- γ was reduced to 52% of the levels found with LPS/IFN- γ induction alone.

Kellerin (**321**) is the major constituent (1.5%, *w/w*) of *F. sinkiangensis*, and its anti-inflammatory mechanism is to inhibit the mRNA expression of inflammatory cytokines such as NO, tumor necrosis factor- α (TNF- α), COX-2, interleukin-6 (IL-6), and interleukin-1 β (IL-1 β) [190]. Zhang et al. [190] found that kellerin (**321**) could transform microglia from a pro-inflammatory M1 phenotype into an anti-inflammatory M2 phenotype, and thus alleviate cognitive impairment in mice. In another research, kellerin (**321**) was found to decrease the levels of pro-inflammatory cytokines, inhibit the NF- κ B signaling pathway, and reduce ROS production and NADPH oxidase activity to exert neuroprotective effects [27].

Motai et al. reported six new sesquiterpene coumarins from *F. fukanensis*, four of which (fukanefuomarin H–K) (**385**, **388**, **396**, and **397**) showed NO-inhibitory activities, with IC₅₀ values of 11.1–55.6 μ M. In addition, fukanefuomarin H (**385**) and fukanefuomarin K (**397**) could inhibit the gene expression of iNOS, IL-6, and TNF- α [156]. The sesquiterpene coumarins ferubungeanol B (**308**) (IC₅₀ 23.6 μ M) and samarcandin acetate (**290**) (IC₅₀ 25.6 μ M) from *F. bungeana* were found to have a strong inhibitory effect on NO production in LPS-induced BV-2 microglia compared with the positive control, minocycline (IC₅₀ 25.6 μ M) [137]. Episamarcandin acetate (**295**) from *Ferula sinkiangensis* was also found to exert its anti-inflammatory activity by significantly decreasing NO production, with an IC₅₀ value of 2.3 μ M, and inhibiting TNF- α , IL-1 β , and IL-6 expression [149].

4.7. Antitumor Effects

In the past few decades, the cytotoxicity of sesquiterpenes and sesquiterpene derivatives from the *Ferula* species, especially sesquiterpene coumarins, has been studied extensively. These compounds have shown significant cytotoxicity against various tumor cell lines, including HCT116 and HT-29 human colon cancer cells; AGS, BGC-823, and MGC-803 human gastric cancer cells; M4Beu human melanoma cells; BxPC3, PANC-1, and Capan-1 human pancreatic cancer cells; HeLa human cervical cancer cells; and MCF-7 and MDA-MB-231 human breast cancer cells. The IC₅₀ values of these compounds against different cancer cell lines in vitro are listed in Table 6.

Umbelliprenin (**150**) is one of the most widely studied sesquiterpene coumarins with antitumor potential. Promoting tumor cell apoptosis is one of the important mechanisms in antitumor therapy. Researchers discovered that umbelliprenin (**150**) could promote apoptosis in tumor cells by annexin V-FITC/PI staining. In the meanwhile, umbelliprenin (**150**) activated caspase-3, -8, and -9 and the proapoptotic protein Bax and reduced the expression of the antiapoptotic protein Bcl-2, caspase-3, -8, and -9, and the proapoptotic protein Bax and reduced the expression of the antiapoptotic protein Bcl2 [230,231], which promoted apoptosis in the Jurkat T-CLL and Raji B-CLL cell lines in a time- and dose-dependent manner [232]. In addition, it could activate the mitochondrial apoptotic pathway and lead to apoptosis of the cancer cells by decreasing the mitochondrial membrane potential, enhancing the P53, P27, P16, and Rb protein expression and diminishing the expression of the proteins of cyclin E, cyclin D, Cdk4, and Cdk6 as well as cell cycle arrest in the G0/G1 phase [233]. Apart from this, umbelliprenin (**150**) could attenuate cell migration through the Wnt signaling pathway by decreasing the expression levels of Wnt-2, β -catenin, GSK-3 β , p-GSK-3 β , survivin, and c-myc [193]. In another study, umbelliprenin was found to induce cytoprotective autophagy by reducing the phosphorylation levels of AKT and mTOR and blocking the Akt signaling pathway [230]. In brief, umbelliprenin (**150**) could exert its antitumor property by inducing apoptosis and autophagy, inhibiting the cell cycle, and attenuating the migration and invasion of cancer cells. In an in vivo study, a double-stage carcinogenicity assay of mouse skin tumors was performed to investigate

the cancer chemopreventive activity of umbelliprenin (150). The results showed that mice treated with umbelliprenin (150) together with peroxyntirite (initiator)/TPA (promoter) had delayed papillary tumor formation, with effects comparable to those of the curcumin control. Furthermore, the tumor development pattern was slower in umbelliprenin-treated mice compared with curcumin treatment. Thus, umbelliprenin (150) may be a potential cancer chemopreventive agent [234].

Galbanic acid (213) is another extensively studied sesquiterpene coumarin. Kim et al. [235] revealed the potential molecular mechanism of galbanic acid (213) in overcoming chemotherapy resistance in drug-resistant lung cancer. As an effective TNF-related apoptosis-inducing ligand (TRAIL) sensitizer, galbanic acid (213) enhanced TRAIL-induced cell apoptosis by inhibiting multidrug resistance 1 (MDR1) and activating caspase and death receptor 5 (DR5) in cisplatin-resistant H460/R non-small-cell lung cancer cells. Galbanic acid (213) induced tumor cell-cycle arrest at G₁, which is associated with the inhibition of the cyclin/cyclin-dependent kinase (CDK)4/6 pathway, particularly cyclin D₁. [236]. It also inhibited tumor cell metastasis. Neoangiogenesis and the activation of matrix metalloproteinases (MMPs) play a crucial role in tumor generation and metastasis. Neovessels are formed during tumor generation and metastasis [237], so the inhibition of angiogenesis may promote cancer cell death [238]. Kim et al. [239] reported that galbanic acid (213) reduced the number of blood vessels in tumor cells by more than 40%, significantly reduced the proliferation of vascular endothelial growth factor-(VEGF)-induced human umbilical-vein endothelial cells (HUVECs), and inhibited VEGF-induced migration and tube formation in HUVECs. It was shown to have an inhibitory effect on tumor-induced angiogenesis. MMPs are capable of degrading the vast majority of proteins in the extracellular matrix and disrupting the extracellular matrix and basement membrane barriers of tissues, which play a crucial role in the invasive and metastatic process of cancer cells [240]. Thus, inhibiting the activity of MMPs is an effective strategy to block the migration of tumor cells. Studies have demonstrated that galbanic acid (213) can inhibit the activity and expression of MMP2 and MMP9 [241]. Hypoxia-inducible factor (HIF) is a transcription factor that regulates the expression of genes involved in the regulation of hypoxic mechanisms (e.g., angiogenesis or apoptosis) as well as tumor growth, invasion, and metastasis [242]. Hypoxia in tumors can stimulate and induce HIF-1 α and HIF-2 α protein expression [243]. EGFR-MAPK is an important signaling pathway with regulatory effects on HIF-1 α expression [244]. Syeda et al. [245] found that galbanic acid (213) downregulated HIF-1 α and HIF-1 β mRNA expression under both hypoxic and normoxic conditions, and it had an inhibitory effect on HIF-1 activation. Under normoxic conditions, it shortened the half-life of the EGFR (HIF-1 downstream genes) and promoted EGFR degradation to inhibit HIF activation. Meantime, it inhibited HIF-1 α accumulation in A549 and OVCAR-3 cells by suppressing the EGFR/HIF-1 α signaling pathway [244].

Table 6. The IC₅₀ values of sesquiterpenes and sesquiterpene derivatives against different cancer cell lines.

Names	No.	Cell Lines	IC ₅₀ (μ M)	References
8-O-Acetyl-sinkiangenorin F	209	AGS	62.7 $\pm$ 2.5	[81]
Coladin	232	HCT116	3.7 $\pm$ 1.5	[75]
		HT-29	5.4 $\pm$ 1.2	[75]
Conferol	267	COLO205	11.19 $\pm$ 0.68	[246]
Conferone	265	COLO205	27.63 $\pm$ 0.69	[246]
		MCF-7	34.02 $\pm$ 0.68	[246]
Episamarcandin	305	AGS	83.8 $\pm$ 1.4	[128]
Farnesiferol A	228	HeLa	20 $\pm$ 0.2	[189]
Farnesiferol C	202	HeLa	25 $\pm$ 0.8	[189]
		AGS	101.6 $\pm$ 1.3	[128]

Table 6. Cont.

Names	No.	Cell Lines	IC ₅₀ (μM)	References
Fekolone	187	AGS	75.4 ± 2.1	[128]
Fekrynol	211	HeLa	35 ± 0.6	[189]
		MGC-803	49 ± 0.8	[189]
		AGS	20 ± 0.5	[189]
Fekrynol acetate	212	HeLa	25 ± 0.6	[189]
		MGC-803	28 ± 1.2	[189]
(+)–Ferulasin	422	PANC-1	2.24 ± 0.83	[195]
		CFPAC-1	6.12 ± 0.52	[195]
		SW1990	11.77 ± 1.57	[195]
		Capan-2	8.57 ± 0.59	[195]
(–)-Ferulasin	423	PANC-1	0.92 ± 0.12	[195]
		CFPAC-1	19.13 ± 2.99	[195]
Feselol	268	COLO205	38.41 ± 0.8	[246]
		MCF-7	35.95 ± 1.29	[246]
Galbanic acid	213	HeLa	43 ± 2.0	[189]
		MCF-7	56.65 ± 1.4	[225]
		MDA-MB-231	48.75 ± 1.16	[225]
4′-Hydroxy kamolonol acetate	277	HeLa	4.5 ± 0.1	[158]
13-Hydroxyfeselol	271	HCT116	34.1 ± 2.3	[75]
		HT-29	35.4 ± 4.0	[75]
Kamolonol	334	HeLa	3.8 ± 0.1	[158]
Kellerin	321	HeLa	37 ± 1.8	[189]
		MCF-7	18.24 ± 0.12	[246]
Lehmannolol	300	HeLa	42 ± 0.9	[189]
		AGS	26.0 ± 0.9	[128]
Lehmannolone	298	HeLa	81.1 ± 1.4	[128]
Mogoltadone	227	COLO205	31.71 ± 0.15	[246]
		MCF-7	30.45 ± 0.6	[246]
		K-562	21.11 ± 0.85	[246]
		HepG2	23.06	[247]
Polyanthinin	226	HeLa	28 ± 0.4	[189]
		MGC-803	45 ± 0.9	[189]
		AGS	45 ± 0.9	[189]
Sinkiangenol E	222	HeLa	16 ± 0.8	[189]
Sinkiangenone A	448	MGC-803	45.05 ± 3.09	[21]
		AGS	48.13 ± 0.87	[21]
Sinkiangenone B	449	MGC-803	18.89 ± 1.32	[21]
		AGS	16.15 ± 0.14	[21]
Sinkiangenorin D	219	HeLa	20.4 ± 1.3	[128]
		AGS	104.8 ± 1.2	[128]
		K562	81.1 ± 1.0	[128]
Sinkiangenorin E	344	AGS	12.7 ± 2.5	[30]
Sinkiangenorin F	208	AGS	27.1 ± 1.4	[81]
Sinkianone	201	HeLa	77.9 ± 0.7	[128]
Umbelliprenin	150	AGS	11.74 ± 1.33	[193]
		BGC-823	24.62 ± 2.45	[193]
		M4Beu	12.4	[232]
		BxPC3	45.15 ± 2.57	[230]
		PANC-1	47.13 ± 5.13	[230]

The anticancer mechanisms of umbelliprenin (150) and galbanic acid (213) are shown in Figure 10.




Figure 10. The anticancer mechanisms of umbelliprenin (150) and galbanic acid (213). MMPs (matrix metalloproteinases), HUVEC (human umbilical vein endothelial cell), HIF (hypoxia-inducible factor), VEGF (vascular endothelial growth factor), EGFR (epithelial growth factor receptor), AKT (protein kinase B). “Red arrow” represent upregulation or downregulation.

4.8. Anti-Acetylcholinesterase Effects

Dastan et al. [163] evaluated the acetylcholinesterase (AChE) inhibitory activity of kamonolol acetate (278) from *F. pseudalliacea*. The results revealed that AChE was suppressed by kamonolol acetate (278), with an IC_{50} value of 63.9 μ M. Moreover, they proved that kamonolol acetate (278) inhibited AChE in the mixed-type model through kinetics together with molecular modeling studies. The findings suggested that kamonolol acetate (278) might be a potential lead compound for designing AChE inhibitors.

4.9. Antidiabetic Effects

Amin et al. [248] conducted antiglycosylation tests on components isolated from the *F. narthex* exudate. The antiglycation activity of the isolated constituents is composed of both oxidative and non-oxidative inhibition modes. In the bovine serum albumin (BSA)-glucose test, ligupersin A (273) displayed a higher activity (IC_{50} : 0.41 mM) than the control, aminoguanidine (IC_{50} : 1.75 mM). In the BSA-methyl glyoxal (MGO) experiment, 5'-acetoxy-8'-hydroxyumbelliprenin (155) showed better activity (IC_{50} : 1.03 mM) than the control, aminoguanidine (IC_{50} : 0.15 mM). In another study, 10'-acetoxy-11'-hydroxyumbelliprenin (167) displayed α -glucosidase inhibitory activity, with an IC_{50} value of 0.05 mM. The results indicated that the antidiabetic activity of the *F. narthex* exudate may be related to the presence of these constituents [249].

5. Conclusions

Several *Ferula* species have a long history of use in traditional medicine due to their potential therapeutic properties in treating various health conditions, such as gastrointestinal disorders, respiratory issues, and inflammatory diseases. In recent years, due to its important edible and medicinal values, extensive research has been conducted on every aspect of *Ferula*, such as its geographical distribution, physiological ecology, genomics, metabolomics, taxonomy, phytoconstituents, biosynthesis, pharmacological activity, traditional uses, clinical efficacy, and industrial applications [250–254].

Ferula is known for its production of sesquiterpenes. Sesquiterpenes are a subclass of terpenes, which are natural hydrocarbons synthesized by plants, including the *Ferula* species, through the mevalonic acid pathway. Sesquiterpenes are composed of three isoprene units, giving them a 15-carbon structure. Sesquiterpenes and their derivatives have antibacterial, antifungal, and antiviral activities, which are characteristically related to plant defense mechanisms [255]. In this work, information on 454 sesquiterpenes and their derivatives from various parts of this plant, including resins, stems, aerial parts, seeds, and roots have been summarized. The specific sesquiterpenes found in *Ferula* species can vary between different plant varieties, and even within the same species, and they are influenced by factors such as environmental conditions and geographic location. These compounds not only give *Ferula* plants their unique aromas but also contribute to their potential therapeutic properties, making them of interest to researchers and practitioners in the fields of herbal medicine. Sesquiterpenes are known for their diverse biological activities, including antioxidative, antibacterial, and anti-inflammatory properties. Ferutin (18), umbelliprenin (150), and galbanic acid (213) are sesquiterpenes from *Ferula* which have undergone extensive pharmacological activity research, and investigating these activities can help uncover potential treatments for a wide range of health conditions. They have also shown promise in drug discovery and development.

It should be noted that the specific biological activity of sesquiterpenes is related to their chemical structure. Researchers should understand their structure–activity relationships to design compounds with better activity, fully tap into their therapeutic potential, and develop standardized applications in medicine industries.

In summary, *Ferula* plants offer a wealth of research opportunities in fields such as phytochemistry, pharmacology, agriculture, ecology, and biotechnology. The diverse sesquiterpenes produced by *Ferula* species have the potential to yield novel drugs, making them a valuable subject of study for researchers across the globe.

Author Contributions: Conceptualization, C.F. and J.S.; software, J.W., Q.Z. and H.W.; writing—original draft preparation, J.W. and Q.Z.; writing—review and editing, J.W., H.W., L.S., G.W., Y.Z., C.F. and J.S.; visualization, Q.Z. and H.W.; supervision, J.S. and J.W.; project administration, J.S. and J.W.; funding acquisition, J.W., Y.Z. and J.S. All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by the CAMS Innovation Fund for Medical Sciences (CIFMS, Grant No. 2022-I2M-1-017), the National Natural Science Foundation of China (Grant No. 81903920), and the Natural Science Foundation of Xinjiang Uygur Autonomous Region, China (Grant No. 2022D01B190).

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: Not applicable.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Akalin, E.; Tuncay, H.O.; Olcay, B.; Miski, M. A New *Ferula* (Apiaceae) Species from Southwest Anatolia: *Ferula Pisidica* Akalin & Miski. *Plants* **2020**, *9*, 740. [\[CrossRef\]](#) [\[PubMed\]](#)
2. Chen, Z.; Zhou, G.; Ma, S. Research Progress of *Ferula Ferulaeoides*: A Review. *Molecules* **2023**, *28*, 3579. [\[CrossRef\]](#) [\[PubMed\]](#)
3. Yaqoob, U.; Nawchoo, I.A. Distribution and Taxonomy of *Ferula* L.: A Review. *Res. Rev. J. Bot. Sci.* **2016**, *5*, 15–23.
4. Mohammadhosseini, M.; Venditti, A.; Sarker, S.D.; Nahar, L.; Akbarzadeh, A. The Genus *Ferula*: Ethnobotany, Phytochemistry and Bioactivities—A Review. *Ind. Crop. Prod.* **2019**, *129*, 350–394. [\[CrossRef\]](#)
5. Ahmed, A.A.; Hegazy, M.E.; Zellagui, A.; Rhouati, S.; Mohamed, T.A.; Sayed, A.A.; Abdella, M.A.; Ohta, S.; Hirata, T. Ferulsinaic Acid, a Sesquiterpene Coumarin with a Rare Carbon Skeleton from *Ferula* Species. *Phytochemistry* **2007**, *68*, 680–686. [\[CrossRef\]](#) [\[PubMed\]](#)
6. Salehi, M.; Naghavi, M.R.; Bahmankar, M. A Review of *Ferula* Species: Biochemical Characteristics, Pharmaceutical and Industrial Applications, and Suggestions for Biotechnologists. *Ind. Crop. Prod.* **2019**, *139*, 111511. [\[CrossRef\]](#)
7. Mahboubi, M. *Ferula Gummosa*, a Traditional Medicine with Novel Applications. *J. Diet. Suppl.* **2016**, *13*, 700–718. [\[CrossRef\]](#) [\[PubMed\]](#)

8. Iranshahy, M.; Iranshahi, M. Traditional Uses, Phytochemistry and Pharmacology of Asafoetida (*Ferula assa-foetida* Oleo-Gum-Resin)—A Review. *J. Ethnopharmacol.* **2011**, *134*, 1–10. [\[CrossRef\]](#)
9. Amalraj, A.; Gopi, S. Biological Activities and Medicinal Properties of Asafoetida: A Review. *J. Tradit. Complement. Med.* **2017**, *7*, 347–359. [\[CrossRef\]](#)
10. Lee, C.L.; Chiang, L.C.; Cheng, L.H.; Liaw, C.C.; Abd El-Razek, M.H.; Chang, F.R.; Wu, Y.C. Influenza A (H1N1) Antiviral and Cytotoxic Agents from *Ferula assa-foetida*. *J. Nat. Prod.* **2009**, *72*, 1568–1572. [\[CrossRef\]](#)
11. Mahendra, P.; Bisht, S. *Ferula asafoetida*: Traditional Uses and Pharmacological Activity. *Pharmacogn. Rev.* **2012**, *6*, 141–146. [\[CrossRef\]](#) [\[PubMed\]](#)
12. Al-Ja'fari, A.-H.; Vila, R.; Freixa, B.; Costa, J.; Cañigueral, S. Antifungal Compounds from the Rhizome and Roots of *Ferula hermonis*. *Phytother. Res. PTR* **2013**, *27*, 911–915. [\[CrossRef\]](#) [\[PubMed\]](#)
13. Elisabetsky, E.; Shanley, P. Ethnopharmacology in the Brazilian Amazon. *Pharmacol. Ther.* **1994**, *64*, 201–214. [\[CrossRef\]](#) [\[PubMed\]](#)
14. Bashir, S.; Alam, M.; Bashir, A.; Aman, A.; Ali, J. Screening of *Ferula narthex* Boiss Crude Methanolic Extract for Analgesic, Gastrointestinal Motility and Insecticidal Activity. *Middle East J. Sci. Res.* **2013**, *14*, 471–475.
15. Akaberi, M.; Iranshahy, M.; Iranshahi, M. Review of the Traditional Uses, Phytochemistry, Pharmacology and Toxicology of Giant Fennel (*Ferula communis* L. subsp. *communis*). *Iran. J. Basic. Med. Sci.* **2015**, *18*, 1050–1062. [\[PubMed\]](#)
16. Sattar, Z.; Iranshahi, M. Phytochemistry and Pharmacology of *Ferula hermonis* Boiss.—A Review. *Drug Res.* **2017**, *67*, 437–446. [\[CrossRef\]](#) [\[PubMed\]](#)
17. Ensiyeh, C.; Ehsan, K.; Ehsan, O.; Mehrdad, I. Antibacterial, Antioxidant and Melanogenesis Inhibitory Activity of Auraptene, a Coumarin from *Ferula szowitsiana* Root. *Nutr. Cancer* **2022**, *74*, 1829–1836. [\[CrossRef\]](#)
18. Motai, T.; Kitanaka, S. Sesquiterpene Phenylpropanoids from *Ferula Fukanensis* and Their Nitric Oxide Production Inhibitory Effects. *J. Nat. Prod.* **2005**, *68*, 365–368. [\[CrossRef\]](#)
19. Bao, X.-H.; Li, Y.-P.; Li, Q.; Cheng, Y.-X.; Jiao, Y.-B.; Zhang, H.-X.; Yan, Y.-M. Racemic Norneolignans from the Resin of *Ferula sinkiangensis* and Their COX-2 Inhibitory Activity. *Fitoterapia* **2023**, *164*, 105341. [\[CrossRef\]](#)
20. Li, G.; Li, X.; Cao, L.; Shen, L.; Zhu, J.; Zhang, J.; Wang, J.; Zhang, L.; Si, J. Steroidal Esters from *Ferula sinkiangensis*. *Fitoterapia* **2014**, *97*, 247–252. [\[CrossRef\]](#)
21. Wang, J.; Gao, Y.; Wang, H.; Chen, L.; Cao, L.; Xu, J.; Li, X.; Zhao, Y.; Zhu, J.; Si, J. Apoptosis Induction and Cell Cycle Arrest Induced by Sinkiangenone B, a Novel Phenylpropanoid Derivative from the Resin of *Ferula sinkiangensis* K. M. Shen. *RSC Adv.* **2018**, *8*, 4093–4103. [\[CrossRef\]](#)
22. Alkhatib, R.; Hennebelle, T.; Joha, S.; Roumy, V.; Güzel, Y.; Biabiany, M.; Idziorek, T.; Preudhomme, C.; Quesnel, B.; Sahpaz, S.; et al. Humulane and Germacrane Sesquiterpenes from *Ferula lycia*. *J. Nat. Prod.* **2010**, *73*, 780–783. [\[CrossRef\]](#) [\[PubMed\]](#)
23. Liu, T.; Wang, L.; Zhang, L.; Jiang, H.; Zhang, Y.; Mao, L. Insecticidal, Cytotoxic and Anti-Phytopathogenic Fungal Activities of Chemical Constituents from the Aerial Parts of *Ferula sinkiangensis*. *Nat. Prod. Res.* **2018**, *34*, 1430–1436. [\[CrossRef\]](#) [\[PubMed\]](#)
24. Yi, X.; Li, Z.; Zheng, Q.; Sang, R.; Li, H.; Gao, G.; Qin, Q.; Zhu, N. Three New Tetrahydrobenzofuran Derivatives from *Ferula sinkiangensis* K.M.Shen and Their Cytotoxic Activities. *Nat. Prod. Res.* **2022**, *37*, 3369–3373. [\[CrossRef\]](#) [\[PubMed\]](#)
25. Eruçar, F.M.; Senadeera, S.P.D.; Wilson, J.A.; Goncharova, E.; Beutler, J.A.; Miski, M. Novel Cytotoxic Sesquiterpene Coumarin Ethers and Sulfur-Containing Compounds from the Roots of *Ferula turcica*. *Molecules* **2023**, *28*, 5733. [\[CrossRef\]](#) [\[PubMed\]](#)
26. Macrì, R.; Bava, I.; Scarano, F.; Mollace, R.; Musolino, V.; Gliozzi, M.; Greco, M.; Foti, D.; Tucci, L.; Maiuolo, J.; et al. In Vitro Evaluation of Ferutinin Rich-*Ferula communis* L., Ssp. *Glauca*, Root Extract on Doxorubicin-Induced Cardiotoxicity: Antioxidant Properties and Cell Cycle Modulation. *Int. J. Mol. Sci.* **2023**, *24*, 12735. [\[CrossRef\]](#) [\[PubMed\]](#)
27. Mi, Y.; Jiao, K.; Xu, J.-K.; Wei, K.; Liu, J.-Y.; Meng, Q.Q.; Guo, T.-T.; Zhang, X.-N.; Zhou, D.; Qing, D.-G.; et al. Kellerin from *Ferula sinkiangensis* Exerts Neuroprotective Effects after Focal Cerebral Ischemia in Rats by Inhibiting Microglia-Mediated Inflammatory Responses. *J. Ethnopharmacol.* **2021**, *269*, 113718. [\[CrossRef\]](#)
28. Arjmand, Z.; Hamburger, M.; Dastan, D. Isolation and Purification of Terpenoid Compounds from *Ferula haussknechtii* and Evaluation of Their Antibacterial Effects. *Nat. Prod. Res.* **2023**, *37*, 1617–1624. [\[CrossRef\]](#)
29. Wang, J.-L.; Sang, C.-Y.; Wang, J.; Li, P.-L.; Chai, T.; Naghavi, M.R.; Zhao, Y.-M.; Yang, J.-L. Sesquiterpene Coumarins from *Ferula sinkiangensis* and Their Anti-Pancreatic Cancer Effects. *Phytochemistry* **2023**, *214*, 113824. [\[CrossRef\]](#)
30. Li, G.Z.; Wang, J.C.; Li, X.J.; Cao, L.; Gao, L.; Lv, N.; Si, J.Y. An Unusual Sesquiterpene Coumarin from the Seeds of *Ferula sinkiangensis*. *J. Asian Nat. Prod. Res.* **2016**, *18*, 891–896. [\[CrossRef\]](#)
31. Dall'Acqua, S.; Linardi, M.A.; Maggi, F.; Nicoletti, M.; Petitto, V.; Innocenti, G.; Basso, G.; Viola, G. Natural Daucane Sesquiterpenes with Antiproliferative and Proapoptotic Activity against Human Tumor Cells. *Bioorg. Med. Chem.* **2011**, *19*, 5876–5885. [\[CrossRef\]](#) [\[PubMed\]](#)
32. Dall'Acqua, S.; Linardi, M.A.; Bortolozzi, R.; Clauser, M.; Marzocchini, S.; Maggi, F.; Nicoletti, M.; Innocenti, G.; Basso, G.; Viola, G. Natural Daucane Esters Induces Apoptosis in Leukaemic Cells through ROS Production. *Phytochemistry* **2014**, *108*, 147–156. [\[CrossRef\]](#) [\[PubMed\]](#)
33. Iranshahi, M.; Amin, G.-R.; Jalalizadeh, H.; Shafiee, A. New Germacrane Derivative from *Ferula persica*. *Pharm. Biol.* **2003**, *41*, 431–433. [\[CrossRef\]](#)
34. González, A.G.; Díaz, J.G.; López, L.A.; Valencia, E.; De Paz, P.P.; Barrera, J.B. Sesquiterpene Esters and Sesquiterpene Coumarin Ethers from *Ferula linkii*-TF. *Phytochemistry* **1993**, *33*, 863–866. [\[CrossRef\]](#)

35. Suzuki, K.; Okasaka, M.; Kashiwada, Y.; Takaishi, Y.; Honda, G.; Ito, M.; Takeda, Y.; Kodzhimatov, O.K.; Ashurmetov, O.; Sekiya, M.; et al. Sesquiterpene Lactones from the Roots of *Ferula varia* and Their Cytotoxic Activity. *J. Nat. Prod.* **2007**, *70*, 1915–1918. [[CrossRef](#)] [[PubMed](#)]
36. Lamnaouer, D.; Martin, M.-T.; Molho, D.; Bodo, B. Isolation of Daucane Esters from *Ferula communis* var. *Brevifolia*. *Phytochemistry* **1989**, *28*, 2711–2716. [[CrossRef](#)]
37. Ahmed, A.A. New Sesquiterpenes from *Ferula sinaica*. *J. Nat. Prod.* **1990**, *53*, 483–486. [[CrossRef](#)]
38. Miski, M.; Mabry, T.J. Daucane Esters from *Ferula communis* subsp. *communis*. *Phytochemistry* **1985**, *24*, 1735–1741. [[CrossRef](#)]
39. Zhu, G.; Zhang, H. Studies on the Chemical Constituents in the Root of *Ferula tunshanica* Su. *J. China Pharm. Univ.* **1996**, *27*, 585–588. [[CrossRef](#)]
40. Miski, M.; Mabry, T.J. Fercolide, a Type of Sesquiterpene Lactone from *Ferula communis* subsp. *communis* and the Correct Structure of Vaginatol. *Phytochemistry* **1986**, *25*, 1673–1675. [[CrossRef](#)]
41. Iranshahi, M.; Ghiadi, M.; Sahebkar, A.; Rahimi, A.; Bassarello, C.; Piacente, S.; Pizza, C. Badrakemonin, a New Eremophilane-Type Sesquiterpene from the Roots of *Ferula badrakema* Kos.-Pol. *Iran. J. Pharm. Res.* **2009**, *8*, 275–279.
42. Fraga, B.M.; Hernández, M.G.; Diaz, J.G. Carotdiol Esters from *Ferula linkii*. *Phytochemistry* **1989**, *28*, 1649–1652. [[CrossRef](#)]
43. Appendino, G.; Jakupovic, J.; Alloatti, S.; Ballero, M. Daucane Esters from *Ferula arrigonii*. *Phytochemistry* **1997**, *45*, 1639–1643. [[CrossRef](#)]
44. Shikishima, Y.; Takaishi, Y.; Honda, G.; Ito, M.; Takeda, Y.; Tori, M.; Takaoka, S.; Kodzhimatov, O.K.; Ashurmetov, O. Sesquiterpenes from *Ferula penninervis*. *J. Nat. Prod.* **2002**, *65*, 1897–1903. [[CrossRef](#)] [[PubMed](#)]
45. Oughlissi-Dehak, K.; Lawton, P.; Michalet, S.; Bayet, C.; Darbour, N.; Hadj-Mahammed, M.; Badjah-Hadj-Ahmed, Y.A.; Dijoux-Franca, M.-G.; Guilet, D. Sesquiterpenes from Aerial Parts of *Ferula vesceritensis*. *Phytochemistry* **2008**, *69*, 1933–1938. [[CrossRef](#)] [[PubMed](#)]
46. Ahmed, A.A.; Abdel-Razek, M.H.; Nassar, M.I.; Izumi, S.; Ohta, S.; Hirata, T. An Eudesmanolide and a Carotane from *Ferula sinaica*. *Phytochemistry* **2001**, *57*, 513–515. [[CrossRef](#)] [[PubMed](#)]
47. Auzi, A.A.; Gray, A.I.; Salem, M.M.; Badwan, A.A.; Sarker, S.D. Feruhermonins A-C: Three Daucane Esters from the Seeds of *Ferula hermonis* (Apiaceae). *J. Asian Nat. Prod. Res.* **2008**, *10*, 711–717. [[CrossRef](#)] [[PubMed](#)]
48. Chen, B.; Teranishi, R.; Kawazoe, K.; Takaishi, Y.; Honda, G.; Itoh, M.; Takeda, Y.; Kodzhimatov, O.K. Sesquiterpenoids from *Ferula kuhistanica*. *Phytochemistry* **2000**, *54*, 717–722. [[CrossRef](#)]
49. Iranshahi, M.; Hosseini, S.T.; Shahverdi, A.R.; Molazade, K.; Khan, S.S.; Ahmad, V.U. Diversolidos A–G, Guaianolides from the Roots of *Ferula diversivittata*. *Phytochemistry* **2008**, *69*, 2753–2757. [[CrossRef](#)]
50. Alkhatib, R.; Hennebelle, T.; Joha, S.; Idziorek, T.; Preudhomme, C.; Quesnel, B.; Sahpaz, S.; Bailleul, F. Activity of Elaeochytrin A from *Ferula elaeochytris* on Leukemia Cell Lines. *Phytochemistry* **2008**, *69*, 2979–2983. [[CrossRef](#)]
51. Naguib, Y.M.A. Pharmaceutically Active Composition Extracted from *Ferula hermonis* and Process of Its Extraction. U.S. Patent US6623768B1, 23 September 2003.
52. Galal, A.M.; Abourashed, E.A.; Ross, S.A.; ElSohly, M.A.; Al-Said, M.S.; El-Ferally, F.S. Daucane Sesquiterpenes from *Ferula hermonis*. *J. Nat. Prod.* **2001**, *64*, 399–400. [[CrossRef](#)]
53. Valle, M.G.; Appendino, G.; Nano, G.M.; Picci, V. Prenylated Coumarins and Sesquiterpenoids from *Ferula communis*. *Phytochemistry* **1986**, *26*, 253–256. [[CrossRef](#)]
54. Fraga, B.M.; González, A.G.; González, P.; Hernandez, M.G.; Larruga, C.L. Carotane Sesquiterpenes from *Ferula lancerottensis*. *Phytochemistry* **1985**, *24*, 501–504. [[CrossRef](#)]
55. Diaz, J.G.; Fraga, B.M.; González, A.G.; González, P.; Hernández, M.G. Eight Carotane Sesquiterpenes from *Ferula linkii*. *Phytochemistry* **1984**, *23*, 2541–2544. [[CrossRef](#)]
56. Lhuillier, A.; Fabre, N.; Cheble, E.; Oueida, F.; Maurel, S.; Valentin, A.; Fourasté, I.; Moulis, C. Daucane Sesquiterpenes from *Ferula hermonis*. *J. Nat. Prod.* **2005**, *68*, 468–471. [[CrossRef](#)] [[PubMed](#)]
57. Wang, N.; Yuan, C.; Kimie, B.; Masahiko, T.; Misunobu, D. Chemical constituents of roots of *Ferula licentiana* var. *tunshanica* and *F. kingdon-wardii* and their systematical significance. *J. Plant Resour. Environ.* **1997**, *6*, 15–18+49.
58. Ahmed, A.A. Daucanes and Other Constituents from *Ferula sinaica*. *Phytochemistry* **1991**, *30*, 1207–1210. [[CrossRef](#)]
59. Singh, M.M.; Agnihotri, A.; Garg, S.N.; Agarwal, S.K.; Gupta, D.N.; Keshri, G.; Kamboj, V.P. Antifertility and Hormonal Properties of Certain Carotane Sesquiterpenes of *Ferula jaeschkeana*. *Planta Med.* **1988**, *54*, 492–494. [[CrossRef](#)]
60. Garg, S.N.; Misra, L.N.; Agarwal, S.K.; Mahajan, V.P.; Rastogi, S.N. Carotane Derivatives from *Ferula jaeschkeana*. *Phytochemistry* **1987**, *26*, 449–450. [[CrossRef](#)]
61. Hu, Y.; Li, X.-D.; Li, G.-Y.; Li, N.; Zuo, W.-J.; Zeng, Y.-M.; Meng, H.; Li, X.; Wang, J.-H. Two Novel Sesquiterpenoids from the Roots of *Ferula ferulaeoides* (Steud.) Korov. *Helv. Chim. Acta* **2010**, *93*, 1019–1024. [[CrossRef](#)]
62. Miski, M.; Ulubelen, A.; Mabry, T.J. Six Sesquiterpene Alcohol Esters from *Ferula elaeochytris*. *Phytochemistry* **1983**, *22*, 2231–2233. [[CrossRef](#)]
63. Razdan, T.K.; Qadri, B.; Qurishi, M.A.; Khuroo, M.A.; Kachroo, K. Sesquiterpene Esters Sesquiterpene-Coumarin Ethers from *Ferula jaeschkeana*. *Phytochemistry* **1989**, *28*, 3389–3393. [[CrossRef](#)]
64. Tamemoto, K.; Takaishi, Y.; Chen, B.; Kawazoe, K.; Shibata, H.; Higuti, T.; Honda, G.; Ito, M.; Takeda, Y.; Kodzhimatov, O.K.; et al. Sesquiterpenoids from the Fruits of *Ferula kuhistanica* and Antibacterial Activity of the Constituents of *F. kuhistanica*. *Phytochemistry* **2001**, *58*, 763–767. [[CrossRef](#)] [[PubMed](#)]
65. Appendino, G.; Cravotto, G.; Sterner, O.; Ballero, M. Oxygenated Sesquiterpenoids from a Nonpoisonous Sardinian Chemotype of Giant Fennel (*Ferula communis*). *J. Nat. Prod.* **2001**, *64*, 393–395. [[CrossRef](#)] [[PubMed](#)]

66. Appendino, G.; Maxia, L.; Bascope, M.; Houghton, P.J.; Sanchez-Duffhues, G.; Muñoz, E.; Sterner, O. A Meroterpenoid NF-kappaB Inhibitor and Drimane Sesquiterpenoids from *Asafetida*. *J. Nat. Prod.* **2006**, *69*, 1101–1104. [[CrossRef](#)] [[PubMed](#)]
67. Garg, S.N.; Agarwal, S.K. Further New Sesquiterpenes from *Ferula jaeschkeana*. *J. Nat. Prod.* **1988**, *51*, 771–774. [[CrossRef](#)] [[PubMed](#)]
68. Ahmed, A.A. Sesquiterpene Coumarins and Sesquiterpenes from *Ferula sinaica*. *Phytochemistry* **1999**, *50*, 109–112. [[CrossRef](#)]
69. Fraga, B.M.; Hernández, M.G.; Díaz, J.G.; González, A.G.; González, P. Carotane Sesquiterpenes from *Ferula linkii*. *Phytochemistry* **1986**, *25*, 2883–2886. [[CrossRef](#)]
70. Tamemoto, K.; Takaishi, Y.; Kawazoe, K.; Honda, G.; Ito, M.; Kiuchi, F.; Takeda, Y.; Kodzhimatov, O.K.; Ashurmetov, O.; Shimizu, K.; et al. An Unusual Sesquiterpene Derivative from *Ferula kuhistanica*. *J. Nat. Prod.* **2002**, *65*, 1323–1324. [[CrossRef](#)]
71. Diaz, J.G.; Fraga, B.M.; González, A.G.; Hernandez, M.G.; Perales, A. Carotane Sesquiterpenes from *Ferula linkii*. *Phytochemistry* **1986**, *25*, 1161–1165. [[CrossRef](#)]
72. Golovina, L.A.; Saidkhodzhaev, A.I. Structure and Stereochemistry of Lapidin. *Chem. Nat. Compd.* **1981**, *17*, 244–248. [[CrossRef](#)]
73. Gamal-Eldeen, A.M.; Hegazy, M.-E.F. A Crystal Lapiferin Derived from *Ferula Vesceritensis* Induces Apoptosis Pathway in MCF-7 Breast Cancer Cells. *Nat. Prod. Res.* **2010**, *24*, 246–257. [[CrossRef](#)] [[PubMed](#)]
74. Motai, T.; Kitanaka, S. Sesquiterpenoids from *Ferula Fukanensis* and Their Inhibitory Effects on Nitric Oxide Production. *J. Nat. Med.* **2006**, *60*, 54–57. [[CrossRef](#)]
75. Jabrane, A.; Jannet, H.B.; Mighri, Z.; Mirjolet, J.F.; Duchamp, O.; Harzallah-Skhiri, F.; Lacaille Dubois, M.A. Two New Sesquiterpene Derivatives from the Tunisian Endemic *Ferula tunetana* Pom. *Chem. Biodivers.* **2010**, *7*, 392–399. [[CrossRef](#)] [[PubMed](#)]
76. Lamnaouer, D.; Fraigui, O.; Martin, M.; Bodo, B. Structure of Ferulenol Derivatives from *Ferula communis* Var. *Genuina*. *Phytochemistry* **1991**, *30*, 2383–2386. [[CrossRef](#)]
77. Appendino, G.; Tagliapietra, S.; Mario Nano, G.; Jakupovic, J. Sesquiterpene Coumarin Ethers from *Asafetida*. *Phytochemistry* **1993**, *35*, 183–186. [[CrossRef](#)]
78. Appendino, G.; Tagliapietra, S.; Gariboldi, P.; Mario Nano, G.; Picci, V. ω -Oxygenated Prenylated Coumarins from *Ferula communis*. *Phytochemistry* **1988**, *27*, 3619–3624. [[CrossRef](#)]
79. Appendino, G.; Tagliapietra, S.; Nano, G.; Picci, V. Ferprenin, a Prenylated Coumarin from *Ferula communis*. *Phytochemistry* **1988**, *27*, 944–946. [[CrossRef](#)]
80. Monti, M.; Pinotti, M.; Appendino, G.; Dallochio, F.; Bellini, T.; Antognoni, F.; Poli, F.; Bernardi, F. Characterization of Anti-Coagulant Properties of Prenylated Coumarin Ferulenol. *Biochim. Biophys. Acta-Gen. Subj.* **2007**, *1770*, 1437–1440. [[CrossRef](#)]
81. Li, G.; Wang, J.; Li, X.; Cao, L.; Lv, N.; Chen, G.; Zhu, J.; Si, J. Two New Sesquiterpene Coumarins from the Seeds of *Ferula sinkiangensis*. *Phytochem. Lett.* **2015**, *13*, 123–126. [[CrossRef](#)]
82. Kajimoto, T.; Yahiro, K.; Nohara, T. Sesquiterpenoid and Disulphide Derivatives from *Ferula assa-foetida*. *Phytochemistry* **1989**, *28*, 1761–1763. [[CrossRef](#)]
83. Abd El-Razek, M.H.; Wu, Y.-C.; Chang, F.-R. Sesquiterpene Coumarins from *Ferula foetida*. *J. Chin. Chem. Soc.* **2007**, *54*, 235–238. [[CrossRef](#)]
84. Ghosh, A.; Banerji, A.; Mandal, S.; Banerji, J. A New Sesquiterpenoid Coumarin from *Ferula assafoetida*. *Nat. Prod. Commun.* **2009**, *4*, 1023–1024. [[CrossRef](#)] [[PubMed](#)]
85. Bandyopadhyay, D.; Banerjee, M.; Laskar, S.; Basak, B. Asimafoetidnol: A New Sesquiterpenoid Coumarin from the Gum Resin of *Ferula assa-foetida*. *Nat. Prod. Commun.* **2011**, *6*, 209–212. [[CrossRef](#)] [[PubMed](#)]
86. Xu, Q.; Li, G.; Liu, Y.; Chen, S.; Xu, L.; Yang, S. Study on Sesquiterpenoid Coumarins from the Root of *Ferula fukanensis* K. M. Shen. *Chin. Pharm.* **2009**, *44*, 255–257. [[CrossRef](#)]
87. Li, X.; Li, G.; Wang, H.; Wang, Y.; Wang, J. Chemical Constituents from *Ferula lehmannii* Boss. *Mod. Chin. Med.* **2010**, *12*, 17–20. [[CrossRef](#)]
88. Banerji, A.; Mallick, B.; Chatterjee, A.; Budzikiewics, H.; Breuer, M. Assafoetidol and Ferocolicin, Two Sesquiterpenoid Coumarins from *Ferula assafoetida* Regel. *Tetrahedron Lett.* **1988**, *29*, 1557–1560. [[CrossRef](#)]
89. Abd El-Razek, M.H.; Ohta, S.; Ahmed, A.A.; Hirata, T. Sesquiterpene Coumarins from the Roots of *Ferula assa-foetida*. *Phytochemistry* **2001**, *58*, 1289–1295. [[CrossRef](#)]
90. Yang, J.; Li, G.; Li, Z.; Qin, H. Chemical Constituents from *Ferula teterrima*. *Nat. Prod. Res. Dev.* **2006**, *18*, 246–248. [[CrossRef](#)]
91. Kiryalov, N.P. Isolation of the Coumarin Badrakemin from the Roots of *Ferula badrakema*. *Chem. Nat. Compd.* **1967**, *3*, 307–309. [[CrossRef](#)]
92. Iranshahi, M.; Amin, G.; Shafiee, A. A New Coumarin from *Ferula persica*. *Pharm. Biol.* **2004**, *42*, 440–442. [[CrossRef](#)]
93. Bagirov, V.Y.; Kir'yalov, N.P. Coumarins from the Roots of *Ferula nevsii*. *Chem. Nat. Compd.* **1970**, *6*, 474. [[CrossRef](#)]
94. Kuliev, Z.A.; Khasanov, T.K. Structures of Cauferin and Cauferidin. *Chem. Nat. Compd.* **1978**, *14*, 271–274. [[CrossRef](#)]
95. Kuliev, Z.A.; Khasanov, T.K.; Malikov, V.M. Structure and Configuration of Cauferinin. *Chem. Nat. Compd.* **1979**, *15*, 127–129. [[CrossRef](#)]
96. Kamoldinov, K.; Li, J.; Eshbakova, K.; Sagdullaev, S.; Xu, G.; Zhou, Y.; Li, J.; Aisa, H.A. Sesquiterpene Coumarins from *Ferula samarkandica* Korovin and Their Bioactivity. *Phytochemistry* **2021**, *187*, 112705. [[CrossRef](#)] [[PubMed](#)]
97. Iranshahi, M.; Masullo, M.; Asili, A.; Hamedzadeh, A.; Jahanbin, B.; Festa, M.; Capasso, A.; Piacente, S. Sesquiterpene Coumarins from *Ferula gumosa*. *J. Nat. Prod.* **2010**, *73*, 1958–1962. [[CrossRef](#)] [[PubMed](#)]
98. Kuliev, Z.A.; Khasanov, T.K.; Malikov, V.M. Terpenoid Coumarin Glycosides of *Ferula conocaule*. *Chem. Nat. Compd.* **1979**, *15*, 414–416. [[CrossRef](#)]
99. Abd El-Razek, M.H.; Ohta, S.; Hirata, T. Terpenoid Coumarins of the Genus *Ferula*. *Heterocycles* **2003**, *60*, 689. [[CrossRef](#)]
100. Dall'Acqua, S.; Maggi, F.; Minesso, P.; Salvagno, M.; Papa, F.; Vittori, S.; Innocenti, G. Identification of Non-Alkaloid Acetylcholinesterase Inhibitors from *Ferulago campestris* (Besser) Grecescu (Apiaceae). *Fitoterapia* **2010**, *81*, 1208–1212. [[CrossRef](#)]

101. Wang, Y.; Si, J.; Li, X.; Jiang, L.; Zhu, J. Chemical Constituents from the Seeds of *Ferula sinkiangensis*. *Mod. Chin. Med.* **2011**, *13*, 26–28. [\[CrossRef\]](#)
102. Pinar, M.; Rodríguez, B. A New Coumarin from *Ferula Loscosii* and the Correct Structure of Coladonin. *Phytochemistry* **1977**, *16*, 1987–1989. [\[CrossRef\]](#)
103. Yoshino, Y.; Tanaka, N.; Tsuji, D.; Itoh, K.; Kashiwada, Y. Communiferulins, Farnesylated Coumarins from the Roots of *Ferula communis* and Their Anti-Neuroinflammatory Activity. *J. Nat. Med.* **2023**, *77*, 173–179. [\[CrossRef\]](#) [\[PubMed\]](#)
104. Iranshahi, M.; Kalategi, F.; Sahebkar, A.; Sardashti, A.; Schneider, B. New Sesquiterpene Coumarins from the Roots of *Ferula flabelliloba*. *Pharm. Biol.* **2010**, *48*, 217–220. [\[CrossRef\]](#) [\[PubMed\]](#)
105. Vandyshev, V.V.; Sklyar, Y.E.; Perel'son, M.E.; Moroz, M.D. Conferdione—A New Coumarin from *Ferula conocaula*. *Chem. Nat. Compd.* **1974**, *10*, 670–671. [\[CrossRef\]](#)
106. Vandyshev, V.V.; Perel'son, M.E.; Sklyar, Y.E.; Moroz, M.D. Conferin—New Coumarin from the Roots of *Ferula conocaula*. *Chem. Nat. Compd.* **1974**, *10*, 673–674. [\[CrossRef\]](#)
107. Vandyshev, V.V.; Sklyar, Y.E.; Perel'son, M.E.; Moroz, M.D.; Pimenov, M.G. Conferol—A New Coumarin from the Roots of *Ferula conocaula* and *F. moschata*. *Chem. Nat. Compd.* **1972**, *8*, 654. [\[CrossRef\]](#)
108. Iranshahi, M.; Rezaee, R.; Sahebkar, A.; Bassarello, C.; Piacente, S.; Pizza, C. Sesquiterpene Coumarins from the Fruits of *Ferula badrakema*. *Pharm. Biol.* **2009**, *47*, 344–347. [\[CrossRef\]](#)
109. Andrianova, V.B.; Sklyar, Y.E.; Perel'son, M.E.; Pimenov, M.G. Kellerin—A New Coumarin from the Roots of *Ferula kelleri*. *Chem. Nat. Compd.* **1973**, *9*, 758–759. [\[CrossRef\]](#)
110. Nabiev, A.A.; Khasanov, T.K.; Malikov, V.M. Terpenoid Coumarins of *Ferula kokanica*. *Chem. Nat. Compd.* **1982**, *18*, 547–549. [\[CrossRef\]](#)
111. Beselovskaya, N.V.; Sklyar, Y.E. Deacetyltaadzhikorin from *Ferula tadshikorum*. *Chem. Nat. Compd.* **1984**, *20*, 363. [\[CrossRef\]](#)
112. El-Bassuony, A.A.; Gohar, A.A.; Kabbash, A.M. Two New Sesquiterpene Coumarins, Ferusinol and Samarcandin Diastereomer, from *Ferula sinaica*. *Iran. J. Pharm. Res.* **2007**, *6*, 217–221.
113. Motai, T.; Daikonya, A.; Kitanaka, S. Sesquiterpene Coumarins from *Ferula fukanensis* and Nitric Oxide Production Inhibitory Effects. *J. Nat. Prod.* **2004**, *67*, 432–436. [\[CrossRef\]](#) [\[PubMed\]](#)
114. Isaka, K.; Nagatsu, A.; Ondognii, P.; Zevgeegiin, O.; Gombosurengyin, P.; Davgiin, K.; Kojima, K.; Ogihara, Y. Sesquiterpenoid Derivatives from *Ferula ferulioides*. *V. Chem. Pharm. Bull.* **2001**, *49*, 1072–1076. [\[CrossRef\]](#) [\[PubMed\]](#)
115. Nagatsu, A.; Isaka, K.; Kojima, K.; Ondognii, P.; Zevgeegiin, O.; Gombosurengyin, P.; Davgiin, K.; Irfan, B.; Iqbal, C.M.; Ogihara, Y. New Sesquiterpenes from *Ferula ferulaeoides* (Steud.) Korovin. VI. Isolation and Identification of Three New Dihydrofuro[2,3-b]Chromones. *Chem. Pharm. Bull.* **2002**, *50*, 675–677. [\[CrossRef\]](#) [\[PubMed\]](#)
116. Liu, T.; Wang, S.; Xu, L.; Fu, W.; Gibbons, S.; Mu, Q. Sesquiterpenoids with Anti-MDR Staphylococcus Aureus Activities from *Ferula ferulioides*. *Chem. Biodivers.* **2015**, *12*, 599–614. [\[CrossRef\]](#) [\[PubMed\]](#)
117. Yang, J. Studies on the Chemical Constituents of *Ferula teterrima*, *F. sinkiangensis* and *F. ferulaeoides*. Master's Thesis, Chinese Academy of Medical Sciences & Peking Union Medical College, Beijing, China, 2006.
118. Yang, J.; Jing, S.; Li, Z.; Qin, H. Chemical Constituents from Roots of *Ferula sinkiangensis*. *China J. Chin. Mater. Med.* **2007**, *32*, 2382–2384. [\[CrossRef\]](#)
119. Ahmed, A.A.; Mohamed, A.E.-H.H.; El-Razek, M.H.A.; Hegazy, M.-E.F. Ferulol and Epi-Samarcandin, Two New Sesquiterpene Coumarins from *Ferula sinaica*. *Nat. Prod. Commun.* **2007**, *2*, 521–524. [\[CrossRef\]](#)
120. Nassar, M.I.; Mohamed, T.K. A New Sesquiterpene Coumarin from *Ferula assa-foetida*. *Fitoterapia* **1998**, *69*, 41–42.
121. Dastan, D.; Salehi, P.; Reza Gohari, A.; Zimmermann, S.; Kaiser, M.; Hamburger, M.; Reza Khavasi, H.; Nejad Ebrahimi, S. Disesquiterpene and Sesquiterpene Coumarins from *Ferula pseudalliacea*, and Determination of Their Absolute Configurations. *Phytochemistry* **2012**, *78*, 170–178. [\[CrossRef\]](#)
122. Iranshahi, M.; Arfa, P.; Ramezani, M.; Jaafari, M.R.; Sadeghian, H.; Bassarello, C.; Piacente, S.; Pizza, C. Sesquiterpene Coumarins from *Ferula szowitsiana* and in Vitro Antileishmanial Activity of 7-Prenyloxycoumarins against Promastigotes. *Phytochemistry* **2007**, *68*, 554–561. [\[CrossRef\]](#)
123. Li, X.; Li, G.; Wang, H.; Wang, Y.; Wang, J. Chemical Constituents from *Ferula lehmannii* Boss. *J. Nong. Med.* **2010**, *32*, 198–200. [\[CrossRef\]](#)
124. Cha, M.-R.; Choi, Y.; Choi, C.; Kim, Y.; Kim, Y.-K.; Ryu, S.; Kim, Y.; Choi, S. Galbanic Acid, a Cytotoxic Sesquiterpene from the Gum Resin of *Ferula asafoetida*, Blocks Protein Farnesyltransferase. *Planta Med.* **2011**, *77*, 52–54. [\[CrossRef\]](#) [\[PubMed\]](#)
125. Sistani, P.; Dehghan, G.; Sadeghi, L. Structural and Kinetic Insights into HIV-1 Reverse Transcriptase Inhibition by Farnesiferol, C. *Int. J. Biol. Macromol.* **2021**, *174*, 309–318. [\[CrossRef\]](#) [\[PubMed\]](#)
126. Golovina, L.A.; Khasanov, T.K.; Saidkhodzhaev, A.I.; Malikov, V.M.; Rakhmankulov, U. Coumarins and Esters of *Ferula microcarpa*. *Chem. Nat. Compd.* **1978**, *14*, 487–490. [\[CrossRef\]](#)
127. Nabiev, A.A.; Khasanov, T.K.; Malikov, V.M. New Terpenoid Coumarins of *Ferula kopetdaghensis*. *Chem. Nat. Compd.* **1978**, *14*, 440. [\[CrossRef\]](#)
128. Li, G.; Li, X.; Cao, L.; Zhang, L.; Shen, L.; Zhu, J.; Wang, J.; Si, J. Sesquiterpene Coumarins from Seeds of *Ferula sinkiangensis*. *Fitoterapia* **2015**, *103*, 222–226. [\[CrossRef\]](#) [\[PubMed\]](#)
129. Veselovskaya, N.V.; Sklyar, Y.E.; Fesenko, D.A.; Pimenov, M.G. Fekrol—A Terpenoid Coumarin from *Ferula krylovii*. *Chem. Nat. Compd.* **1979**, *15*, 755. [\[CrossRef\]](#)
130. Veselovskaya, N.V.; Sklyar, Y.E.; Savina, A.A. Fekrynol and Its Acetate from *Ferula krylovii*. *Chem. Nat. Compd.* **1981**, *17*, 589–590. [\[CrossRef\]](#)
131. Miski, M.; Jakupovic, J. Cyclic Farnesyl-Coumarin and Farnesyl-Chromone Derivatives from *Ferula communis* subsp. *communis*. *Phytochemistry* **1990**, *29*, 1995–1998. [\[CrossRef\]](#)

132. Kuliev, Z.A.; Khasanov, T.K. The Structures of Ferocaulin, Ferocaulinin, Ferocaulidin, and Ferocaulicin. *Chem. Nat. Compd.* **1978**, *14*, 267–271. [\[CrossRef\]](#)
133. Saidkhodzhaev, A.I.; Malikov, V.M. The Stereochemistry of Feropolol, Feropolin, Feropolone, and Feropolidin. *Chem. Nat. Compd.* **1978**, *14*, 681–682. [\[CrossRef\]](#)
134. Khasanov, T.K.; Saidkhodzhaev, A.I.; Nikonov, G.K. The Structures of Feropolin, Feropolol, Feropolone, and Feropolidin. *Chem. Nat. Compd.* **1976**, *12*, 79–80. [\[CrossRef\]](#)
135. Saidkhodzhaev, A.I.; Malikov, V.M.; Pimenov, M.G.; Melibaev, S. Terpenoid Coumarins of *Ferula lipskvi* and *F. vicaria*. *Chem. Nat. Compd.* **1991**, *27*, 242–243. [\[CrossRef\]](#)
136. Kadyrov, A.S.; Saidkhodzhaev, A.I.; Nikonov, G.K. The Structures of Feroside and of Reoselin a—New Glycosides from the Roots of *Ferula korshinskyi*. *Chem. Nat. Compd.* **1975**, *11*, 604–608. [\[CrossRef\]](#)
137. Guo, T.; Dang, W.; Zhou, Y.; Zhou, D.; Meng, Q.; Xu, L.; Chen, G.; Lin, B.; Qing, D.; Sun, Y.; et al. Sesquiterpene Coumarins Isolated from *Ferula bungeana* and Their Anti-Neuroinflammatory Activities. *Bioorg. Chem.* **2022**, *128*, 106102. [\[CrossRef\]](#) [\[PubMed\]](#)
138. Kir'yanova, I.A.; Sklyar, Y.E. Ferucrin Isobutyrate and Ferucrinone from *Ferula foetidissima*. *Chem. Nat. Compd.* **1984**, *20*, 617–618. [\[CrossRef\]](#)
139. Nabiev, A.A.; Khasanov, T.K.; Malikov, V.M. A Chemical Study of the Roots of *Ferula kopetdagensis*. *Chem. Nat. Compd.* **1979**, *15*, 14–16. [\[CrossRef\]](#)
140. Perel'son, M.E.; Sklyar, Y.E.; Veselovskaya, N.V.; Pimenov, M.G. Ferukrin, a New Terpinoid Coumarin from *Ferula krylovii*. *Pharm. Chem. J.* **1977**, *11*, 366–369. [\[CrossRef\]](#)
141. Lamnaouer, D.; Bodo, B.; Martin, M.-T.; Molho, D. Ferulenol and ω -Hydroxyferulenol, Toxic Coumarins from *Ferula communis* Var. *Genuina*. *Phytochemistry* **1987**, *26*, 1613–1615. [\[CrossRef\]](#)
142. Carboni, S.; Malaguzzi, V.; Marsili, A. Ferulenol a New Coumarin Derivative from *Ferula communis*. *Tetrahedron Lett.* **1964**, *5*, 2783–2786. [\[CrossRef\]](#)
143. Meng, H.; Li, G.; Huang, J.; Zhang, K.; Wang, H.; Wang, J. Sesquiterpene Coumarin and Sesquiterpene Chromone Derivatives from *Ferula ferulaeoides* (Steud.) Korov. *Fitoterapia* **2013**, *86*, 70–77. [\[CrossRef\]](#)
144. Obul, M.; Wang, X.; Zhao, J.; Li, G.; Aisa, H.A.; Huang, G. Structural Modification on Rupestonic Acid Leads to Highly Potent Inhibitors against Influenza Virus. *Mol. Divers.* **2019**, *23*, 1–9. [\[CrossRef\]](#) [\[PubMed\]](#)
145. El-Bassuony, A.A. Antibacterial Activity of a Novel Sesquiterpene Coumarin from *Ferula sinaica*. *Asian J. Chem.* **2007**, *19*, 2053–2058.
146. Guo, T.; Zhou, D.; Yang, Y.; Zhang, X.; Chen, G.; Lin, B.; Sun, Y.; Ni, H.; Liu, J.; Hou, Y.; et al. Bioactive Sesquiterpene Coumarins from the Resin of *Ferula sinkiangensis* Targeted on over-Activation of Microglia. *Bioorg. Chem.* **2020**, *104*, 104338. [\[CrossRef\]](#)
147. Nabiev, A.A.; Khasanov, T.K.; Melibaev, S. Coumarins of *Ferula diversivittata*. *Chem. Nat. Compd.* **1978**, *14*, 441. [\[CrossRef\]](#)
148. Nabiev, A.A.; Malikov, V.M.; Khasanov, T.K. Karatavicin—A New Coumarin from *Ferula karatavica*. *Chem. Nat. Compd.* **1983**, *19*, 498. [\[CrossRef\]](#)
149. Wang, J.; Huo, X.; Wang, H.; Dong, A.; Zheng, Q.; Si, J. Undescribed Sesquiterpene Coumarins from the Aerial Parts of *Ferula sinkiangensis* and Their Anti-Inflammatory Activities in Lipopolysaccharide-Stimulated RAW 264.7 Macrophages. *Phytochemistry* **2023**, *210*, 113664. [\[CrossRef\]](#) [\[PubMed\]](#)
150. Perel'son, M.E.; Sokolova, A.I.; Sklyar, Y.E. The Structure of Feterin—A New Terpenoid Coumarin from *Ferula teterrima*. *Chem. Nat. Compd.* **1978**, *14*, 263–267. [\[CrossRef\]](#)
151. Bashir, S.; Alam, M.; Adhikari, A.; Shrestha, R.L.; Yousuf, S.; Ahmad, B.; Parveen, S.; Aman, A.; Iqbal Choudhary, M. New Antileishmanial Sesquiterpene Coumarins from *Ferula narthex* Boiss. *Phytochem. Lett.* **2014**, *9*, 46–50. [\[CrossRef\]](#)
152. Buddrus, J.; Bauer, H.; Abumustafa, E.; Khattab, A.; Mishaal, S.; Elkhriy, E.; Linscheid, M. Foetidin, a Sesquiterpenoid Coumarin from *Ferula-assa-foetida*. *Phytochemistry* **1985**, *24*, 869–870. [\[CrossRef\]](#)
153. Wu, Y.-C.; Abd El-Razek, M.H.; Chang, F.-R.; Liaw, C.-C.; Nassar, M.I.; Huang, H.-C.; Chen, Y.-H.; Yang, Y.-L. Two Sesquiterpene-Coumarins from the Roots of *Ferula marmarica*. *Heterocycles* **2004**, *63*, 2101. [\[CrossRef\]](#)
154. Kadyrov, A.S.; Saidkhodzhaev, A.I.; Malikov, V.M. The Structure of Foliferin. *Chem. Nat. Compd.* **1978**, *14*, 442–443. [\[CrossRef\]](#)
155. Motai, T.; Kitanaka, S. Sesquiterpene Coumarins from *Ferula Fukanensis* and Nitric Oxide Production Inhibitory Effects (2). *Chem. Pharm. Bull.* **2004**, *52*, 1215–1218. [\[CrossRef\]](#) [\[PubMed\]](#)
156. Motai, T.; Daikonya, A.; Kitanaka, S. Sesquiterpene Coumarins from *Ferula Fukanensis* and Their Pro-Inflammatory Cytokine Gene Expression Inhibitory Effects. *Chem. Pharm. Bull.* **2013**, *61*, 618–623. [\[CrossRef\]](#) [\[PubMed\]](#)
157. Bagirov, V.Y.; Sheichenko, V.I.; Veselovskaya, N.V.; Sklyar, Y.E.; Savina, A.A.; Kir'yanova, I.A. Structure and Stereochemistry of Galbanic Acid. *Chem. Nat. Compd.* **1980**, *16*, 439–441. [\[CrossRef\]](#)
158. Dastan, D.; Salehi, P.; Reza Gohari, A.; Ebrahimi, S.; Aliahmadi, A.; Hamburger, M. Bioactive Sesquiterpene Coumarins from *Ferula pseudalliacea*. *Planta Med.* **2014**, *80*, 1118–1123. [\[CrossRef\]](#) [\[PubMed\]](#)
159. Lamnaouer, D.; Fraigui, O.; Martin, M.-T.; Gallard, J.-F.; Bodo, B. Structure of Isoferprenin, a 4-Hydroxycoumarin Derivative from *Ferula communis* Var. *Genuina*. *J. Nat. Prod.* **1991**, *54*, 576–578. [\[CrossRef\]](#)
160. Yang, J.-R.; An, Z.; Li, Z.-H.; Jing, S.; Qin, H.-L. Sesquiterpene Coumarins from the Roots of *Ferula sinkiangensis* and *Ferula teterrima*. *Chem. Pharm. Bull.* **2006**, *54*, 1595–1598. [\[CrossRef\]](#)
161. Nabiev, A.A.; Malikov, V.M. Microlobin—A New Coumarin from *Ferula microloba*. *Chem. Nat. Compd.* **1983**, *19*, 664–667. [\[CrossRef\]](#)
162. Paknikar, S.; Kirtany, J. Terpenoid Parts of Kamolone and Kamolol of *Ferula penninervis*. *Experientia* **1974**, *30*, 224–225. [\[CrossRef\]](#)

163. Dastan, D.; Validi, S.; Ebadi, A. Kamonolol Acetate from *Ferula Pseudalliacea* as AChE Inhibitor: In Vitro and in Silico Studies. *Struct. Chem.* **2020**, *31*, 965–973. [\[CrossRef\]](#)
164. Bagirov, V.Y.; Sheichenko, V.I. The Structure of Karatavic Acid. *Chem. Nat. Compd.* **1975**, *11*, 734–736. [\[CrossRef\]](#)
165. Kir'yalov, N.P.; Bagirov, V.Y. The Structure of Karatavic Acid. *Chem. Nat. Compd.* **1968**, *4*, 240–242. [\[CrossRef\]](#)
166. Veselovskaya, N.V.; Sklyar, Y.E.; Pimenov, M.G. Terpenoid Coumarins of *Ferula aitchisonii*. *Chem. Nat. Compd.* **1982**, *18*, 368. [\[CrossRef\]](#)
167. Kir'yalov, N.P.; Bagirov, V.Y. The Structure of Karatavicinol. *Chem. Nat. Compd.* **1969**, *5*, 191–192. [\[CrossRef\]](#)
168. Saidkhodzhaev, A.I.; Mukumova, D.U.; Kamilov, K.M.; Malikov, V.M.; Pimenov, M.G. Terpenoid Coumarins of *Ferula cummosa*. *Chem. Nat. Compd.* **1991**, *27*, 243–244. [\[CrossRef\]](#)
169. Nabiev, A.A.; Khasanov, T.K.; Malikov, V.M. A New Terpenoid Coumarin from *Ferula kopetdaghensis*. *Chem. Nat. Compd.* **1982**, *18*, 44–46. [\[CrossRef\]](#)
170. Kamilov, K.M.; Nikonov, G.K. Coumarins of *Ferula Kopetdaghensis* and the Structure of Kopeolin and Kopeoside. *Chem. Nat. Compd.* **1973**, *9*, 294–298. [\[CrossRef\]](#)
171. Kamilov, K.M.; Nikonov, G.K. Coumarins of *Ferula Kopetdaghensis*—Kopetdaghin and Kopeodin (Farnesiferol B). *Chem. Nat. Compd.* **1974**, *10*, 450–453. [\[CrossRef\]](#)
172. Gu, X.-Y.; Zhang, H.-Q.; Xue, Y.-C.; Lv, Y.; Zheng, Q.-T. A study of chemical structure of lehmannolone. *J. Plant Resour. Environ.* **1995**, *4*, 13–16.
173. Sagitdinova, G.V.; Saidkhodzhaev, A.I.; Malikov, V.M. Structure and Stereochemistry of the Coumarins of *Ferula lehmannii*. *Chem. Nat. Compd.* **1983**, *19*, 672–675. [\[CrossRef\]](#)
174. Nabiev, A.A.; Malikov, V.M. Microlobidene—A Terpenoid Coumarin from *Ferula Microloba* with a New Type of Terpenoid Skeleton. *Chem. Nat. Compd.* **1983**, *19*, 743–744. [\[CrossRef\]](#)
175. Khasanov, T.K.; Saidkhodzhaev, A.I.; Nikonov, G.K. Structure and Configuration of the Coumarins Mogoltadone and Mogoltadin. *Chem. Nat. Compd.* **1974**, *10*, 20–23. [\[CrossRef\]](#)
176. Khasanov, T.K.; Saidkhodzhaev, A.I.; Nikonov, G.K. Structure and Configuration of New Coumarins of the Roots of *Ferula mogoltavica*. *Chem. Nat. Compd.* **1974**, *10*, 8–11. [\[CrossRef\]](#)
177. Zakharov, P.I.; Terent'ev, P.B.; Nikonov, G.K.; Ban'kovskii, A.I. Mass-Spectrometric Study of the Coumarins Mogoltin, Mogoltavin, and Mogoltavinin. *Chem. Nat. Compd.* **1974**, *10*, 15–19. [\[CrossRef\]](#)
178. Kir'yanova, I.A.; Sklyar, Y.E.; Pimenov, M.G.; Baranova, Y.V. Terpenoid Coumarins of *Ferula foetidissima* and *F. inciso-serrata*. *Chem. Nat. Compd.* **1982**, *18*, 489. [\[CrossRef\]](#)
179. Su, B.N.; Takaishi, Y.; Honda, G.; Itoh, M.; Takeda, Y.; Kodzhimatov, O.K.; Ashurmetov, O. Sesquiterpene Coumarins and Related Derivatives from *Ferula pallida*. *J. Nat. Prod.* **2000**, *63*, 436–440. [\[CrossRef\]](#)
180. Iranshahi, M.; Mojarab, M.; Sadeghian, H.; Hanafi-Bojd, M.Y.; Schneider, B. Polar Secondary Metabolites of *Ferula persica* Roots. *Phytochemistry* **2008**, *69*, 473–478. [\[CrossRef\]](#)
181. Perel'son, M.E. The Configuration of Polyanthin and Polyanthinin. *Chem. Nat. Compd.* **1975**, *11*, 254. [\[CrossRef\]](#)
182. Bukreeva, T.V. Components of the Resin of the Roots of *Ferula kirialovii*. *Chem. Nat. Compd.* **1987**, *23*, 633–634. [\[CrossRef\]](#)
183. Kir'yalov, N.P.; Bukreeva, T.V.; Gindin, V.A.; Mamatov, G.; Kozhina, I.S. The Structure of the Coumarin Glucoside Reoselin from the Roots of *Ferula pseudooreoselinum*. *Chem. Nat. Compd.* **1975**, *11*, 88–90. [\[CrossRef\]](#)
184. Kamilov, K.M.; Nikonov, G.K. Structure of the Coumarin Reoselin. *Chem. Nat. Compd.* **1974**, *10*, 82–83. [\[CrossRef\]](#)
185. Kir'yalov, N.P.; Movchan, S.D. The Structure of Samarcandin and Samarcandone, Coumarin Compounds from *Ferula samarcandica*. *Chem. Nat. Compd.* **1968**, *4*, 63–65. [\[CrossRef\]](#)
186. Kir'yalov, N.P.; Bukreeva, T.V. Samarcandin Acetate from the Roots of *Ferula pseudooreoselinum*. *Chem. Nat. Compd.* **1972**, *8*, 777–778. [\[CrossRef\]](#)
187. Bandyopadhyay, D.; Basak, B.; Chatterjee, A.; Lai, T.K.; Banerji, A.; Banerji, J.; Neuman, A.; Prangé, T. Saradaferin, a New Sesquiterpenoid Coumarin from *Ferula assafoetida*. *Nat. Prod. Res.* **2006**, *20*, 961–965. [\[CrossRef\]](#) [\[PubMed\]](#)
188. Shomirzoeva, O.; Xu, M.-Y.; Sun, Z.-J.; Li, C.; Nasriddinov, A.; Muhidinov, Z.; Zhang, K.; Gu, Q.; Xu, J. Chemical Constituents of *Ferula seravschanica*. *Fitoterapia* **2021**, *149*, 104829. [\[CrossRef\]](#) [\[PubMed\]](#)
189. Wang, J.; Wang, H.; Zhang, M.; Li, X.; Zhao, Y.; Chen, G.; Si, J.; Jiang, L. Sesquiterpene Coumarins from *Ferula sinkiangensis* K.M.Shen and Their Cytotoxic Activities. *Phytochemistry* **2020**, *180*, 112531. [\[CrossRef\]](#)
190. Xing, Y.; Li, N.; Zhou, D.; Chen, G.; Jiao, K.; Wang, W.; Si, Y.; Hou, Y. Sesquiterpene Coumarins from *Ferula sinkiangensis* Act as Neuroinflammation Inhibitors. *Planta Med.* **2016**, *83*, 135–142. [\[CrossRef\]](#)
191. Khalilova, É.K.; Saidkhodzhaev, A.I. Terpenoid Coumarins of *Ferula sumbul*. *Chem. Nat. Compd.* **1998**, *34*, 506–507. [\[CrossRef\]](#)
192. Perel'son, M.E.; Bandyshev, V.V.; Sklyar, Y.E.; Vezhkhovska-Renke, K.; Veselovskaya, N.V.; Pimenov, M.G. New Terpenoid Coumarins from *Ferula tadshikorum*. *Chem. Nat. Compd.* **1976**, *12*, 533–537. [\[CrossRef\]](#)
193. Zhang, L.; Sun, X.; Si, J.; Li, G.; Cao, L. Umbelliprenin Isolated from *Ferula sinkiangensis* Inhibits Tumor Growth and Migration through the Disturbance of Wnt Signaling Pathway in Gastric Cancer. *PLoS ONE* **2019**, *14*, e0207169. [\[CrossRef\]](#)
194. Al-Hazimi, H.M.G. Terpenoids and a Coumarin from *Ferula sinaica*. *Phytochemistry* **1986**, *25*, 2417–2419. [\[CrossRef\]](#)
195. Wang, J.-L.; Meng, X.-H.; Zheng, Y.-D.; Sang, C.-Y.; Wang, W.-F.; Ma, J.-Y.; Ha, T.-K.-Q.; Zhao, Y.-M.; Yang, J.-L. (±)-Ferulasin, Unusual Sesquiterpene Chromones from *Ferula sinkiangensis*. *Tetrahedron* **2022**, *122*, 132953. [\[CrossRef\]](#)
196. Motai, T.; Kitanaka, S. Sesquiterpene Chromones from *Ferula Fukanensis* and Their Nitric Oxide Production Inhibitory Effects. *J. Nat. Prod.* **2005**, *68*, 1732–1735. [\[CrossRef\]](#) [\[PubMed\]](#)

197. Su, B.-N.; Takaishi, Y.; Honda, G.; Itoh, M.; Takeda, Y.; Kodzhimatov, O.K.; Ashurmetov, O. Sesquiterpene Phenylpropanoid and Sesquiterpene Chromone Derivatives from *Ferula pallida*. *J. Nat. Prod.* **2000**, *63*, 520–522. [\[CrossRef\]](#) [\[PubMed\]](#)
198. Meng, H.; Li, G.; Huang, J.; Zhang, K.; Wei, X.; Ma, Y.; Zhang, C.; Wang, J. Sesquiterpenoid Derivatives from *Ferula ferulaeoides* (Steud.) Korov. *Phytochemistry* **2013**, *86*, 151–158. [\[CrossRef\]](#) [\[PubMed\]](#)
199. Kojima, K.; Isaka, K.; Ondognii, P.; Zevgeegiin, O.; Davgiin, K.; Mizukami, H.; Ogihara, Y. Sequiterpenoid Derivatives from *Ferula ferulioides*. III. *Chem. Pharm. Bull.* **1999**, *47*, 1145–1147. [\[CrossRef\]](#)
200. Liu, T.; Osman, K.; Kaatz, G.W.; Gibbons, S.; Mu, Q. Antibacterial Sesquiterpenoid Derivatives from *Ferula ferulaeoides*. *Planta Med.* **2013**, *79*, 701–706. [\[CrossRef\]](#)
201. Kojima, K.; Isaka, K.; Purev, O.; Jargalsaikhan, G.; Suran, D.; Mizukami, H.; Ogihara, Y. Sesquiterpenoid Derivatives from *Ferula ferulioides*. *Chem. Pharm. Bull.* **1998**, *46*, 1781–1784. [\[CrossRef\]](#)
202. Xu, H.; Dickschat, J.S. Germacrene A—A Central Intermediate in Sesquiterpene Biosynthesis. *Chemistry* **2020**, *26*, 17318–17341. [\[CrossRef\]](#)
203. Xu, H.; Dickschat, J.S. Germacrene B—A Central Intermediate in Sesquiterpene Biosynthesis. *Beilstein J. Org. Chem.* **2023**, *19*, 186–203. [\[CrossRef\]](#) [\[PubMed\]](#)
204. Zhou, D.; Li, N.; Zhang, Y.; Yan, C.; Jiao, K.; Sun, Y.; Ni, H.; Lin, B.; Hou, Y. Biotransformation of Neuro-Inflammation Inhibitor Kelllerin Using *Angelica Sinensis* (Oliv.) Diels Callus. *RSC Adv.* **2016**, *6*, 97302–97312. [\[CrossRef\]](#)
205. Ibraheim, Z.Z.; Abdel-Mageed, W.M.; Dai, H.; Guo, H.; Zhang, L.; Jaspars, M. Antimicrobial Antioxidant Daucane Sesquiterpenes from *Ferula hermonis* Boiss. *Phytother. Res.* **2012**, *26*, 579–586. [\[CrossRef\]](#) [\[PubMed\]](#)
206. Abourashed, E.A.; Galal, A.M.; Shibl, A.M. Antimycobacterial Activity of Ferutinin Alone and in Combination with Antitubercular Drugs against a Rapidly Growing Surrogate of *Mycobacterium Tuberculosis*. *Nat. Prod. Res.* **2011**, *25*, 1142–1149. [\[CrossRef\]](#) [\[PubMed\]](#)
207. Appendino, G.; Mercalli, E.; Fuzzati, N.; Arnoldi, L.; Stavri, M.; Gibbons, S.; Ballero, M.; Maxia, A. Antimycobacterial Coumarins from the Sardinian Giant Fennel (*Ferula communis*). *J. Nat. Prod.* **2004**, *67*, 2108–2110. [\[CrossRef\]](#) [\[PubMed\]](#)
208. Al-Yahya, M.A.; Muhammad, I.; Mirza, H.H.; El-Ferally, F.S. Antibacterial Constituents from the Rhizomes of *Ferula communis*. *Phytother. Res.* **1998**, *12*, 335–339. [\[CrossRef\]](#)
209. Shahverdi, A.R.; Iranshahi, M.; Mirjani, R.; JAMALIFAR, H.; Amin, G. Bioassay-Guided Isolation and Identification of an Antibacterial Compound from *Ferula persica* Var. *Persica* Roots. *DARU J. Pharm. Sci.* **2005**, *13*, 17–19.
210. Pavlović, I.; Petrović, S.; Milenković, M.; Stanojković, T.; Nikolić, D.; Krnić, A.; Niketić, M. Antimicrobial and Cytotoxic Activity of Extracts of *Ferula heuffelii* Griseb. Ex Heuff. and Its Metabolites. *Chem. Biodivers.* **2015**, *12*, 1585–1594. [\[CrossRef\]](#)
211. Sun, Z.-L.; Liu, T.; Wang, S.-Y.; Ji, X.-Y.; Mu, Q. TLC-Bioautography Directed Isolation of Antibacterial Compounds from Active Fractionation of *Ferula ferulioides*. *Nat. Prod. Res.* **2019**, *33*, 1761–1764. [\[CrossRef\]](#)
212. Shahverdi, A.R.; Fakhimi, A.; Zarrini, G.; Dehghan, G.; Iranshahi, M. Galbanic Acid from *Ferula szowitsiana* Enhanced the Antibacterial Activity of Penicillin G and Cephalexin against *Staphylococcus Aureus*. *Biol. Pharm. Bull.* **2007**, *30*, 1805–1807. [\[CrossRef\]](#)
213. Bazzaz, B.S.F.; Du, A.R.; Iranshahi, M.; Naderinasab, M.; Karamodin, M.K. Evaluating the Potentiating Effect of Galbanic Acid from *Ferula szowitsiana* on Three Common Antibiotics against Resistant Hospital Isolates of *Staphylococcus Aureus*. *Iran. J. Pharm. Res.* **2009**, *8*, 217–221. [\[CrossRef\]](#)
214. Safdari, H.; Neshani, A.; Sadeghian, A.; Ebrahimi, M.; Iranshahi, M.; Sadeghian, H. Potent and Selective Inhibitors of Class A β -Lactamase: 7-Prenyloxy Coumarins. *J. Antibiot.* **2014**, *67*, 373–377. [\[CrossRef\]](#) [\[PubMed\]](#)
215. Dastan, D.; Salehi, P.; Aliahmadi, A.; Gohari, A.R.; Maroofi, H.; Ardalan, A. New Coumarin Derivatives from *Ferula Pseudalliacea* with Antibacterial Activity. *Nat. Prod. Res.* **2016**, *30*, 2747–2753. [\[CrossRef\]](#) [\[PubMed\]](#)
216. Pfaller, M.A.; Diekema, D.J. Epidemiology of Invasive Candidiasis: A Persistent Public Health Problem. *Clin. Microbiol. Rev.* **2007**, *20*, 133–163. [\[CrossRef\]](#) [\[PubMed\]](#)
217. Ghannadi, A.; Fattahian, K.; Shokoohinia, Y.; Behbahani, M.; Shahnoush, A. Anti-Viral Evaluation of Sesquiterpene Coumarins from *Ferula assa-foetida* against HSV-1. *Iran. J. Pharm. Res.* **2014**, *13*, 523–530. [\[CrossRef\]](#) [\[PubMed\]](#)
218. Rezaee, R.; Behravan, E.; Behravan, J.; Soltani, F.; Naderi, Y.; Emami, B.; Iranshahi, M. Antigenotoxic Activities of the Natural Dietary Coumarins Umbelliferone, Herniarin and 7-Isopentenylxy Coumarin on Human Lymphocytes Exposed to Oxidative Stress. *Drug Chem. Toxicol.* **2014**, *37*, 144–148. [\[CrossRef\]](#) [\[PubMed\]](#)
219. Raafat, K.; El-Lakany, A. Acute and Subchronic In-Vivo Effects of *Ferula hermonis* L. and *Sambucus nigra* L. and Their Potential Active Isolates in a Diabetic Mouse Model of Neuropathic Pain. *BMC Complement. Altern. Med.* **2015**, *15*, 257. [\[CrossRef\]](#)
220. Naji Reyhani Garmroudi, S.; Karimi, E.; Oskoueian, E.; Homayouni-Tabrizi, M.; Iranshahi, M. Ferutinin: A Phytoestrogen from *Ferula* and Its Anticancer, Antioxidant, and Toxicity Properties. *J. Biochem. Mol. Toxicol.* **2021**, *35*, e22713. [\[CrossRef\]](#)
221. Soltani, F.; Mosaffa, F.; Iranshahi, M.; Karimi, G.; Malekaneh, M.; Haghighi, F.; Behravan, J. Evaluation of Antigenotoxicity Effects of Umbelliprenin on Human Peripheral Lymphocytes Exposed to Oxidative Stress. *Cell Biol. Toxicol.* **2009**, *25*, 291–296. [\[CrossRef\]](#)
222. Mahaki, H.; Tanzadehpanah, H.; Abou-Zied, O.K.; Moghadam, N.H.; Bahmani, A.; Salehzadeh, S.; Dastan, D.; Saidijam, M. Cytotoxicity and Antioxidant Activity of Kamolonol Acetate from *Ferula pseudalliacea*, and Studying Its Interactions with Calf Thymus DNA (Ct-DNA) and Human Serum Albumin (HSA) by Spectroscopic and Molecular Docking Techniques. *Process Biochem.* **2019**, *79*, 203–213. [\[CrossRef\]](#)
223. Kogure, K.; Yamauchi, I.; Tokumura, A.; Kondou, K.; Tanaka, N.; Takaishi, Y.; Fukuzawa, K. Novel Antioxidants Isolated from Plants of the Genera *Ferula*, *inula*, *Prangos* and *Rheum* Collected in Uzbekistan. *Phytomedicine* **2004**, *11*, 645–651. [\[CrossRef\]](#) [\[PubMed\]](#)
224. Sayed, A.A.R. Ferulsinaic Acid Attenuation of Advanced Glycation End Products Extends the Lifespan of *Caenorhabditis Elegans*. *J. Pharm. Pharmacol.* **2011**, *63*, 423–428. [\[CrossRef\]](#) [\[PubMed\]](#)

225. Sajjadi, M.; Karimi, E.; Oskoueian, E.; Iranshahi, M.; Neamati, A. Galbanic Acid: Induced Antiproliferation in Estrogen Receptor-Negative Breast Cancer Cells and Enhanced Cellular Redox State in the Human Dermal Fibroblasts. *J. Biochem. Mol. Toxicol.* **2019**, *33*, e22402. [[CrossRef](#)] [[PubMed](#)]
226. Geroushi, A.; Auzi, A.A.; Elhwuegi, A.S.; Elzawam, F.; Elsherif, A.; Nahar, L.; Sarker, S.D. Antiinflammatory Sesquiterpenes from the Root Oil of *Ferula hermonis*. *Phytother. Res.* **2011**, *25*, 774–777. [[CrossRef](#)] [[PubMed](#)]
227. Zamani Taghizadeh Rabe, S.; Iranshahi, M.; Mahmoudi, M. In Vitro Anti-Inflammatory and Immunomodulatory Properties of Umbelliprenin and Methyl Galbanate. *J. Immunotoxicol.* **2016**, *13*, 209–216. [[CrossRef](#)] [[PubMed](#)]
228. Iranshahi, M.; Askari, M.; Sahebkar, A.; Hadjipavlou-Litina, D. Evaluation of Antioxidant, Anti-Inflammatory and Lipoxygenase Inhibitory Activities of the Prenylated Coumarin Umbelliprenin. *Daru-J. Pharm. Sci.* **2009**, *17*, 99–103. [[CrossRef](#)]
229. Kohno, S.; Murata, T.; Sugiura, A.; Ito, C.; Iranshahi, M.; Hikita, K.; Kaneda, N. Methyl Galbanate, a Novel Inhibitor of Nitric Oxide Production in Mouse Macrophage RAW264.7 Cells. *J. Nat. Med.* **2011**, *65*, 353–359. [[CrossRef](#)] [[PubMed](#)]
230. Wang, H.; Liu, Y.; Wang, Y.; Xu, T.; Xia, G.; Huang, X. Umbelliprenin Induces Autophagy and Apoptosis While Inhibits Cancer Cell Stemness in Pancreatic Cancer Cells. *Cancer Med.* **2023**, *12*, 15277–15288. [[CrossRef](#)]
231. Gholami, O.; Jeddi-Tehrani, M.; Iranshahi, M.; Zarnani, A.H.; Ziai, S.A. Umbelliprenin from *Ferula szowitsiana* Activates Both Intrinsic and Extrinsic Pathways of Apoptosis in Jurkat T-CLL Cell Line. *Iran. J. Pharm. Res.* **2013**, *12*, 371–376. [[CrossRef](#)]
232. Ziai, S.A.; Gholami, O.; Iranshahi, M.; Zamani, A.H.; Jeddi-Tehrani, M. Umbelliprenin Induces Apoptosis in CLL Cell Lines. *Iran. J. Pharm. Res.* **2012**, *11*, 653–659. [[CrossRef](#)]
233. Zhang, L.; Si, J.; Li, G.; Li, X.; Zhang, L.; Gao, L.; Huo, X.; Liu, D.; Sun, X.; Cao, L. Umbelliprenin and Lariciresinol Isolated from a Long-Term-Used Herb Medicine *Ferula sinkiangensis* Induce Apoptosis and G0/G1 Arresting in Gastric Cancer Cells. *RSC Adv.* **2015**, *5*, 91006–91017. [[CrossRef](#)]
234. Iranshahi, M.; Sahebkar, A.; Takasaki, M.; Konoshima, T.; Tokuda, H. Cancer Chemopreventive Activity of the Prenylated Coumarin, Umbelliprenin, in Vivo. *Eur. J. Cancer Prev.* **2009**, *18*, 412–415. [[CrossRef](#)] [[PubMed](#)]
235. Kim, Y.H.; Shin, E.A.; Jung, J.H.; Park, J.E.; Koo, J.; Koo, J.I.; Shim, B.S.; Kim, S.-H. Galbanic Acid Potentiates TRAIL Induced Apoptosis in Resistant Non-Small Cell Lung Cancer Cells via Inhibition of MDR1 and Activation of Caspases and DR5. *Eur. J. Pharmacol.* **2019**, *847*, 91–96. [[CrossRef](#)] [[PubMed](#)]
236. Zhang, Y.; Kim, K.-H.; Zhang, W.; Guo, Y.; Kim, S.-H.; Lü, J. Galbanic Acid Decreases Androgen Receptor Abundance and Signaling and Induces G1 Arrest in Prostate Cancer Cells. *Int. J. Cancer* **2012**, *130*, 200–212. [[CrossRef](#)] [[PubMed](#)]
237. Folkman, J. Fundamental Concepts of the Angiogenic Process. *Curr. Mol. Med.* **2003**, *3*, 643–651. [[CrossRef](#)] [[PubMed](#)]
238. Harper, J.; Moses, M.A. Molecular Regulation of Tumor Angiogenesis: Mechanisms and Therapeutic Implications. In *Cancer: Cell Structures, Carcinogens and Genomic Instability*; Birkhäuser: Basel, Switzerland, 2006; pp. 223–268.
239. Kim, K.-H.; Lee, H.-J.; Jeong, S.-J.; Lee, H.-J.; Lee, E.-O.; Kim, H.-S.; Zhang, Y.; Ryu, S.-Y.; Lee, M.-H.; Lü, J.; et al. Galbanic Acid Isolated from *Ferula assafoetida* Exerts In Vivo Anti-Tumor Activity in Association with Anti-Angiogenesis and Anti-Proliferation. *Pharm. Res.* **2011**, *28*, 597–609. [[CrossRef](#)] [[PubMed](#)]
240. Mannello, F.; Gazzanelli, G. Tissue Inhibitors of Metalloproteinases and Programmed Cell Death: Conundrums, Controversies and Potential Implications. *Apoptosis* **2001**, *6*, 479–482. [[CrossRef](#)]
241. Shahcheraghi, S.H.; Lotfi, M.; Soukhtanloo, M.; Ghayour-Mobarhan, M.; Jaliani, H.Z.; Sadeghnia, H.R.; Ghorbani, A. Effects of Galbanic Acid on Proliferation, Migration, and Apoptosis of Glioblastoma Cells Through the PI3K/Akt/MTOR Signaling Pathway. *Curr. Mol. Pharmacol.* **2020**, *14*, 79–87. [[CrossRef](#)]
242. Fallah, J.; Rini, B.I. HIF Inhibitors: Status of Current Clinical Development. *Curr. Oncol. Rep.* **2019**, *21*, 6. [[CrossRef](#)]
243. Wicks, E.E.; Semenza, G.L. Hypoxia-Inducible Factors: Cancer Progression and Clinical Translation. *J. Clin. Investig.* **2022**, *132*, e159839. [[CrossRef](#)]
244. Eskandani, M.; Abdolalizadeh, J.; Hamishehkar, H.; Nazemiyeh, H.; Barar, J. Galbanic Acid Inhibits HIF-1 α Expression via EGFR/HIF-1 α Pathway in Cancer Cells. *Fitoterapia* **2015**, *101*, 1–11. [[CrossRef](#)] [[PubMed](#)]
245. Syeda, S.; Abbas, S.; Naveed, D.S.; Qamar, F.; Saba, Hussain, J.; Jawed, S. Exploration & Wonders of Medicinal Plant “Asafoetida” (Heeng) Constituent Galbanic Acid as Anticancerous and Antimutagenic Agent. *IESRJ* **2015**, *1*, 15–17.
246. Eruçar, F.M.; Kuran, F.K.; Altıparmak Ülbegi, G.; Özbey, S.; Karavuş, Ş.N.; Arcan, G.G.; Yazıcı Tütüniş, S.; Tan, N.; Aksoy Sağır, P.; Miski, M. Sesquiterpene Coumarin Ethers with Selective Cytotoxic Activities from the Roots of *Ferula huber-morathii* Peşmen (Apiaceae) and Unequivocal Determination of the Absolute Stereochemistry of Samarcandin. *Pharmaceuticals* **2023**, *16*, 792. [[CrossRef](#)] [[PubMed](#)]
247. Zhang, X.; Li, M.; Wang, Y.; Chen, Q.; Liu, J.; Li, S.; Tang, S. Study on chemical constituents from *Ferula teterrima*. *Chin. Tradit. Herbal Drugs* **2023**, *54*, 2373–2378.
248. Amin, A.; Tuenter, E.; Cos, P.; Maes, L.; Exarchou, V.; Apers, S.; Pieters, L. Antiprotozoal and Antiglycation Activities of Sesquiterpene Coumarins from *Ferula narthex* Exudate. *Molecules* **2016**, *21*, 1287. [[CrossRef](#)] [[PubMed](#)]
249. Amin, A.; Hanif, M.; Rafey, A.; Zaib, S.; Bakhsh, S.; Ramzan, M.; Zaman, A.; Rehman, F.U.; Iqbal, J.; Pieters, L. Sesquiterpene Coumarins from *Ferula narthex* 15-LOX, α -Glucosidase Inhibition and Molecular Docking Studies. *Rev. Bras. Farmacogn.* **2020**, *30*, 12–17. [[CrossRef](#)]
250. Foadoddini, M.; Javdani, H.; Farahi, A.; Hosseini, M. Therapeutic Potential of *Ferula foetida* (Bunge) Regel on Gastric Ulcer Model in Rats. *Environ. Sci. Pollut. Res. Int.* **2022**, *29*, 12147–12156. [[CrossRef](#)] [[PubMed](#)]

251. Qin, J.; Si, J.; Jia, B.; Zhao, C.; Zhou, D.; He, X.; Wang, C.; Zhu, X. Water Use Strategies of *Ferula bungeana* on Mega-Dunes in the Badain Jaran Desert. *Front. Plant Sci.* **2022**, *13*, 957421. [[CrossRef](#)]
252. Qin, H.-H.; Cai, J.; Liu, C.-K.; Zhou, R.-X.; Price, M.; Zhou, S.-D.; He, X.-J. The Plastid Genome of Twenty-Two Species from *Ferula*, *Talassia*, and *Soranthus*: Comparative Analysis, Phylogenetic Implications, and Adaptive Evolution. *BMC Plant Biol.* **2023**, *23*, 9. [[CrossRef](#)]
253. Farhadi, F.; Iranshahi, M.; Mohtashami, L.; Shakeri Asil, S.; Iranshahy, M. Metabolic Differences of Two *Ferula* Species as Potential Sources of Galbanum: An NMR-Based Metabolomics Study. *Phytochem. Anal.* **2021**, *32*, 811–819. [[CrossRef](#)]
254. Hasanpour, M.; Safari, H.; Mohammadpour, A.H.; Iranshahy, M.; Dehghan Nayyeri, M.J.; Farhadi, F.; Emami, B.; Iranshahi, M. Efficacy of Covexir® (*Ferula foetida* Oleo-Gum) Treatment in Symptomatic Improvement of Patients with Mild to Moderate COVID-19: A Randomized, Double-Blind, Placebo-Controlled Trial. *Phytother. Res.* **2022**, *36*, 4504–4515. [[CrossRef](#)]
255. Abu-Izneid, T.; Rauf, A.; Shariati, M.A.; Khalil, A.A.; Imran, M.; Rebezov, M.; Uddin, M.S.; Mahomoodally, M.F.; Rengasamy, K.R.R. Sesquiterpenes and Their Derivatives-Natural Anticancer Compounds: An Update. *Pharmacol. Res.* **2020**, *161*, 105165. [[CrossRef](#)]

Disclaimer/Publisher’s Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.