

Table S1. Distribution of EORTC scores according to the variables of interest. Data are median [I quartile-III quartile].

	Functional Scales			Symptomatic Scales		
	Global QoL	Physical Functioning	Social Functioning	Disease-Related Worries	Pain	Upper-GI Symptoms
	Median (IQR)	Median (IQR)	Median (IQR)	Median (IQR)	Median (IQR)	Median (IQR)
Gender						
Male	0.83 (0.67-1.00)	1.00 (0.94-1.00)	0.94 (0.80-1.00)	0.25 (0.08-0.33)	0.05 (0.00-0.10)	0.03 (0.00-0.09)
Female	0.75 (0.58-0.88)	0.87 (0.72-0.94)	0.89 (0.78-0.94)	0.27 (0.16-0.40)	0.10 (0.05-0.23)	0.06 (0.00-0.09)
Age						
>= 65 y	0.67 (0.50-0.83)	0.83 (0.67-0.94)	0.89 (0.76-0.94)	0.27 (0.17-0.40)	0.10 (0.05-0.24)	0.06 (0.03-0.15)
< 65 y	0.83 (0.67-1.00)	0.94 (0.89-1.00)	0.89 (0.83-0.94)	0.25 (0.08-0.33)	0.05 (0.00-0.14)	0.03 (0.00-0.08)
Comorbidities						
No	0.67 (0.67-1.00)	1.00 (0.94-1.00)	0.89 (0.83-0.94)	0.27 (0.13-0.33)	0.05 (0.00-0.10)	0.06 (0.03-0.09)
Single	0.83 (0.73-1.00)	0.94 (0.83-1.00)	0.94 (0.83-0.97)	0.17 (0.07-0.29)	0.08 (0.04-0.15)	0.03 (0.00-0.06)
Multiple	0.83 (0.58-0.92)	0.89 (0.72-1.00)	0.89 (0.72-0.94)	0.27 (0.17-0.40)	0.10 (0.00-0.24)	0.05 (0.00-0.12)
Type of NEN						
NF	0.83 (0.58-1.00)	0.94 (0.75-1.00)	0.89 (0.78-0.94)	0.27 (0.17-0.40)	0.05 (0.00-0.19)	0.03 (0.00-0.09)
F	0.83 (0.67-1.00)	0.94 (0.78-1.00)	0.89 (0.82-0.94)	0.25 (0.08-0.33)	0.10 (0.05-0.19)	0.06 (0.00-0.09)
Tumor grade						
G2	0.83 (0.79-1.00)	0.94 (0.88-1.00)	0.91 (0.88-0.94)	0.20 (0.13-0.37)	0.05 (0.00-0.10)	0.03 (0.00-0.06)
G1	0.75 (0.58-0.92)	0.94 (0.78-1.00)	0.89 (0.78-0.94)	0.27 (0.13-0.33)	0.10 (0.00-0.19)	0.06 (0.00-0.09)
Type of surgery						
Limited	0.83 (0.67-1.00)	0.94 (0.82-1.00)	0.89 (0.83-1.00)	0.20 (0.07-0.27)	0.10 (0.00-0.19)	0.03 (0.00-0.07)
Standard	0.83 (0.58-0.92)	0.94 (0.67-1.00)	0.89 (0.72-0.94)	0.27 (0.17-0.40)	0.08 (0.00-0.19)	0.06 (0.00-0.12)

Legend. QoL, quality of life. GI, gastrointestinal. IQR, interquartile range. NEN, neuroendocrine neoplasm. NF, nonfunctioning. F, functioning.