



Supplementary Materials

880 nm NIR-Triggered Organic Small Molecular-Based Nanoparticles for Photothermal Therapy of Tumor

Yunying Zhao ^{1,†}, Zheng He ^{1,†}, Qiang Zhang ¹, Jing Wang ¹, Wenying Jia ¹, Long Jin ¹, Linlin Zhao ^{1,2,*} and Yan Lu ¹

¹ School of Materials Science & Engineering, Tianjin Key Laboratory for Photoelectric Materials and Devices, Key Laboratory of Display Materials & Photoelectric Devices, Ministry of Education, Tianjin University of Technology, Tianjin 300384, China; 17853483674@163.com (Y.Z.); 13752723282@163.com (Z.H.); zhangqiang@email.tjut.edu.cn (Q.Z.); wangjing@iccas.ac.cn (J.W.); jia15249238411@163.com (W.J.); KimYong0205@163.com (L.J.); luyan@tjut.edu.cn (Y.L.)

² State Key Laboratory of Molecular Engineering of Polymers, Fudan University, Shanghai 200433, China

* Correspondence: linlinzhao@email.tjut.edu.cn

† These authors contributed equally to this work.




Figure S1. Zeta potential of TNPs in water.




Figure S2. Changes of hydrodynamic diameters of TNPs in DMEM with time, [TNPs] = 180 $\mu\text{g/mL}$.




Figure S3. (A) Temperature elevation of TNPs (180 $\mu\text{g/mL}$) under 880 nm irradiation at 0.7 W/cm² for 5 min, followed by subsequent cooling to room temperature and (B) Linear time data versus- $\ln(\theta)$ obtained from the cooling period of NIR laser off.