

Article

Effect of Solvent on Fluorescence Emission from Polyethylene Glycol-Coated Graphene Quantum Dots under Blue Light Illumination

Po-Chih Yang ^{1,*}, Yu-Xuan Ting ¹, Siyong Gu ², Yasser Ashraf Gandomi ³, Jianlin Li ⁴, Chien-Te Hsieh ^{1,*}

¹ Department of Chemical Engineering and Materials Science, Yuan Ze University, Taoyuan 32003, Taiwan; danny0935331003@gmail.com

² Fujian Provincial Key Laboratory of Functional Materials and Applications, School of Materials Science and Engineering, Xiamen University of Technology, Xiamen 361024, China; gu-siyong@163.com

³ Department of Chemical Engineering, Massachusetts Institute of Technology, Cambridge, MA 02142, USA; ygandomi@mit.edu

⁴ Electrification and Energy Infrastructure Division, Oak Ridge National Laboratory, Oak Ridge, TN 37831, USA; lij4@ornl.gov

* Correspondence: pcyang@saturn.yzu.edu.tw (P.-C.Y.); cthsieh@saturn.yzu.edu.tw (C.-T.H.)




Figure S1. Particle size distributions of different GQD samples determined from DLS technique.




Figure S2. Survey-scan XPS spectra of pristine GQD and PEG-coated GQD cluster samples.




Figure S3. FT-IR spectra of pristine GQD and PEG-coated GQD cluster samples.




Figure S4. Photographs of different GQD suspensions at 450 nm: (a) water, (b) ethanol, and (c) PGMEA.