

Supplementary Material

Comprehensive Study of the Chemistry behind the Stability of Carboxylic SWCNT Dispersions in the Development of a Transparent Electrode

Jovana Stanojev ^{1,*}, Stevan Armaković ², Sara Joksović ¹, Branimir Bajac ¹, Jovan Matović ¹ and Vladimir V. Srdić ³

¹ BioSense Institute, University of Novi Sad, Dr Zorana Djindjica 1, 21000 Novi Sad, Serbia;

sara.joksovic@biosense.rs (S.J.); branimir.bajac@biosense.rs (B.B.); jovan.matovic@biosense.rs (J.M.)

² Department of Physics, Faculty of Sciences, University of Novi Sad, Trg Dositeja Obradovica 4, 21000 Novi Sad, Serbia; stevan.armakovic@df.uns.ac.rs

³ Faculty of Technology Novi Sad, University of Novi Sad, Bulevar Cara Lazara 1, 21000 Novi Sad, Serbia; srdicvv@uns.ac.rs

* Correspondence: jovana.stanojev@biosense.rs; Tel.: +381-214852137 In the supplementary material is given the Raman spectra of SWCNT-COOH, analyzed with the red laser (780 nm) (Figure S1).




Figure S1. Raman spectra of SWCNT-COOH after the HCl treatment, analyzed with the red laser (780 nm).