Biocompatible Materials Based on Self-Assembling Peptides on Ti25Nb10Zr Alloy: Molecular Structure and Organization Investigated by Synchrotron Radiation Induced Techniques.

Valeria Secchi¹, Stefano Franchi^{1,*,†}, Marta Santi¹, Alina Vladescu², Mariana Braic², Tomáš Skála³, Jaroslava Nováková³, Monica Dettin⁴, Annj Zamuner⁴, Giovanna Iucci¹ and Chiara Battocchio^{1,*}

- ¹ Department of Science, Roma Tre University of Rome, Via della Vasca Navale 79, 00146 Rome, Italy; valeria.secchi@uniroma3.it (V.S.); santimarta3@gmail.com (M.S.); giovanna.iucci@uniroma3.it (G.I.)
- ² National Institute for Optoelectronics, 409 Atomistilor St., 077125 Magurele, Romania; alinava@inoe.ro (A.V.); mariana.braic@inoe.ro (M.B.)
- ³ Department of Surface and Plasma Science, Faculty of Mathematics and Physics, Charles University, V Holešovičkách 2, 18000 Prague, Czech Republic; tomas.skala@elettra.eu (T.S.); jaroslava.lavkova@gmail.com (J.N.)
- ⁴ Department of Industrial Engineering, University of Padua, Via Marzolo, 9, Padua 35131, Italy; monica.dettin@unipd.it (M.D.); annj.zamuner@studenti.unipd.it (A.Z.)
- * Correspondence: stefano.franchi@elettra.eu (S.F.); chiara.battocchio@uniroma3.it (C.B.); Tel.: +39-040-3758059 (S.F.); +39-06-5733-3400 (C.B.)
- + Present Address: Elettra-Sincrotrone Trieste S.C.p.A. di interesse nazionale, Strada Statale 14-km 163,5 in AREA Science Park, 34149 Basovizza, Trieste, Italy.

Supplementary Material

Figure S1. Total EDS mapping spectrum and relative distribution of the concentration of each constituent element on the Ti25Nb10Zr alloy surface performed in one scanned zone on the surface (zone 4 of Table S1)



Table S1. The elemental composition of the alloy determined by EDS measurements

Zone	Ti (wt%)	Nb(wt %)	Zr (wt %)
Zone 1	68 7	21.9	94
Zone 1	00.7	21.9	2.4
Zone 2	69.2	21.6	9.2
Zone 3	68.7	22.1	9.2
Zone 4	68.8	21.8	9.4
Average	68.85±0.24	21.85±0.21	9.3±0.11

Table S2. Binding Energies in eV of the atomic species present on Clean Surface Sample and relative atomic ratios. Me-Ox: oxygen of metal oxides; O-org: organic oxygen.

		С	1s			O1s				
	C-C	C-0	C=O	СООН	Me-Ox	O-org	H ₂ O	Ti2p _{3/2}	Nb3d _{5/2}	Zr3d _{5/2}
B.E. (eV)	285.0	286.4	287.9	288.1	530.1	531.8	533.3	458.5	207.1	182.2
FWH M (eV)	1.6	1.6	1.6	1.6	1.8	1.8	1.8	1.8	1.7	1.6
		C-0/	С=О/	COOH/		O org/	C tot			
		C-C	C-C	C-C		TiO ₂	/Ti	Ti/Nb	Ti/Zr	Nb/Zr
Atomi c Ratio (%)		31	12	13		61	676	380	522	159

Table S3. Binding Energies in eV of the atomic species present on MUL samples surface. N+: protonated nitrogen; Me-Ox: oxygen of metaloxides; O-org: organic oxygen; ----: not detected.

	C1s			N1s		O1s						
MUL	C-C	C-N	N-C=O	СООН	Ν	N+	M-Ox	O-org	H ₂ O	Ti2p _{3/2}	Nb3d5/2	Zr3d5/2
sample	(eV)	(eV)	(eV)	(eV)								
pH 2	285.0	286.3	288.1	289.1	400.1	401.7		531.4	533.3			
FWHM	1.5	1.5	1.5	1.5	1.6	1.6		2.2	2.2			
pH 4	285.0	286.3	288.3	289.5	400.2	401.8		531.7	533.5			
FWHM	1.6	1.6	1.6	1.6	1.6	1.6		2.1	2.1			
pH 7	285.0	286.3	287.6	289.0	401.3		530.8	532.7	534.2	459.5	208.2	
FWHM	1.6	1.6	1.6	1.6	1.9		2.0	2.0	2.0	2.3	1.9	
pH 10	285.0	286.3	288.2	289.3	400.1	401.8		531.2	533.0			
FWHM	1.6	1.6	1.6	1.6	1.6	1.6		2.1	2.1			
pH 12	285.0	286.3	288.1	289.1	400.2	401.9	530.8	532.6	534.6	458.5	207.1	182.1
FWHM	1.6	1.6	1.6	1.6	1.6	1.6	2.3	2.3	2.3	1.9	2.0	2.1

	C1s				N1s			O1s				
ML	C-C	C-N	N-C=O	COOH		N+	M-O _x	O-org	H ₂ O	Ti 2p 3/2	Nb 3d	Zr 3d
Sample	(eV)	(eV)	(eV)	(eV)	N (eV)	(eV)	(eV)	(eV)	(eV)	(eV)	5/2 (eV)	5/2 (eV)
pH 2	255.0	286.4	288.1	289.2	400.2	401.8	530.2	531.7	533.3	458.8	207.1	182.2
FWHM	1.6	1.6	1.6	1.6	1.6	1.6	1.9	1.9	1.9	1.9	1.6	1.4
pH 4	285.0	288.0	286.3	289.1	400.0	401.4	530.3	531.6	533.2	458.5	207.0	182.1
FWHM	1.5	1.5	1.5	1.5	1.5	1.5	2.1	2.1	2.1	1.7	2.3	2.1
pH 7	285.0	286.3	287.3	289.0	400.1	401.4	530.3	531.9	533.5	458.6	207.3	182.3
FWHM	1.6	1.6	1.6	1.6	1.7	1.7	2.0	2.0	2.0	1.9	1.7	1.7
pH 10	285.0	286.3	288.0	289.2	400.2		530.4	531.7	533.3	458.7	207.3	182.4
FWHM	1.6	1.6	1.6	1.6	1.8		2.1	2.1	2.1	1.8	1.5	1.8
pH 12	285.0	286.4	287.9	289.1	400.3		530.6	532.1	533.7	458.7	207.4	182.5
FWHM	1.6	1.6	1.6	1.6	1.8		2.0	2.0	2.0	1.7	1.7	2.0

Table S4. Binding Energies in eV of the atomic species present on ML sample surface. N+: protonated nitrogen; Me-Ox: oxygen of metaloxides; O-org: organic oxygen.

Figure S2. AFM image of Clean Surface Sample

