Plasmon-Enhanced Blue-Light Emission of Stable Perovskite Quantum Dot Membranes

Kai Gu¹, Hongshang Peng^{1,2}, Siwei Hua¹, Yusong Qu¹ and Di Yang^{1,*}

* Correspondence: diyang@muc.edu.cn; Tel.: +86-1352-274-3208



Figure S1. TEM and HRTEM images of PVDF-MAPbBr_{1.2}Cl_{1.8} QDs nanofibers indicating that almost all PQDs are encapsulated inside the PVDF fibers.



Figure S2. (**a**) The diameter statistics of the PVDF-MAPbBr_{1.2}Cl_{1.8} fibers and (**b**) the size statistics of the MAPbBr_{1.2}Cl_{1.8} QDs.



Figure S3. Normalized PL spectra of PVDF-MAPbBr_{3-x} Cl_x with different the ratio of Cl⁻ and Br⁻.



Figure S4. (**a**) Schematic diagram of five points on the PVDF-PQDs membrane; (**b**) The PL spectra of five points on the membrane. The inset shows overlap of the PL spectra of these five points.



Figure S5. The optical images of (**a**) PVDF-MAPbBr_{1.2}Cl_{1.8} and (**b**) PVDF-MAPbBr₃ films on FTO substrates under ultraviolet irradiation at ~365 nm wavelength.