

Article

# Fragments of a Wheat Hevein-Like Antimicrobial Peptide Augment the Inhibitory Effect of a Triazole Fungicide on Spore Germination of *Fusarium oxysporum* and *Alternaria solani*


Larisa Shcherbakova <sup>1,\*</sup>, Tatyana Odintsova <sup>2,\*</sup>, Tatyana Pasechnik <sup>1</sup>, Lenara Arslanova <sup>1</sup>, Tatyana Smetanina <sup>1</sup>, Maxim Kartashov <sup>1</sup>, Marina Slezina <sup>2</sup> and Vitaly Dzhavakhiya <sup>1</sup>

<sup>1</sup> All-Russian Research Institute of Phytopathology, Bolshie Vyazemy, 143050 Moscow reg., Russia; beefarmer@yandex.ru (T.P.); linr-13@yandex.ru (L.A.); smetaneinatatyana@yandex.ru (T.S.); maki505@mail.ru (M.K.); dzhavakhiya@yahoo.com (V.D.)

<sup>2</sup> Vavilov Institute of General Genetics RAS, Gubkina Str. 3, 119333 Moscow, Russia; omey@list.ru

\* Correspondence: larisavniif@yahoo.com (L.S.); odintsova2005@rambler.ru (T.O.)

Received: 14 October 2020; Accepted: 2 December 2020; Published: date



**Figure S1.** Detached tomato seedlings in (a) sterile distilled water (SDW) and SDW supplemented with WAMP-2-derived peptides: (b) WAMP-C, (c) WAMP-G1 (each at a concentration of 200 µg/mL). Seedlings were photographed after 2 days of incubation.