

## Effectiveness of a Novel Compound HAIR & SCALP COMPLEX on Hair Follicle Regeneration

Greta Ferruggia <sup>1</sup>, Martina Contino <sup>1</sup>, Massimo Zimbone <sup>2</sup> and Maria Violetta Brundo <sup>1,\*</sup>

<sup>1</sup> Department of Biological, Geological and Environmental Sciences, University of Catania, Via Androne 81, 95124 Catania, Italy; greta.ferruggia@phd.unict.it (G.F.); martina.contino@phd.unict.it (M.C.)

<sup>2</sup> Institute for Microelectronics and Microsystems, National Research Council of Italy (CNR-IMM), 95123 Catania, Italy; massimo.zimbone@ct.infn.it

\* Correspondence: mariavioletta.brundo@unict.it; Tel.: +39-0957306039

### Supplementary materials

**Table S1.** Concentration of 20 bioactive molecules of GF20 characterized by ELISA test.

Bioactive molecules	Log Concentration (pg/mL)
GM-CSF	3.963 ± 0.38
bFGF	3.955 ± 0.49
TGF- $\beta$	3.5 ± 0.15
HGF	3.42 ± 0.15
IFN- $\gamma$	3.35 ± 0.38
SDF-1 $\alpha$	3.32 ± 0.3
VEGF	3.31 ± 0.426
IL-2	3.29 ± 0.3
IGF-1	3.1 ± 0.08
BMP-2	3.1 ± 0.27
IL-17A	3.04 ± 0.477
G-CSF	2.85 ± 0.27
PDGF	2.766 ± 0.354
NGF	2.622 ± 0.832
KGF	2.61 ± 0.4
EGF	2.41 ± 0.36
EOTAXIN	2.4 ± 0.37
TNF- $\alpha$	2.305 ± 0.356
IL-6	1.81 ± 0.3
IL-4	0.95 ± 0.049

Data were reported as Log Concentration ± standard deviation.