

Interactions Under Crowding Milieu: Chemical-Induced Denaturation of Myoglobin is Determined by the Extent of Heme Dissociation on Interaction with Crowders

Khalida Nasreen ^{1,†}, Zahoor Ahmad Parray ^{1,†}, Shahzaib Ahamad ¹, Faizan Ahmad ¹,
Anwar Ahmed ^{2,3}, Salman Freeh Alamery ², Tajamul Hussain ², Md. Imtaiyaz Hassan ¹ and
Asimul Islam ^{1,*}

¹ Centre for Interdisciplinary Research in Basic Sciences, Jamia Millia Islamia, Jamia Nagar, New Delhi 110025, India

² Center of Excellence in Biotechnology Research, College of Science, King Saud University, Riyadh, Saudi Arabia

³ Protein Research Chair, Department of Biochemistry, College of Science, King Saud University, Riyadh, Saudi Arabia

[†] Both the authors have contributed equally

^{*} Correspondence: aislam@jmi.ac.in; Tel.: 0091-9312812007




Figure S1. The docked complex of myoglobin (heme and its amino acid residues) with dextran 70 (dimer).




Figure S2. Two-dimensional plot interaction of myoglobin (heme and its amino acid residues) with dextran 70 (dimer) generated by maestro (schrodinger).