



Average pseudospectra. Principal Component Analysis. Top: loadings of components 1, 2 and 3 accounting for 89.3, 6.7, 2.1 % of the total variance, respectively. Bottom: Scatter plots of components 1-2 (left) and 2-3 (right).



Large PCA. Principal Component 2. Montages of the 40 score images of the series. Field of view of each image montage: 7.89 cm x 11.45 cm corresponding roughly to 2 x 2 cm<sup>2</sup> per section. Intensities can be compared.

Montage 1: images 1 to 12: inbred line M21 with 2, 2, 3, 3 and 2 sections per internode, respectively. Images 13 to 20: inbred line M22 with 2 sections per internode.


Montage 2: images 1 to 2: inbred line M22 with 2 sections per internode. images 3 to 12: inbred line M23 with 2 sections per internode. Images 13 to 20: inbred line M24 with 2 sections per internode. Only 4 internodes could be analysed for this line



Large PCA. Principal Component 2. Montages of the 40 rind zoomed score images of the series. Fields of view: one image: 4.17\*4.17 mm<sup>2</sup>, montage: 1.89 cm x 2.28 cm. Intensities can be compared.

Montage 1: images 1 to 12: inbred line M21 with 2, 2, 3, 3 and 2 sections per internode, respectively. Images 13 to 20: inbred line M22 with 2 sections per internode.

Montage 2: images 1 to 2: inbred line M22 with 2 sections per internode. images 3 to 12: inbred line M23 with 2 sections per internode. Images 13 to 20: inbred line M24 with 2 sections per internode. Only 4 internodes could be analysed for this line



Large PCA. Principal Component 2. Montages of the 40 centre zoomed score images of the series. Fields of view: one image: 4.17\*4.17 mm<sup>2</sup>, montage: 1.89 cm x 2.28 cm. Intensities can be compared.

Montage 1: images 1 to 12: inbred line M21 with 2, 2, 3, 3 and 2 sections per internode, respectively. Images 13 to 20: inbred line M22 with 2 sections per internode.

Montage 2: images 1 to 2: inbred line M22 with 2 sections per internode. images 3 to 12: inbred line M23 with 2 sections per internode. Images 13 to 20: inbred line M24 with 2 sections per internode. Only 4 internodes could be analysed for this line.