

Article

Lie Symmetry Analysis of the Black-Scholes-Merton Model for European Options with Stochastic Volatility

Andronikos Paliathanasis ^{1,*}, K. Krishnakumar ^{2,†}, K.M. Tamizhmani ^{2,†} and Peter G.L. Leach ^{3,4,5,†}

¹ Instituto de Ciencias Físicas y Matemáticas, Universidad Austral de Chile, Valdivia 5090000, Chile

² Department of Mathematics, Pondicherry University, Kalapet 605014, India; krishapril09@gmail.com (K.K.); kmtmani54@gmail.com (K.M.T.)

³ Institute of Systems Science, Department of Mathematics, Durban University of Technology, Durban 4000, South Africa; leach@ucy.ac.cy

⁴ School of Mathematics, Statistics and Computer Science, University of KwaZulu-Natal, Durban 4000, South Africa

⁵ Department of Mathematics and Statistics, University of Cyprus, Lefkosia 1678, Cyprus

* Correspondence: anpaliat@phys.uoa.gr; Tel.: +56-63-2221200

† These authors contributed equally to this work.

Academic Editor: Indranil SenGupta

Received: 30 January 2016; Accepted: 15 April 2016; Published: 3 May 2016

Abstract: We perform a classification of the Lie point symmetries for the Black-Scholes-Merton Model for European options with stochastic volatility, σ , in which the last is defined by a stochastic differential equation with an Orstein-Uhlenbeck term. In this model, the value of the option is given by a linear (1 + 2) evolution partial differential equation in which the price of the option depends upon two independent variables, the value of the underlying asset, S , and a new variable, y . We find that for arbitrary functional form of the volatility, $\sigma(y)$, the (1 + 2) evolution equation always admits two Lie point symmetries in addition to the automatic linear symmetry and the infinite number of solution symmetries. However, when $\sigma(y) = \sigma_0$ and as the price of the option depends upon the second Brownian motion in which the volatility is defined, the (1 + 2) evolution is not reduced to the Black-Scholes-Merton Equation, the model admits five Lie point symmetries in addition to the linear symmetry and the infinite number of solution symmetries. We apply the zeroth-order invariants of the Lie symmetries and we reduce the (1 + 2) evolution equation to a linear second-order ordinary differential equation. Finally, we study two models of special interest, the Heston model and the Stein-Stein model.

Keywords: lie point symmetries; financial mathematics; stochastic volatility; Black-Scholes-Merton equation

MSC: 22E60; 35Q91

1. Introduction

The Black-Scholes-Merton Model for European options is based upon some Ansatz for the stock price. Specifically, the process for the stock price is characterized by continuity, and it has the ability to hedge continuously with transaction costs and has constant volatility [1–3].

In the Black-Scholes-Merton Model, the price of a financial asset is given by the solution of the stochastic differential equation

$$dS_t = rS_t dt + \sigma S_t dW_t \quad (1)$$

where W_t is a Brownian motion, and the value $u = u(t, S)$ of the option is given by the solution of the (1 + 1) evolution equation,

$$\frac{1}{2}\sigma^2 S^2 u_{,SS} + rSu_{,S} - ru + u_{,t} = 0 \tag{2}$$

in which t is time, S is the current value of the underlying asset, for example a stock price, and r is the rate of return on a safe investment. The value of the option is subject to the satisfaction of the terminal condition, $u(T, S) = U(S)$, when $t = T$. Finally, σ is the volatility of the model.

The Black-Scholes-Merton Model assumes constant volatility σ . However, in real problems, σ is not a constant. One possible generalisation of the model, Equation (2), is to consider that the volatility depends upon the time, t , and on the value of the stock, S , i.e., $\sigma = \sigma(t, S)$. It has been proposed that σ is a function of a mean Orstein-Uhlenbeck process [4].

Consider that $\sigma = f(y)$, where y is given by the stochastic differential equation with the Orstein-Uhlenbeck term [5-7]:

$$dy_t = \alpha(m - y_t) dt + \beta d\hat{Z}_t \tag{3}$$

The new Brownian motion \hat{Z}_t can be correlated with W_t and be expressed as follows:

$$\hat{Z}_t = \rho W_t + \sqrt{1 - \rho^2} Z_t \tag{4}$$

in which Z_t describes a Brownian motion independent of W_t and ρ is the correlation factor with values $|\rho| \leq 1$.

Hence, the Black-Scholes Equation (2) in the case of stochastic volatility is modified and the value u of the option is given by the (1 + 2) evolution equation

$$(\hat{M}_1 + \hat{M}_2 + \hat{M}_3 + \hat{M}_4) u(t, S, y) = 0 \tag{5}$$

where the operators, $\hat{M}_1, \hat{M}_2, \hat{M}_3, \hat{M}_4$, are defined as follows:

$$\hat{M}_1 = \frac{1}{2}f^2(y) S^2 \frac{\partial^2}{\partial S^2} + rS \frac{\partial}{\partial S} - r + \frac{\partial}{\partial t} \tag{6}$$

$$\hat{M}_2 = \rho\beta S f(y) \frac{\partial^2}{\partial S \partial y}, \quad \hat{M}_3 = -\beta \Lambda(t, S, y) \frac{\partial}{\partial y}, \quad \text{and} \tag{7}$$

$$\hat{M}_4 = \frac{1}{2}\beta^2 \frac{\partial^2}{\partial y^2} + \alpha(m - y) \frac{\partial}{\partial y} \tag{8}$$

The function $\Lambda(t, S, y)$ is

$$\Lambda(t, S, y) = \rho \frac{\mu - r}{f(y)} + \gamma(t, S, y) \sqrt{1 - \rho^2} \tag{9}$$

and $u(t, S, y)$ satisfies the terminal condition $u(T, S, y) = U(S)$ at time $t = T$.

The operator \hat{M}_1 gives the Black-Scholes-Merton Equation (2) with volatility $\sigma = f(y)$, \hat{M}_2 expresses the correlation term between the two Brownian motions, W_t and \hat{Z}_t , of the European option and of the volatility, respectively, and \hat{M}_4 is the Orstein-Uhlenbeck process term. Finally, the term \hat{M}_3 , the so called premium term, expresses the market price of the volatility risk [6]. The function $\gamma(t, S, y)$ in Equation (9) is the risk-premium factor which drives the volatility and follows from the second Brownian motion, Z_t , where in the case of absolute correlation, i.e., $|\rho| = 1$, $\gamma(t, S, y)$ does not play any role in the model. The first term of the *rhs* side of Equation (9) is called the excess return-to-risk ratio [6]. The statistical importance of stochastic volatility has been confirmed in [8].

The purpose of this work is the study of the Black-Scholes-Merton Model with stochastic volatility, Equation (5), by using the method of group invariant transformations, specifically the Lie (point) symmetries of the equation. The importance of Lie symmetries is that they provide a systematic method

to facilitate the solution of differential equations because they provide first-order invariants which can be used to reduce the differential equations. Moreover, Lie symmetries can be used for the classification of differential equations. Furthermore, we can extract important information for the differential equation, consequently for the model, from the group of invariant transformations admitted.

The first application of the Lie symmetries in financial modeling was performed by Gazizov & Ibragimov in [9]. They studied the admitted group of invariant transformations for the Black-Scholes-Merton Equation (2), with constant volatility and they proved that Equation (2) admits as Lie symmetries the elements of Lie algebra, $\{A_{3,8} \oplus_s A_{3,1}\} \oplus_s \infty A_1$ (In the Mubarakzyanov Classification Scheme [10–13]). This means that Equation (2) is maximally symmetric and according to the Theorem of Sophus Lie [14] there exists a transformation on the space of variables $\{t, S, u\}$ in which Equation (2) can be written in the form of the heat equation. The last was an important result because the mathematical methods from physical science can be used for the study of differential equations in financial mathematics. A similar result has been found for the one-factor model of commodities [15], which means that the three different equations, the heat equation, the Black-Scholes-Merton equation and the one-factor model of commodities equation, are equivalent at the mathematical level even if they describe different subjects.

In recent years, Lie symmetries have covered a big range of applications in financial mathematics. For instance, the group invariants of the Cox-Ingersoll-Ross Pricing Equation have been studied in [16] and the nonlinear Merton model in [17]. As far as concerns the Asian option, a Lie symmetry classification has been performed in [18]. As for generalisations of the Black-Scholes-Merton Model, the Lie symmetries and the reduction process of the nonautonomous model can be found in [19,20], while another generalisation of Equation (2) with a “source” was studied in [21].

Furthermore, in [22,23], the symmetry analysis of the space- and time-dependent one-factor model of commodities and of the nonautonomous two-dimensional Black-Scholes-Merton Equations were performed. For other applications of Lie symmetries in financial mathematics, see, for instance, [24–26], and references therein.

The stochastic volatility model, Equation (5), is a $(1 + 2)$ evolution equation. Below, we perform a symmetry analysis and we determine the group invariant solutions. In particular, we restrict our analysis to the model in which the risk premium factor vanishes without necessarily $|\rho| = 1$ and from Equation (9), only the term which expresses the return-to-risk ratio survives. Moreover, we study two models for European options with stochastic volatility, the Heston model [27] and the Stein–Stein model [28]. The latter is a model without correlation between the two Brownian motions, W_t and \hat{Z}_t , *i.e.*, $\rho = 0$ in Equation (4). The plan of the paper is as follows.

In Section 2, we give the basic properties and definitions for the Lie point symmetries of differential equations and we perform the symmetry classification for our model. We find that Equation (5) without the risk premium factor is always invariant under the $\{3A_1\} \oplus_s \infty A_1$ Lie algebra. However, when $f(y)$ is constant, Equation (5) is invariant under a larger Lie algebra. The application of the Lie symmetries to Equation (5) can be found in Section 3, in which we reduce the $(1 + 2)$ evolution equation by using the zeroth-order invariants provided by the Lie symmetries and we derive invariant solutions. In Sections 4 and 5 we study two models of stochastic volatility for European options, the Heston model and the Stein-Stein model, respectively. For these two models, we find that both are invariant under the Lie algebra $\{3A_1\} \oplus_s \infty A_1$, and we apply the Lie symmetries to solve the equations of the two models. For the Heston model, the closed-form solution is expressed in terms of Kummer Functions, whereas for the Stein–Stein model, the closed-form solution is expressed in terms of Hypergeometric Functions. Furthermore, we give some numerical solutions for the two models. Finally, in Section 6, we discuss our results and draw our conclusions

2. Lie Symmetry Analysis

We consider the Black-Scholes-Merton Equation with stochastic volatility governed by the evolution Equation (5) for which the premium term depends only upon the return-to-risk ratio. For a time-independent rate-of-return, Equation (5) becomes

$$\begin{aligned} \mathcal{H} : 0 = & \frac{1}{2}f^2(y) S^2 u_{,SS} + \rho\beta S f(y) u_{,Sy} + \frac{1}{2}\beta^2 u_{,yy} \\ & + rSu_{,S} + \left[\alpha(m-y) - \beta\rho\frac{\mu-r}{f(y)} \right] u_{,y} - ru + u_{,t} \end{aligned} \tag{10}$$

Let Φ be the map of an one-parameter point transformation such as

$$\Phi(u(t, S, y)) = u'(t', S', y') \tag{11}$$

with infinitesimal transformation (ε is the parameter of smallness.)

$$t' = t + \varepsilon\zeta^1(t, S, y, u) \tag{12}$$

$$S' = S + \varepsilon\zeta^2(t, S, y, u) \tag{13}$$

$$y' = y + \varepsilon\zeta^3(t, S, y, u) \tag{14}$$

$$u' = u + \varepsilon\eta(t, S, y, u) \tag{15}$$

and generator

$$X = \frac{\partial t'}{\partial \varepsilon} \partial_t + \frac{\partial S'}{\partial \varepsilon} \partial_S + \frac{\partial y'}{\partial \varepsilon} \partial_y + \frac{\partial u'}{\partial \varepsilon} \partial_u \tag{16}$$

Consider now that $u(t, S, y)$ is a solution of Equation (10) and under the map Φ , Equation (11), $u'(t', S', y')$ is also a solution of Equation (10). Then, we say that the generator X , of the infinitesimal transformation of the one-parameter point transformation, Φ , is a Lie (point) symmetry of (Equation 10) and Equation (10) is invariant under the action of the map Φ . That means that there exists a function ψ such that the following condition holds [29]

$$X^{[2]}(\mathcal{H}) = \psi\mathcal{H}, \text{ mod } (\mathcal{H}) = 0 \tag{17}$$

or, equivalently,

$$X^{[2]}(\mathcal{H}) = 0 \tag{18}$$

where $X^{[2]}$ is the second prolongation/extension of X in the space of variables $\{t, S, y, u, u_{,S}, u_{,y}, u_{,SS}, u_{,Sy}, u_{,yy}\}$. Specifically $X^{[2]}$ is defined from the following formula

$$X^{[2]} = X + \eta_i^A \partial_{u_i^A} + \eta_{ij}^A \partial_{u_{ij}^A} \tag{19}$$

where η_i^A, η_{ij}^A are given by the relations

$$\eta_i^A = \eta_{,i}^A + u_{,i}^B \eta_{,B}^A - \zeta_{,i}^j u_{,j}^A - u_{,i}^A u_{,j}^B \zeta_{,B}^j \tag{20}$$

and

$$\begin{aligned} \eta_{ij}^A = & \eta_{,ij}^A + 2\eta_{,B(i}^A u_{,j)}^B - \zeta_{,ij}^k u_{,k}^A + \eta_{,BC}^A u_{,i}^B u_{,j}^C - 2\zeta_{,(i|B|}^k u_{,j)}^B u_{,k}^A \\ & - \zeta_{,BC}^k u_{,i}^B u_{,j}^C u_{,k}^A + \eta_{,B}^A u_{,ij}^B - 2\zeta_{,(j,i)k}^A u_{,k}^A - \zeta_{,B}^k \left(u_{,k}^A u_{,ij}^B + 2u_{,(j,i)k}^B u_{,k}^A \right) \end{aligned} \tag{21}$$

The importance of the existence of a Lie symmetry for a partial differential equation is that from the associated Lagrange’s system,

$$\frac{dx^i}{\zeta^i} = \frac{du^A}{\eta^A} \tag{22}$$

Zeroth-order invariants, $U^{[0]}(x^k, u^A)$, can be determined which can be used to reduce the number of the independent variables of the differential equation.

In the following, we perform a classification of the Lie symmetries of Equation (10). Function $f(y)$ is defined by the requirement that Equation (10) admit Lie symmetries. The latter requirement can be seen as a geometric selection rule as the Lie symmetries are generated from the elements of the Homothetic Algebra [30] of the (pseudo)Riemannian space, which defines the Laplace operator in the (1 + 2) evolution Equation (10). In our case, the (pseudo)Riemannian manifold is defined by the Brownian motions, W_t, \hat{Z}_t , of the stock price, S , and of the volatility, σ , respectively.

Before we proceed with the symmetry analysis, we remark that Equation (10) is a linear equation which means that it always admits the linear symmetry, $X_u = u\partial_u$ and the infinite-dimensional abelian subalgebra of solutions, $X_b = b(t, S, y)\partial_u$, where function $b(t, S, y)$, is a solution of the original Equation (10) [31].

2.1. Classification

From the symmetry condition Equation (17), we get a system of thirty-one equations (for the derivation of the system, we used the symbolic package SYM of Mathematica [32,33]) in which the solution of the system gives the form of the generator Equation (16) of the transformation Equation (11), that transforms solutions into solutions. From the latter system, we have the following results.

For arbitrary function, $f(y)$, Equation (10) admits the Lie symmetries

$$X_1 = \partial_t, X_2 = S\partial_S \tag{23}$$

plus the vector fields X_u, X_b . The algebra in which the Lie symmetries form is the $\{3A_1\} \oplus_s \infty A_1$.

When $f(y) = f_0$, Equation (10) admits the Lie symmetries

$$\bar{X}_1 = \partial_t, \bar{X}_2 = S\partial_S, \bar{X}_3 = e^{-\alpha t}\partial_y \tag{24}$$

$$\bar{X}_4 = 2f_0tS\partial_S + 2\frac{\beta}{\alpha}f_0\rho\partial_y + \left((f_0^2 - 2r)t + 2\ln S \right) u\partial_u \tag{25}$$

$$\bar{X}_5 = e^{\alpha t} \left(2f_0^2\beta\rho S\partial_S + \beta^2 f_0\partial_y \right) + (2\alpha f_0(y - m) + 2\beta(\mu - r)) u\partial_u \tag{26}$$

plus the vector fields X_u, X_b . The Lie Brackets of the Lie algebra are given in Table 1.

Table 1. Lie brackets of the Lie symmetries of Equation (10) for $f(y) = f_0$.

$[\bar{X}_I, \bar{X}_J]$	\bar{X}_1	\bar{X}_2	\bar{X}_3	\bar{X}_4	\bar{X}_5	X_u
\bar{X}_1	0	0	$-a\bar{X}_3$	$2f_0^2\bar{X}_2 + (f_0^2 - 2r) X_u$	$\alpha\bar{X}_5$	0
\bar{X}_2	0	0	0	$2X_u$	0	0
\bar{X}_3	$\alpha\bar{X}_3$	0	0	0	$2\alpha f_0 X_u$	0
\bar{X}_4	$-2f_0^2\bar{X}_2 - (f_0^2 - 2r) X_u$	$2X_u$	0	0	0	0
\bar{X}_5	$-a\bar{X}_5$	0	$-2\alpha f_0 X_u$	0	0	0
X_u	0	0	0	0	0	0

We remark that the two-factor model of commodities is invariant under the same algebra of point transformations [15,23]. That is an expected result because the two-factor model of commodities follows from the one-factor model in which the second factor, product, follows an Orstein-Uhlenbeck

process. Moreover, as we discussed in the Introduction, the one-factor model is maximally symmetric just like the Black-Scholes-Merton Equation.

On the other hand $f(y) = f_0$ means that the volatility σ is constant. However, the second Brownian motion, \hat{Z}_t , in the space wherein σ is defined, interacts with the Brownian motion W_t and modifies the Black-Scholes-Merton Model. However, in the case for which the correlation ρ vanishes, i.e., $\rho = 0$, Equation (10) is not reduced to Equation (2) but only when the Orstein-Uhlenbeck process is identically zero, that is, $\beta = 0, \alpha = 0$. Otherwise, the price u depends upon the Orstein-Uhlenbeck process.

We continue with the reduction of Equation (10) by applying the zeroth-order Lie invariants. Furthermore for every reduced equation we study the Lie symmetries.

3. Group Invariant Solutions

In this section, we apply the Lie symmetries in order to reduce Equation (10). We study the two cases, $f(y) = f_0$, and $f(y)$ to be an arbitrary function. In order to perform the reduction and the later equation to give a solution of the original problem, there should be a constraint between the Lie symmetry vector and the terminal condition. However, we perform the reduction without considering the terminal condition at the moment because the initial conditions can be modified from different options. As far as the invariant solutions of the Black-Scholes-Merton Equation (2) are concerned, see [34].

3.1. Arbitrary Function $f(y)$

For an arbitrary functional form of $f(y)$, as we saw above, Equation (10) admits three Lie point symmetries in addition to the infinite number of solution symmetries. The last cannot be used for the reduction. Hence, we do not consider them. Moreover, a solution in which u does not depend upon one of the independent variables is not an acceptable solution, that is, the static solution is of no interest. Therefore, we perform reductions with the symmetry vectors $Y_1 = X_1 + \kappa_1 X_u, Y_2 = X_1 + \kappa_2 X_u$ and $Y_{12} = X_1 + cX_2 + \kappa_3 X_u$.

Reduction with respect to the Lie invariants of the symmetry vector Y_1 gives

$$u(t, S, y) = \exp[\kappa_1 t] v(S, y) \tag{27}$$

where $v(S, y)$ satisfies the equation

$$0 = \frac{1}{2} f^2(y) S^2 v_{SS} + \rho \beta S f(y) v_{Sy} + \frac{1}{2} \beta^2 v_{yy} + r S v_{S} + \left[\alpha(m - y) - \beta \rho \frac{\mu - r}{f(y)} \right] v_{y} - (r - \kappa_1) v \tag{28}$$

For this equation, we have that except the linear symmetry and the infinite number of solution symmetries (we call them trivial symmetries) the equation admits the vector field $Y_2 = S \partial_S$, which is a reduced symmetry. Therefore, the application of Y_2 to Equation (28) gives the second-order ordinary differential equation

$$\beta^2 w_{yy} + \left[2\alpha(m - y) + \frac{2\beta}{f(y)} (\rho \kappa_2 f^2(y) - \mu + r) \right] w_y + [(\kappa_2^2 - 1) f^2(y) + 2(r\kappa_2 - r + \kappa_1)] w = 0 \tag{29}$$

where $w = w(y)$ and

$$u(t, S, y) = S^{\kappa_2} \exp[\kappa_1 t] w(y) \tag{30}$$

Equation (29) is a linear second-order differential equation, and it is well known that it is maximally symmetric and is invariant under the special linear (sl) algebra $sl(3, R)$ Lie algebra.

Similarly, if we perform a reduction with Y_2 , the reduced equation admits the Lie Symmetries X_1, X_v, X_b , and finally the solution is again given by Equation (30) with the constraint Equation (29)

Consider the application of the Lie symmetry vector Y_{12} to Equation (29). We have that

$$u(t, S, y) = \exp[k_3 t] v(z, y), \quad z = S \exp[-ct] \tag{31}$$

where

$$0 = z^2 f(y) v_{,zz} + 2\rho\beta f(y) v_{,zy} + \beta^2 v_{,yy} + 2(r - c) z v_{,z} + 2 \left[\alpha(m - y) - \beta\rho \frac{\mu - r}{f(y)} \right] v_{,y} - (r - \kappa_3) v \tag{32}$$

One can easily find that this equation only admits the Lie symmetry, $z\partial_z$, except the trivial symmetries, which is a reduced symmetry. Therefore, the application of the zeroth-order invariants of the symmetry vector ($z\partial_z + \kappa_4 v\partial_v$) in Equation (32) gives solution of the form Equation (30) with the constraint Equation (29)

We continue with the determination of the group invariant solutions for constant $f(y)$.

3.2. Constant Volatility

For $f(y) = f_0$, Equation (10) admits six Lie point symmetries, plus the infinite number of solution symmetries. Moreover, Equation (10) is an $(1 + 2)$ evolution equation, and, in order to reduce it to an ordinary differential equation, we have to apply the zeroth-order invariants of two Lie symmetries. From Table 1, we select reducing Equation (10) by using the following two-dimensional subalgebras $A_I = \{Y_1, Y_2\}$, $B_I = \{Y_2, \bar{X}_3 + \kappa X_u\}$, $B_{II} = \{Y_2, \bar{X}_5 + \kappa X_u\}$, $C_I = \{\bar{X}_3 + \kappa X_u, \bar{X}_4\}$ and $C_{II} = \{\bar{X}_4, \bar{X}_5\}$.

The reduction with the subalgebra A_I we studied in the previous subsection and the solution is Equation (30), where now from Equation (29) we have

$$w(y) = w_1 \mathbf{K} \left(-\frac{c_2}{4\alpha'} \frac{1}{2'} \frac{(\alpha(m - y) + \frac{c_1}{2})^2}{\alpha} \right) + w_2 \mathbf{\Lambda} \left(-\frac{c_2}{4\alpha'} \frac{1}{2'} \frac{(\alpha(m - y) + \frac{c_1}{2})^2}{\alpha} \right)$$

We have that w_1 and w_2 are constants,

$$c_1 = \frac{2\beta}{f_0} (\rho\kappa_2 f_0^2 - \mu + r), \quad c_2 = [(\kappa^2 - 1) f_0^2 + 2(r\kappa_2 - r + \kappa_1)]$$

and $\mathbf{K}, \mathbf{\Lambda}$ are Kummer Functions.

We continue with the application of the remaining subalgebras.

The application of B_I gives

$$u(t, S, y) = S^{\kappa_2} \exp(\kappa e^{\alpha t} y) \phi_I(t) \tag{33}$$

where $\phi_I(t)$ is given by the first-order ordinary differential equation

$$2f_0\phi_{I,t} + \left[(f_0^3\kappa_2 + 2f_0r)(\kappa_2 - 1) + 2\kappa\beta \left(f_0^2\rho\kappa_2 - \mu + r + \frac{\alpha\mu f_0}{\beta} \right) e^{\alpha t} + f_0\beta^2\kappa^2 e^{2\alpha t} \right] \phi_I = 0 \tag{34}$$

with solution

$$\ln \left(\frac{\phi_I(t)}{\phi_{I0}} \right) = -\frac{(f_0^2\kappa_2 + 2r)}{2} (\kappa_2 - 1) t - \frac{2\kappa\beta}{2\alpha f_0} \left(f_0^2\rho\kappa_2 - \mu + r + \frac{\alpha\mu f_0}{\beta} \right) e^{\alpha t} - \frac{\beta^2\kappa^2}{4\alpha} e^{2\alpha t} \tag{35}$$

From the subalgebra B_{II} , we find the solution

$$u(t, S, y) = \phi_{II}(t) S^{\kappa_2} \exp \left(\frac{\alpha}{\beta^2} y^2 + \left(\frac{\kappa e^{-\alpha t} - 2\alpha m}{\beta^2 f_0} - \frac{2(r - \mu + \rho\kappa_2 f_0^2)}{\beta f_0} \right) y \right) \tag{36}$$

where $\phi_{II}(t)$ is given by the expression

$$\ln\left(\frac{\phi_{II}(t)}{\phi_{II0}}\right) = -\frac{1}{2}\left[\left(f_0^2\kappa_2 + 2r\right)\left(\kappa_2 - 1\right) + 2\alpha\right]t + \left[-\frac{\kappa}{\alpha\beta^2f_0^2}\left(\beta\left(r - \mu + \rho\kappa_2f_0^2\right) + \alpha mf_0\right)e^{-\alpha t} + \frac{\kappa^2}{4\alpha f_0^2\beta^2}e^{-2\alpha t}\right] \tag{37}$$

For the subalgebra C_I , we have the invariant solution

$$u(t, S, y) = S^{\psi(t)} \exp(\kappa e^{\alpha t} y) \phi_{III}(t)$$

where

$$\psi(t) = -\frac{\kappa\beta\rho}{\alpha f_0 t} - \frac{r}{f_0^2} + \frac{1}{2} + \frac{2}{4f_0^2 t} \tag{38}$$

and $\phi_{III}(t)$ is given by the expression

$$\ln\left(\frac{\phi_{III}(t)}{\phi_{III0}}\right) = -\frac{1}{2}\ln t + \frac{(2\rho^2 - \alpha t)}{\alpha^2 t} \beta^2 \kappa^2 e^{2\alpha t} - \frac{\kappa}{2\alpha f_0} e^{\alpha t} \left(2\alpha mf + 2\beta\left(r - \mu - 2\rho r + \frac{\rho}{2}f_0^2\right) + \frac{(2r + f_0^2)^2}{8f_0^2}t\right) \tag{39}$$

Finally, from the subalgebra C_{II} , we find the invariant solution

$$u(t, S, y) = \frac{\phi_{IV}(t)}{\sqrt{\alpha t - 2\rho^2}} S^{\bar{\psi}(t,y)} \exp(V(t, y)) \tag{40}$$

where

$$\bar{\psi}(t, y) = \frac{4\rho f_0 a(m - y) + \beta(\alpha(f_0^2 - 2r)t - 4\rho(\mu - r)) + \alpha\beta^2}{2\beta f_0^2(\alpha t - 2\rho^2)} \tag{41}$$

$$V(t, y) = \left[\frac{\alpha^2}{\beta^2(\alpha t - 2\rho^2)} y - a \frac{2\beta\left(r - \mu + \frac{f_0^2}{2}\rho - 2r\rho\right) + 2f_0\alpha m}{\beta^2 f_0^2(\alpha t - 2\rho^2)} \right] ty \tag{42}$$

and function $\phi_{IV}(t)$ is given by the expression

$$\ln\left(\frac{\phi_{IV}(t)}{\phi_{IV0}}\right) = -\frac{f_0^2(8\alpha - 4r - f_0^2) - 4r^2}{8f_0^2(\alpha t - 2\rho^2)}(\alpha t^2 - 2\rho^2 t) + \frac{(f_0^2 - 2r)^2}{2f_0^2\alpha(\alpha t - 2\rho^2)}\rho^4 + \frac{2(f_0^2 - 2r^2)(f_0\alpha m - \beta(\mu - r))}{f_0^2\alpha\beta(\alpha t - 2\rho^2)}\rho^3 + \frac{(4\beta(r - \mu) + 2f_0\alpha m)}{\beta^2 f_0(\alpha t - 2\rho^2)}m\rho^2 + \frac{2(\mu - r)^2}{f_0^2\alpha(\alpha t - 2\rho^2)}\rho^2 \tag{43}$$

In the following section, we study a special model for stochastic volatility which has been proposed by Heston [27].

4. Heston Model

In the Heston model for stochastic volatility the stock price, S , and the volatility, $\sigma = f(y)$, satisfy the stochastic differential equation given below

$$dS_t = rS_t dt + S_t \sqrt{Y_t} dW_t \quad \text{and,} \tag{44}$$

$$dY_t = \theta(\bar{m} - Y_t) dt + \delta \sqrt{Y_t} d\hat{Z}_t \tag{45}$$

where, in comparison with Equations (1) and (3), we observe that $f(y) = \sqrt{y}$ and $\beta = \delta \sqrt{y}$. The differential equation which corresponds to that model is

$$0 = \frac{1}{2} Y S^2 u_{,SS} + \rho \delta Y S u_{,SY} + \frac{1}{2} \delta^2 Y u_{,YY} + r S u_{,S} + (\theta(\bar{m} - Y) - \lambda Y) u_{,Y} - ru + u_{,t} \tag{46}$$

Before we proceed with the symmetry analysis of Equation (46), we perform the coordinate transformation $Y = y^2$. Then, Equation (46) becomes

$$0 = \frac{1}{2} y^2 S^2 u_{,SS} + \beta \rho y S u_{,Sy} + \frac{1}{2} \beta^2 u_{,yy} + r S u_{,S} + \frac{1}{2} \left[c_1 y + \left(\frac{c_2}{y} \right) \right] u_{,y} - ru + u_{,t} \tag{47}$$

in which we have made the replacements $\frac{\delta}{2} \rightarrow \beta$, $c_1 = (\theta - \lambda)$ and $c_2 = \theta m - \beta^2$. Equation (47) can be compared with Equation (5) for $f(y) = y$. However, the risk premium factor of Equation (9) is not zero and has absorbed the term of the Orstein–Uhlenbeck process.

From the Lie symmetry condition, Equations (17) for (47) we find that this equation admits the Lie symmetries

$$X_1 = \partial_t, \quad X_2 = S \partial_S \tag{48}$$

plus the X_u, X_b , that is, Equation (47) is invariant under the Lie algebra $\{3A_1\} \oplus_s \infty A_1$.

We continue with the application of the Lie symmetries in order to reduce Equation (48). We follow the results of Section 3.1, that is, we apply the group invariants of the subalgebra $\{Y_1, Y_2\}$.

We find that the corresponding invariant solution of Equation (47) is

$$u(t, S, y) = S^{\kappa_2} e^{\kappa_1 t} W(y) \tag{49}$$

where $W(y)$ satisfies the linear second-order differential equation:

$$\beta^2 W_{,yy} + \left(2\kappa_2 \beta \rho + c_1 y + \frac{c_2}{y} \right) W_{,y} + \left(2\kappa_1 - 2r(1 - \kappa_2) + y^2 (\kappa_2^2 - \kappa_1^2) \right) W = 0 \tag{50}$$

the solution in closed form of which is expressed in terms of Kummer Functions.

In Figures 1 and 2, we give numerical solutions of Equation (50). Figure 1 is for negative value of $\Delta\kappa$, whereas Figure 2 is for positive value of $\Delta\kappa$, where $\Delta\kappa = \kappa_2 - \kappa_1$.




Figure 1. Evolution of the solution Equation (50) of the Heston model. For the numerical solutions, we select $\rho = 0.5$, $\beta = |0.7|$, $r = 0.5$, $\kappa_1 = 1$ and $\kappa_2 = 0.5$. The left figures are for negative c_1 , c_2 , while the right figures are for negative c_1 and positive c_2 . The solid lines are for $|c_2| = 5|c_1|$, the dotted lines are for $|c_2| = 0.2|c_1|$, and the dash-dash lines are for $|c_2| = |c_1|$. The top figures are for $\beta > 0$, while the lower figures for $\beta < 0$.




Figure 2. Evolution of the solution Equation (50) of the Heston model. For the numerical solutions, we select $\rho = 0.5$, $\beta = |0.7|$, $r = 0.5$, $\kappa_1 = 1$ and $\kappa_2 = 1.01$. The left figures are for negative c_1 , c_2 , while the right figures are for negative c_1 and positive c_2 . The solid lines are for $|c_2| = \frac{3}{2}|c_1|$, the dotted lines are for $|c_2| = \frac{2}{3}|c_1|$, and the dash-dash lines are for $|c_2| = |c_1|$. The top figures are for $\beta > 0$, while the lower figures for $\beta < 0$.

5. Stein-Stein Model

The model which has been proposed by Elias M. Stein and Jeremy C. Stein [28] describes an European option with stochastic volatility for which the correlation among the two Brownian motions vanishes, *i.e.*, $\rho = 0$, in Equation (4). Moreover, they considered that the risk premium factor is constant, *i.e.*, $\gamma(t, S, y) = \gamma_0$ and the volatility is $f(y) = y$, while the stochastic differential equation is

$$dS_t = rS_t dt + S_t Y_t dW_t \quad \text{and} \quad (51)$$

$$dY_t = \theta(\bar{m} - Y_t) dt + \delta d\hat{Z}_t \quad (52)$$

Therefore, from Equation (5), we have that the (1 + 2) evolution differential equation of the Stein-Stein model is

$$\frac{1}{2}y^2S^2u_{,SS} + \frac{1}{2}\beta^2u_{,yy} + rSu_{,S} + [\alpha(m - y) - \beta\gamma_0]v_{,y} - ru + u_{,t} = 0 \tag{53}$$

From the Lie symmetry condition Equation (17), we find that Equation (53) admits the Lie symmetries

$$X_1 = \partial_t, X_2 = S\partial_S, X_u = u\partial_u, X_b = b(t, S, y)\partial_u \tag{54}$$

which form the Lie algebra $\{3A_1\} \oplus_s \infty A_1$, and it is the admitted algebra of the Heston model and of Equation (10) for the arbitrary function $f(y)$.

Following the steps of the previous sections, we find that the invariant solution of the Stein-Stein model with respect to the Lie algebra $\{Y_1, Y_2\}$ is

$$u(t, S, y) = S^{\kappa_2}e^{\kappa_1 t}Y(y) \tag{55}$$

where $Y(y)$ is given by the linear second-order differential equation

$$\beta^2Y_{,yy} + (2\omega - 2ay)Y_{,y} + \left(2r(\kappa_2 - 1) + 2\kappa_1 + y^2(\kappa_2^2 - \kappa_1^2)\right)Y = 0 \tag{56}$$

The closed-form solution of this equation can be expressed in terms of the Hypergeometric Functions, where $\omega = \alpha m - \beta\gamma_0$. In Figures 3 and 4, we give the numerical evolution of $Y(y)$ for various values of the parameters, ω and α , for negative $\Delta\kappa$ and positive $\Delta\kappa$, respectively, where $\Delta\kappa = \kappa_2 - \kappa_1$.




Figure 3. Evolution of the solution Equation (56) of the Stein-Stein model. For the numerical solutions, we select $\beta = 0.5, r = 0.5, \kappa_1 = 1$ and $\kappa_2 = 0.5$. The figures are for $\omega = 0, \omega = 0.5, \omega = -0.5$ and $\omega = \alpha$, respectively. The solid lines are for $\alpha = 0.1$, the dotted lines are for $\alpha = 0.3$ and the dash-dash lines are for $\alpha = 0.4$.




Figure 4. Evolution of the solution Equation (56) of the Stein–Stein model. For the numerical solutions, we select $\beta = 0.5$, $r = 0.5$, $\kappa_1 = 1$ and $\kappa_2 = 1.01$. The figures are for $\omega = 0$, $\omega = 0.5$, $\omega = -0.5$ and $\omega = \alpha$, respectively. The solid lines are for $\alpha = 1.1$, the dotted lines are for $\alpha = 1.3$ and the dash-dash lines are for $\alpha = 1.4$.

6. Conclusions

Volatility with a stochastic process has been shown to be essential for Financial Mathematics. In this work, we studied the algebraic properties, *i.e.*, the Lie symmetries, of the modified Black-Scholes-Merton Equation for European options with a stochastic volatility. We have shown that the autonomous model without the risk premium factor is invariant under a group of point transformations which form the $\{3A_1\} \oplus_s \infty A_1$ Lie Algebra for an arbitrary functional form of the volatility, σ . Moreover, when the volatility is constant but the price of the option depends on the second Brownian motion, in which the volatility is defined, the modified Black-Scholes-Merton Model is invariant under six, plus the infinity, Lie point symmetries and it is not maximally symmetric as the Black-Scholes-Merton Equation with nonstochastic volatility is.

Furthermore, we showed that the Black-Scholes-Merton Model, in which the volatility is constant, but is defined by an Orstein-Uhlenbeck process, is invariant under the same group of point transformations as that of the two-factor model of commodities. The reason for that is that the two models have in common the terms which follow from the Orstein-Uhlenbeck process.

Moreover, we applied the zeroth-order invariants of the Lie symmetries, and we reduced the model to a linear second-order differential equation. As far as the case of constant volatility is concerned, we found the closed forms of the group invariant solutions.

Finally, we studied the algebraic properties and the invariant solutions of two models, the Heston model and the Stein-Stein model, with stochastic volatility of special interest. For each model, we found the invariant solution and we gave some figures for the evolution of the models. Of course because Equation (5) is a linear equation, the general solution is given by the linear combination of the invariant solutions that we have found, while the latter are constrained by the initial conditions and the boundary conditions of the model.

A general consideration of Equation (5), in which the risk premium factor plays a role is still in progress, and the results will be published elsewhere.

Acknowledgments: The research of Andronikos Paliathanasis was supported by FONDECYT grant No. 3160121. K. Krishnakumar thanks the University Grants Commission for providing a University Grants Commission-Basic Scientific Research Fellowship to perform this research work.

Author Contributions: Andronikos Paliathanasis and Peter G.L. Leach determined the problem and the method for the solution; K. Krishnakumar and K.M. Tamizhmani did the calculations. Andronikos Paliathanasis and Peter G.L. Leach wrote the paper and they gave interpretation to the results.

Conflicts of Interest: The authors declare no conflict of interest.

References

- Black, F.; Scholes, M. The valuation of option contracts and a test of market efficiency. *J. Financ.* **1972**, *27*, 399–417.
- Black, F.; Scholes, M. The pricing of options and corporate liabilities. *J. Polit. Econ.* **1973**, *81*, 637–659.
- Merton, R.C. On the pricing of corporate data: The risk structure of interest rates. *J. Financ.* **1974**, *29*, 449–470.
- Sophocleous, C.; O'Hara, J.G.; Leach, P.G.L. Algebraic solution of the Stein-Stein Model for stochastic volatility. *Commun. Nonlinear Sci. Numer. Simul.* **2010**, *16*, 1752–1759.
- Achdou, Y.; Pironneau, O. *Computational Methods for Option Pricing*; SIAM USA Philadelphia: Seattle, WA, USA, 2005.
- Fouque, J.P.; Papanicolaou, G.; Sircar, K.R. *Derivatives in Financial Markets with Stochastic Volatility*; Cambridge University Press: Cambridge, UK, 2000.
- Hull, J.; White, A. The pricing of options on assets with stochastic volatilities. *J. Financ.* **1987**, *42*, 281–300.
- Andersen, T.G.; Benzoni, L.; Lund, J. An empirical investigation of continuous-time equity return models. *J. Financ.* **2002**, *57*, 1239–1284.
- Gazizov, R.K.; Ibragimov, N.H. Lie symmetry analysis of differential equations in Finance. *Nonlinear Dyn.* **1997**, *17*, 387–407.
- Morozov, V.V. Classification of six-dimensional nilpotent Lie algebras. *Izv. Vyss. Uchebn Zavendenii Mat.* **1958**, *5*, 161–171.
- Mubarakzyanov, G.M. On solvable Lie algebras. *Izv. Vyss. Uchebn Zavendenii Mat.* **1963**, *32*, 114–123.
- Mubarakzyanov, G.M. Classification of real structures of five-dimensional Lie algebras. *Izv. Vyss. Uchebn Zavendenii Mat.* **1963**, *34*, 99–106.
- Mubarakzyanov, G.M. Classification of solvable six-dimensional Lie algebras with one nilpotent base element. *Izv. Vyss. Uchebn Zavendenii Mat.* **1963**, *35*, 104–116.
- Lie, S. *Lectures on Differential Equations with Known Infinitesimal Transformations*; Bibliotheca Teubneriana: Leipzig, Germany, 1891.
- Sophocleous, C.; Leach, P.G.L.; Andriopoulos, K. Algebraic properties of evolution partial differential equations modelling prices of commodities. *Math. Methods Appl. Sci.* **2008**, *31*, 679–694.
- Sinkala, W.; Leach, P.G.L.; O'Hara, J.G. An optimal system and group-invariant solutions of the Cox-Ingersoll-Ross pricing equation. *Appl. Math. Comput.* **2008**, *201*, 95–107.
- Naicker, V.; O'Hara, J.G.; Leach, P.G.L. A note on the integrability of the classical portfolio selection model. *Appl. Math. Lett.* **2010**, *23*, 1114–1119.
- Caister, N.C.; O'Hara, J.G.; Govinder, K.S. Solving the Asian option PDE using Lie symmetry methods. *Int. J. Theor. Appl. Financ.* **2010**, *13*, 1256–1277.
- Sophocleous, C.; Leach, P.G.L. Algebraic aspects of evolution partial differential equations arising in financial mathematics. *Appl. Math. Inf. Sci.* **2010**, *4*, 289–305.
- Tamizhmani, K.M.; Krishnakumar, K.; Leach, P.G.L. Algebraic resolution of equations of the Black-Scholes type with arbitrary time-dependent parameters. *Appl. Math. Comput.* **2014**, *247*, 115–124.
- Bozhkov, Y.; Dimas, S. Group classification of a generalized Black-Scholes-Merton equation. *Commun. Nonlinear Sci. Numer. Simul.* **2014**, *19*, 2200–2211.
- Paliathanasis, A.; Morris, R.M.; Leach, P.G.L. The algebraic properties of the space- and time-dependent one-factor problem of commodities. *Quaest. Math.* **2016**, submitted.
- Paliathanasis, A.; Morris, R.M.; Leach, P.G.L. Lie symmetries of $(1 + 2)$ nonautonomous evolution equations in Financial Mathematics. *Mathematics* **2016**, submitted.

24. Bozhkov, Y.; Dimas, S. Group classification of a generalization of the Heath equation. *Appl. Math. Comput.* **2014**, *243*, 121–131.
25. Okelola, M.O.; Govinder, K.S.; O'Hara, J.G. Solving a partial differential equation associated with the pricing of power options with time-dependent parameters. *Math. Methods Appl. Sci.* **2015**, *38*, 2901–2910.
26. Motsepa, T.; Khalique, C.M.; Molati, M. Group classification of a general bond-option pricing equation of mathematical finance. *Abstr. Appl. Anal.* **2014**, *2014*, 709871.
27. Heston, S. A closed form solution for options with stochastic volatility with application to bond and currency options. *Rev. Financ. Stud.* **1993**, *6*, 327–343.
28. Stein, E.M.; Stein, J.C. Stock-price distributions with stochastic volatility: An analytic approach. *Rev. Financ. Stud.* **1991**, *4*, 727–752.
29. Bluman, G.W.; Kumei, S. *Symmetries of Differential Equations*; Springer-Verlag: New York, NY, USA, 1989.
30. Paliathanasis, A.; Tsamparlis, M. Lie point symmetries of a general class of PDEs: The heat equation. *J. Geom. Phys.* **2012**, *62*, 2443–2456.
31. Bluman, G.W. Simplifying the form of Lie groups admitted by a given differential equation. *J. Math. Anal. Appl.* **1990**, *145*, 52–62.
32. Dimas, S.; Tsoubelis, D. SYM: A new symmetry-finding package for mathematica. In *Group Analysis of Differential Equations*; Ibragimov, N.H., Sophocleous, C., Damianou, P.A., Eds.; University of Cyprus: Nicosia, Cyprus, 2005; pp. 64–70.
33. Dimas, S.; Tsoubelis, D. A new mathematica-based program for solving overdetermined systems of PDEs. In *Proceedings of the 8th International Mathematica Symposium, Avignon, France, 19–23 June 2006*.
34. Pooe, C.A.; Mahomed, F.M.; Soh, C.W. Invariant solutions of the Black-Scholes equation. *Math. Comput. Appl.* **2003**, *8*, 63–70.



© 2016 by the authors; licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC-BY) license (<http://creativecommons.org/licenses/by/4.0/>).