

Article

A Numerical Solution of Fractional Lienard's Equation by Using the Residual Power Series Method

Muhammed I. Syam

Department of Mathematical Sciences, United Arab Emirates University, Al-Ain 15551, United Arab Emirates; m.syam@uaeu.ac.ae

Received: 1 October 2017; Accepted: 21 November 2017; Published: 22 December 2017

Abstract: In this paper, we investigate a numerical solution of Lienard's equation. The residual power series (RPS) method is implemented to find an approximate solution to this problem. The proposed method is a combination of the fractional Taylor series and the residual functions. Numerical and theoretical results are presented.

Keywords: Lienard's equation; Caputo derivative; Taylor series; residual power series (RPS)

1. Introduction

The ordinary Lienard's equation is given by:

$$y''(x) + f(y)y'(x) + g(y) = r(x). \quad (1)$$

Different choices of f , g , and r will produce different models. For example, if $f(y)y'(x)$ is the damping force, $g(y)$ is the restoring force, and $r(x)$ is the external force, we get the damped pendulum equation. However, if we choose $f(y) = \epsilon(y^2 - 1)$, $g(y) = y$, and $r(x) = 0$, we get a nonlinear model of electronic oscillation, see [1,2].

Several researchers have studied the exact solution of special cases of Equation (1). For example, Feng [3] investigated the exact solution of:

$$y''(x) + a y(x) + b y^3(x) + c y^5(x) = 0. \quad (2)$$

He found that one of the solutions to Equation (2) is given by:

$$y(x) = \sqrt{-\frac{2a}{b}(1 + \tan(\sqrt{-ax}))} \quad (3)$$

when $\frac{b^2}{4} - 4\frac{ac}{3} = 0$, $b > 0$, and $a < 0$. Several methods are used to investigate the solution of nonlinear equation such as the homotopy analysis method (HAM), Adomian decomposition method, and variational method [4–12]. Equation (2) was studied by many researchers, notably Kong [13], Matinfar et al. [14,15], and others. In this paper, we generalize Equation (2) to the fractional case. It is not an easy task to solve highly nonlinear differential equations of fractional order. Many of the researchers tried to solve nonlinear equations by using different techniques. For example, Liao studied an analytical method termed as the HAM [9–11] to examine nonlinear problems. Furthermore, the HAM is used by many researchers to solve various types of nonlinear problems such as fractional Black–Scholes equation [12], natural convective heat and mass transfer in a steady 2-D MHD fluid flow over a stretching vertical surface via porous media [16], micropolar flow in a porous channel in the presence of mass injection [17], etc. The standard classical analytic schemes require more computational time and computer memory. Some researchers combine

analytical techniques with Laplace transform to study nonlinear problems such as a class of nonlinear differential equations [18], the nonlinear boundary value problem on a semi infinite domain [19], the fractional convection–diffusion equation [20], the fractional Keller–Segel model [21], fractional coupled Burgers equations [16], and the synthesis of FO-PID controllers [22]. In this paper, we consider the following class of fractional Lienard’s equation of the form:

$$D^{2\alpha}y(x) + a y(x) + b y^3(x) + c y^5(x) = 0, \quad x > 0, \quad \frac{1}{2} < \alpha \leq 1, \tag{4}$$

subject to:

$$y(0) = y_0, \quad D^\alpha y(0) = y_1 \tag{5}$$

where $a, b, c, y_0,$ and y_1 are constants. The derivative in Equation (4) is in the Caputo sense. The Caputo derivative is defined as follows; see [23,24].

Definition 1. Let n be the smallest integer greater than or equal to α . The Caputo fractional derivative of order $\alpha > 0$ is defined as:

$$D^\alpha y(x) = \begin{cases} \frac{1}{\Gamma(n-\alpha)} \int_0^x (x-t)^{n-\alpha-1} y^{(n)}(t) dt, & n-1 < \alpha < n, \\ y^{(n)}(x), & \alpha = n \in \mathbb{N}. \end{cases}$$

The power rule of the Caputo derivative is given as follows.

Theorem 1. The Caputo fractional derivative of the power function is given by:

$$D^\alpha x^p = \begin{cases} \frac{\Gamma(p+1)}{\Gamma(p-\alpha+1)} x^{p-\alpha}, & n-1 < \alpha < n, \quad p > n-1, \quad p \in \mathbb{R}, \\ 0, & n-1 < \alpha < n, \quad p \leq n-1, \quad p \in \mathbb{N}. \end{cases}$$

We implement the residual power series (RPS) method [13,25–27] to solve Equation (4). We start by the following definition and some theorems related to the RPS, [13,27].

Definition 2. A power series expansion of the form:

$$\sum_{m=0}^{\infty} c_m (x - x_0)^{m\alpha} = c_0 + c_1 (x - x_0)^\alpha + c_2 (x - x_0)^{2\alpha} + \dots$$

where $0 \leq n - 1 < \alpha \leq n, x \leq x_0,$ is called fractional power series FPS about $x = x_0$.

Theorem 2. Suppose that f has a RPS representation at $x = x_0$ of the form:

$$f(x) = \sum_{m=0}^{\infty} c_m (x - x_0)^{m\alpha}, \quad x_0 \leq x < x_0 + R,$$

where R is the radius of convergence. If $D^{m\alpha} f(x), m = 0, 1, 2, \dots$ are continuous on $(x_0, x_0 + R),$ then $c_m = \frac{D^{m\alpha} f(x_0)}{\Gamma(1+m\alpha)}.$

2. The Residual Power Series Method for Fractional Lienard’s Equation

Write the solution of Equation (4) as a fractional power series of the form:

$$y(x) = \sum_{n=0}^{\infty} y_n \frac{x^{n\alpha}}{\Gamma(1+n\alpha)}. \tag{6}$$

Using the initial conditions in Equation (5), we approximate $y(x)$ in Equation (6) by:

$$y_k(x) = y_0 + y_1 \frac{x^\alpha}{\Gamma(1+\alpha)} + \sum_{n=2}^k y_n \frac{x^{n\alpha}}{\Gamma(1+n\alpha)}, \quad k = 2, 3, \dots \tag{7}$$

where $y_1(x) = y_0 + y_1 \frac{x^\alpha}{\Gamma(1+\alpha)}$ is considered as the first RPS approximate solution of $y(x)$. To find the values of the RPS-coefficients $y_n, n = 2, 3, 4, \dots$, we solve the equation:

$$D^{(n-2)\alpha} Res_n(0) = 0, \quad n = 2, 3, 4, \dots \tag{8}$$

where $Res_k(x)$ is the k th residual function [13,27] and it is defined by:

$$Res_k(x) = D^{2\alpha} y_k(x) + a y_k(x) + b y_k^3(x) + c y_k^5(x). \tag{9}$$

To find y_2 in Equation (7), we substitute the second RPS approximate solution $y_2(x) = y_0 + y_1 \frac{x^\alpha}{\Gamma(1+\alpha)} + y_2 \frac{x^{2\alpha}}{\Gamma(1+2\alpha)}$ into:

$$\begin{aligned} Res_2(x) &= D^{2\alpha} y_2(x) + a y_2(x) + b y_2^3(x) + c y_2^5(x) \\ &= y_2 + a \left(y_0 + y_1 \frac{x^\alpha}{\Gamma(1+\alpha)} + y_2 \frac{x^{2\alpha}}{\Gamma(1+2\alpha)} \right) \\ &\quad + b \left(y_0 + y_1 \frac{x^\alpha}{\Gamma(1+\alpha)} + y_2 \frac{x^{2\alpha}}{\Gamma(1+2\alpha)} \right)^3 \\ &\quad + c \left(y_0 + y_1 \frac{x^\alpha}{\Gamma(1+\alpha)} + y_2 \frac{x^{2\alpha}}{\Gamma(1+2\alpha)} \right)^5. \end{aligned} \tag{10}$$

Then, we solve $Res_2(0) = 0$ to get:

$$y_2 = - \left(a y_0 + b y_0^3 + c y_0^5 \right). \tag{11}$$

To find y_3 in Equation (7), we substitute the third RPS approximate solution $y_3(x) = y_0 + y_1 \frac{x^\alpha}{\Gamma(1+\alpha)} + y_2 \frac{x^{2\alpha}}{\Gamma(1+2\alpha)} + y_3 \frac{x^{3\alpha}}{\Gamma(1+3\alpha)}$ into:

$$\begin{aligned} Res_3(x) &= D^{2\alpha} y_3(x) + a y_3(x) + b y_3^3(x) + c y_3^5(x) \\ &= y_3 + y_3 \frac{x^\alpha}{\Gamma(1+\alpha)} + a \left(y_0 + y_1 \frac{x^\alpha}{\Gamma(1+\alpha)} + y_2 \frac{x^{2\alpha}}{\Gamma(1+2\alpha)} + y_3 \frac{x^{3\alpha}}{\Gamma(1+3\alpha)} \right) \\ &\quad + b \left(y_0 + y_1 \frac{x^\alpha}{\Gamma(1+\alpha)} + y_2 \frac{x^{2\alpha}}{\Gamma(1+2\alpha)} + y_3 \frac{x^{3\alpha}}{\Gamma(1+3\alpha)} \right)^3 \\ &\quad + c \left(y_0 + y_1 \frac{x^\alpha}{\Gamma(1+\alpha)} + y_2 \frac{x^{2\alpha}}{\Gamma(1+2\alpha)} + y_3 \frac{x^{3\alpha}}{\Gamma(1+3\alpha)} \right)^5. \end{aligned} \tag{12}$$

Then, we solve $D^\alpha Res_3(0) = 0$ to get:

$$y_3 = - \left(a y_1 + 3b y_0^2 y_1 + 5c y_0^4 y_1 \right). \tag{13}$$

In general, to find $y_k(x)$, we substitute the k^{th} RPS approximate solution $y_k(x)$ into:

$$Res_k(x) = D^{2\alpha} y_k(x) + a y_k(x) + b y_k^3(x) + c y_k^5(x). \tag{14}$$

Then, we solve $D_t^{(k-2)\alpha} Res_k(0) = 0$ to get:

$$y_k = - \left(\begin{aligned} &a y_{k-2} + b \sum_{i=0}^{k-2} \sum_{j=0}^{k-2-i} \frac{y_i y_j y_{k-2-i-j} \Gamma(1+(k-2)\alpha)}{\Gamma(1+i\alpha)\Gamma(1+j\alpha)\Gamma(1+(k-2-i-j)\alpha)} \\ &+ c \sum_{i=0}^{k-2} \sum_{j=0}^{k-2-i} \sum_{l=0}^{k-2-i-j} \sum_{m=0}^{k-2-i-j-l} \frac{y_i y_j y_l y_m y_{k-2-i-j-l-m} \Gamma(1+(k-2)\alpha)}{\Gamma(1+i\alpha)\Gamma(1+j\alpha)\Gamma(1+l\alpha)\Gamma(1+m\alpha)\Gamma(1+(k-2-i-j-l-m)\alpha)} \end{aligned} \right). \tag{15}$$

3. Convergence Analysis

In this section, we prove the convergence of the proposed method. We start by the following lemma.

Lemma 1. *The classical power series $\sum_{n=0}^{\infty} y_n x^n$, $-\infty < x < \infty$, has a radius of convergence R if and only if the fractional power series $\sum_{n=0}^{\infty} y_n x^{\alpha n}$, $x \geq 0$, has a radius of convergence $R^{\frac{1}{\alpha}}$.*

Proof. See [28]. \square

Theorem 3. *The fractional power series:*

$$y(x) = \sum_{n=0}^{\infty} y_n \frac{x^{n\alpha}}{\Gamma(1 + n\alpha)} \tag{16}$$

where the coefficients are defined in Equation (15) has a positive radius of convergence.

Proof. From Equation (15), one can see that:

$$\frac{|y_k|}{\Gamma(1 + k\alpha)} \leq \left(\begin{array}{c} A y_{k-2} + B \sum_{i=0}^{k-2} \sum_{j=0}^{k-2-i} |y_i| |y_j| |y_{k-2-i-j}| \\ + C \sum_{i=0}^{k-2} \sum_{j=0}^{k-2-i} \sum_{l=0}^{k-2-i-j} \sum_{m=0}^{k-2-i-j-l} |y_i| |y_j| |y_l| |y_m| |y_{k-2-i-j-l-m}| \end{array} \right) \tag{17}$$

where:

$$\begin{aligned} A &= \frac{|a|}{\Gamma(1 + k\alpha)}, \\ B &= \underset{\substack{0 \leq j \leq k-2-i \\ 0 \leq i \leq k-2}}{\text{Max}} \left\{ \frac{\Gamma(1 + (k-2)\alpha)}{\Gamma(1 + i\alpha)\Gamma(1 + j\alpha)\Gamma(1 + (k-2-i-j)\alpha)\Gamma(1 + k\alpha)} \right\} |b|, \\ C &= \underset{\substack{0 \leq m \leq k-2-i-j-l \\ 0 \leq l \leq k-2-i-j \\ 0 \leq j \leq k-2-i \\ 0 \leq i \leq k-2}}{\text{Max}} \left\{ \frac{\Gamma(1 + (k-2)\alpha)}{\rho_{i,j,k,l,m}} \right\} |c|, \end{aligned}$$

$$\rho_{i,j,k,l,m} = \Gamma(1 + i\alpha)\Gamma(1 + j\alpha)\Gamma(1 + l\alpha)\Gamma(1 + m\alpha)\Gamma(1 + (k-2-i-j-l-m)\alpha)\Gamma(1 + k\alpha).$$

Let:

$$f(x) = \sum_{k=0}^{\infty} a_k x^k \tag{18}$$

where $a_0 = |y_0|$, $a_1 = \frac{|y_1|}{\Gamma(1+\alpha)}$, and

$$\begin{aligned} a_k &= A a_{k-2} + B \sum_{i=0}^{k-2} \sum_{j=0}^{k-2-i} a_i a_j a_{k-2-i-j} \\ &+ C \sum_{i=0}^{k-2} \sum_{j=0}^{k-2-i} \sum_{l=0}^{k-2-i-j} \sum_{m=0}^{k-2-i-j-l} a_i a_j a_l a_m a_{k-2-i-j-l-m} \end{aligned} \tag{19}$$

for $k = 2, 3, \dots$. Then,

$$\begin{aligned} \omega &= f(x) = a_0 + a_1 x + x^2 \sum_{k=0}^{\infty} a_{k+2} x^k \\ &= a_0 + a_1 x + x^2 \left(\begin{aligned} &A \sum_{k=0}^{\infty} a_k x^k + B \sum_{k=0}^{\infty} \left(\sum_{i=0}^k \sum_{j=0}^{k-i} a_i a_j a_{k-i-j} \right) x^k \\ &+ C \sum_{k=0}^{\infty} \left(\sum_{i=0}^k \sum_{j=0}^{k-i} \sum_{l=0}^{k-i-j} \sum_{m=0}^{k-i-j-l} a_i a_j a_l a_m a_{k-i-j-l-m} \right) x^k \end{aligned} \right). \end{aligned} \tag{20}$$

Let:

$$H(x, \omega) = \omega - a_0 - a_1 x - x^2 (A\omega + B\omega^3 + C\omega^5). \tag{21}$$

It is easy to see that $H(x, \omega)$ is an analytic function in the (x, ω) - plane and

$$H(0, a_0) = 0, H_{\omega}(0, a_0) = 1 \neq 0. \tag{22}$$

By implicit function theorem [29], $f(x)$ is analytic function in a neighborhood of the point $(0, a_0)$ of the (x, ω) - plane with a positive radius of convergence. Thus, the series in Equation (6) is convergent by Lemma 1. \square

4. Numerical Results

In this section, we present four examples to show the efficiency of the proposed approach. Comparison with the exact solution presented in Equation (3) is reported in Tables 1–4 for different choices of $a, b,$ and c with $\alpha = 1$. Let:

$$error(x) = |y_{exact}(x) - y_5(x)|, x \geq 0. \tag{23}$$

In these two examples, we use $k = 3$. Then,

$$\begin{aligned} y_2 &= - (a y_0 + b y_0^3 + c y_0^5), \\ y_3 &= - (a y_1 + 3b y_0^2 y_1 + 5c y_0^4 y_1). \end{aligned} \tag{24}$$

Example 1. Consider the following class of fractional differential equation:

$$D^{2\alpha} y(x) - y(x) + 3y^3(x) - \frac{27}{16} y^5(x) = 0, \frac{1}{2} < \alpha \leq 1, \tag{25}$$

subject to:

$$y(0) = y_0 = \sqrt{\frac{2}{3}}, D^{\alpha} y(0) = y_1 = \frac{1}{\sqrt{6}}. \tag{26}$$

Then, the error, when $\alpha = 1$, is reported in Table 1. Figure 1 shows the effect of α on the solution for $\alpha = 0.6, 0.7, 0.8, 0.9, 1$.

Table 1. Error when $\alpha = 1$.

x	Exact Solution	$y_3(x)$	$error(x)$
0.00	0.816497	0.816497	0
0.02	0.824622	0.824620	2.2×10^{-6}
0.04	0.832675	0.832658	1.7×10^{-5}
0.06	0.840663	0.840606	5.8×10^{-5}
0.08	0.848595	0.848460	1.4×10^{-4}
0.10	0.856479	0.856216	2.6×10^{-4}




Figure 1. The approximate solution for $\alpha = 0.6, 0.7, 0.8, 0.9, 1$.

Example 2. Consider the following class of fractional differential equation:

$$D^{2\alpha}y(x) - y(x) + 4y^3(x) - 3y^5(x) = 0, \quad \frac{1}{2} < \alpha \leq 1, \tag{27}$$

subject to:

$$y(0) = y_0 = \sqrt{\frac{1}{2}}, \quad D^\alpha y(0) = y_1 = \frac{1}{\sqrt{8}}. \tag{28}$$

Then, the error, when $\alpha = 1$, is reported in Table 2. Figure 2 shows the effect of α on the solution for $\alpha = 0.6, 0.7, 0.8, 0.9, 1$.

Table 2. Error when $\alpha = 1$.

x	Exact Solution	$y_3(x)$	$error(x)$
0.00	0.707107	0.707107	0
0.02	0.714142	0.714144	1.9×10^{-6}
0.04	0.721103	0.721118	1.5×10^{-5}
0.06	0.727986	0.728036	4.9×10^{-5}
0.08	0.734789	0.734905	1.1×10^{-4}
0.10	0.741508	0.741733	2.2×10^{-4}




Figure 2. The approximate solution for $\alpha = 0.6, 0.7, 0.8, 0.9, 1$.

Example 3. Consider the following class of fractional differential equation:

$$D^{2\alpha}y(x) - y(x) + 2y^3(x) - \frac{3}{4}y^5(x) = 0, \quad \frac{1}{2} < \alpha \leq 1, \tag{29}$$

subject to:

$$y(0) = y_0 = 1, D^\alpha y(0) = y_1 = 1. \tag{30}$$

Then, the error, when $\alpha = 1$, is reported in Table 3. Figure 3 shows the effect of α on the solution for $\alpha = 0.6, 0.7, 0.8, 0.9, 1$.

Table 3. Error when $\alpha = 1$.

x	Exact Solution	$y_3(x)$	$error(x)$
0.00	1.41421356	1.41421356	0
0.02	1.43407937	1.43407951	1.4×10^{-8}
0.04	1.45370489	1.45370501	1.2×10^{-6}
0.06	1.47313071	1.47313091	2.0×10^{-6}
0.08	1.49239558	1.49239579	2.1×10^{-6}
0.10	1.51153681	1.51153707	2.6×10^{-6}




Figure 3. The approximate solution for $\alpha = 0.6, 0.7, 0.8, 0.9, 1$.

Example 4. Consider the following class of fractional differential equation:

$$D^{2\alpha}y(x) - 4y(x) + 8y^3(x) - 3y^5(x) = 0, \frac{1}{2} < \alpha \leq 1, \tag{31}$$

subject to:

$$y(0) = y_0 = 1, D^\alpha y(0) = y_1 = 1. \tag{32}$$

Then, the error, when $\alpha = 1$, is reported in Table 4. Figure 4 shows the effect of α on the solution for $\alpha = 0.6, 0.7, 0.8, 0.9, 1$.

Table 4. Error when $\alpha = 1$.

x	Exact Solution	$y_3(x)$	$error(x)$
0.00	1	1	0
0.02	1.01981437	1.01981441	4.0×10^{-8}
0.04	1.03931280	1.03931294	1.4×10^{-7}
0.06	1.05857420	1.05857444	2.4×10^{-7}
0.08	1.07767317	1.07767348	3.1×10^{-7}
0.10	1.09668137	1.09668171	3.4×10^{-7}




Figure 4. The approximate solution for $\alpha = 0.6, 0.7, 0.8, 0.9, 1$.

5. Conclusions

In this paper, we have investigated the analytical solution of Lienard's equation based on the RPS method. Convergence of the proposed infinite series is presented. Four examples of our numerical results are presented. Comparison with the exact solution when $\alpha = 1$ is reported in Tables 1–4. From Tables 1–4, we see that the approximate solute is close to the exact solution with four terms only. From Figures 1 and 2, we see that as α is increasing, the approximate solution is increasing while from Figures 3 and 4, we see that as α is increasing, the approximate solution is decreasing. We see that this approach is cheap compared with other methods and we obtain accurate results using four terms only. A reasonably accurate solution can be achieved with only a few terms. Moreover, the proposed method can be applied to several nonlinear models in science and engineering.

Acknowledgments: In addition, The authors would like to express their appreciation for the valuable comments of the reviewers.

Conflicts of Interest: The author declares no conflict of interest.

References

- Guckenheimer, J. Dynamics of the van der Pol equation. *IEEE Trans. Circuits Syst.* **1980**, *27*, 983.
- Zhang, Z.F.; Ding, T.; Huang, H.W.; Dong, Z.X. *Qualitative Theory of Differential Equations*; Science Press: Peking, China, 1985.
- Feng, Z. On explicit exact solutions for the Lienard equation and its applications. *Phys. Lett. A* **2002**, *293*, 50–56.
- Syam, M.; Attili, B. Numerical solution of singularly perturbed fifth order two point boundary value problem. *Appl. Math. Comput.* **2005**, *170*, 1085–1094.
- Attilia, B.; Furati, K.; Syam, M. An efficient implicit Runge–Kutta method for second order systems. *Appl. Math. Comput.* **2006**, *178*, 229–238.
- Syam, M.; Siyyam, H. Numerical differentiation of implicitly defined curves. *J. Comput. Appl. Math.* **1999**, *108*, 131–144.
- El-Sayed, M.; Syam, M. Electrohydrodynamic instability of a dielectric compressible liquid sheet streaming into an ambient stationary compressible gas. *Arch. Appl. Mech.* **2007**, *77*, 613–626.
- Syam, M. The modified Broyden-variational method for solving nonlinear elliptic differential equations. *Chaos Solitons Fractals* **2007**, *32*, 392–404.
- Liao, S.J. *Beyond Perturbation: Introduction to Homotopy Analysis Method*; Chapman and Hall/CRC Press: Boca Raton, FL, USA, 2003.
- Liao, S.J. On the homotopy analysis method for nonlinear problems. *Appl. Math. Comput.* **2004**, *147*, 499–513.
- Liao, S.J. An approximate solution technique not depending on small parameters: A special example. *Int. J. Non-Linear Mech.* **1995**, *30*, 371–380.

12. Kumar, S.; Kumar, D.; Singh, J. Numerical computation of fractional Black–Scholes equation arising in financial market. *Egypt. J. Basic Appl. Sci.* **2014**, *1*, 177–183.
13. Kong, D. Explicit exact solutions for the Lienard equation and its applications. *Phys. Lett. A* **1995**, *196*, 301–306.
14. Matinfar, M.; Hosseinzadeh, H.; Ghanbari, M. A numerical implementation of the variational iteration method for the Lienard equation. *World J. Model. Simul.* **2008**, *4*, 205–210.
15. Matinfar, M.; Mahdavi, M.; Raeesy, Z. Exact and numerical solution of Lienard’s equation by the variational homotopy perturbation method. *J. Inf. Comput. Sci.* **2011**, *6*, 73–80.
16. Rashidi, M.M.; Rostami, B.; Freidoonimehr, N.; Abbasbandy, S. Free convective heat and mass transfer for MHD fluid flow over a permeable vertical stretching sheet in the presence of the radiation and buoyancy effects. *Ain Shams Eng. J.* **2014**, *5*, 901–912.
17. Rashidi, M.M.; Hassan, H. An analytic solution of micropolar flow in a porous channel with mass injection using homotopy analysis method. *Int. J. Numer. Methods Heat Fluid Flow* **2014**, *24*, 419–437.
18. Khuri, S.A. A Laplace decomposition algorithm applied to a class of nonlinear differential equations. *J. Appl. Math.* **2001**, *1*, 141–155.
19. Khan, M.; Gondal, M.A.; Hussain, I.; Karimi Vanani, S. A new comparative study between homotopy analysis transform method and homotopy perturbation transform method on semi-infinite domain. *Math. Comput. Model.* **2012**, *55*, 1143–1150.
20. Singh, J.; Swroop, R.; Kumar, D. A computational approach for fractional Convection–Diffusion equation via integral transforms. *Ain Shams Eng. J.* **2016**, doi:10.1016/j.asej.2016.04.014.
21. Kumar, S.; Kumar, A.; Argyros, I.K. A new analysis for the Keller–Segel model of fractional order. *Numer. Algorithms* **2016**, doi:10.1007/s11075-016-0202-z.
22. Caponetto, R.; Dongola, G.; Fortuna, L.; Gallo, A. New results on the synthesis of FO-PID controllers. *Commun. Nonlinear Sci. Numer. Simul.* **2010**, *15*, 997–1007.
23. Syam, M. Analytical solution of the Fractional Fredholm integro-differential equation using the modified residual power series method. *Complexity* **2017**, *2017*, 6, doi:10.1155/2017/4573589.
24. Caputo, M. Linear models of dissipation whose Q is almost frequency independent-II. *Geophys. J. R. Astron. Soc.* **1967**, *13*, 529–539.
25. Alquran, M.; Al-Khaled, K.; Sarda, T.; Chattopadhyay, J. Revisited Fisher’s equation in a new outlook: A fractional derivative approach. *Phys. A Stat. Mech. Appl.* **2015**, *438*, 81–93.
26. Jaradat, H.M.; Al-Shara, S.; Khan, Q.J.A.; Alquran, M.; Al-Khaled, K. Analytical solution of time-fractional Drinfeld-Sokolov-Wilson System using residual power series method. *IAENG Int. J. Appl. Math.* **2016**, *46*, 64–70.
27. Abu Arqub, O. Series solution of fuzzy differential equations under strongly generalized differentiability. *J. Adv. Res. Appl. Math.* **2013**, *5*, 31–52.
28. El-Ajou, A.; Abu Arqub, O.; Al zhour, Z.; Momani, S. New results on fractional power series: Theories and applications. *Entropy* **2013**, *15*, 5305–5323.
29. Rudin, W. *Principles of Mathematical Analysis*, 3rd ed.; China Machine Press: Beijing, China, 2004.



© 2017 by the author. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).