

Development of a Screening Method for Sulfamethoxazole in Environmental Water by Digital Colorimetry Using a Mobile Device

Patrícia S. Peixoto ¹, Pedro H. Carvalho ², Ana Machado ^{3,4}, Luisa Barreiros ¹, Adriano A. Bordalo ^{3,4},
Hélder P. Oliveira ^{2,5} and Marcela A. Segundo ^{1,*}

¹ LAQV, REQUIMTE, Faculty of Pharmacy, University of Porto, 4050-313 Porto, Portugal; patriciaspeixoto@gmail.com (P.S.P.); lbarreiros@ff.up.pt (L.B.)

² INESC TEC—Institute for Systems and Computer Engineering, Technology and Science, 4200-465 Porto, Portugal; phmq.carvalho@gmail.com (P.H.C.); helder.f.oliveira@inesctec.pt (H.P.O.)

³ Department of Population Studies, ICBAS, Instituto de Ciências Biomédicas Abel Salazar, Universidade do Porto, 4050-313 Porto, Portugal; ammachado@icbas.up.pt (A.M.); bordalo@icbas.up.pt (A.A.B.)

⁴ CIIMAR—Interdisciplinary Centre of Marine and Environmental Research, 4450-208 Matosinhos, Portugal

⁵ Faculty of Sciences, University of Porto, 4169-007 Porto, Portugal

* Correspondence: msegundo@ff.up.pt; Tel.: +351-220428676




Figure S1. Experimental apparatus for solid-phase extraction performed in the laboratory (A. Extraction disk, B. Placement of disk in the holder, C. Flow set-up for SPE).



Figure S2. Detail of the color palette with numbers assigned to each patch.