

Table S1. Biochemical hyperandrogenism analysed by weight class. Controls vs women with PCOS.

	Controls	PCOS	
	n (%)	n (%)	<i>p</i> -value
Normal weight	n=21	n=21	
Testosterone, ng/mL, median (min, max)	0.27 (0.13, 0.51)	0.34 (0.20, 0.59)	0.012
SHBG, mmol/L, median (min, max)	71.0 (38.0, 142.0)	57.5 (35.0, 110.0)	0.148
FAI, median (min, max)	0.004 (0.00, 0.01)	0.006 (0.00, 0.01)	0.010
Overweight	n=13	n=36	
Testosterone, ng/mL, median (min, max)	0.32 (0.13, 0.70)	0.36 (0.02, 1.29)	0.402
SHBG, mmol/L, median (min, max)	76.0 (32.0, 125.0)	43.0 (19.0, 94.0)	0.005
FAI, median (min, max)	0.005 (0.00, 0.01)	0.009 (0.00, 0.02)	0.003
Overweight and obese	n=18	n=106	
Testosterone, ng/mL, median (min, max)	0.35 (0.13, 0.81)	0.35 (0.02, 1.29)	0.791
SHBG, mmol/L, median (min, max)	59.0 (27.0, 125.0)	39.0 (6.4, 108.0)	0.004
FAI, median (min, max)	0.007 (0.00, 0.01)	0.009 (0.00, 0.06)	0.027

Data were not normally distributed. Mann-Whitney U-test was used.