Supplementary Materials: Cysteine Derivatized ^{99m}Tc-Labelled Fatty Acids as β-Oxidation Markers

Theodoros Tsotakos, Charalambos Triantis, Christos Kiritsis, Aggeliki Panagiotopoulou, Dimitrios Psimadas, Patricia Kyprianidou, Maria Pelecanou, Minas Papadopoulos and Ioannis Pirmettis

Table of Contents

Page S2	Figure S1 HSQC spectrum of Re-Cys-Val
Page S3	Figure S2 HSQC spectrum of Re-Cys-FA11
Page S4	Figure S3 HSQC spectrum of Re-Cys-FA16
Page S5	Figure S4 COSY spectrum of Re-Cys-Val
Page S6	Figure S5 COSY spectrum of Re-Cys-FA11
Page S7	Figure S6 COSY spectrum of Re-Cys-FA16



Figure S1. HSQC spectrum of complex **Re-Cys-Val** in DMSO-*d*₆, at 25 °C. The numbering of the atoms is given in Scheme 1.



Figure S2. HSQC spectrum of complex **Re-Cys-FA11** in DMSO-*d*₆, at 25 °C. The numbering of the atoms is given in Scheme 1.



Figure S3. HSQC spectrum of complex Re-Cys-FA16 in DMSO- d_6 , at 25 °C. The numbering of the atoms is given in Scheme 1.



Figure S4. COSY spectrum (range 6.5–3.8 ppm) of complex Re-Cys-Val in DMSO-d₆ at 25 °C.



Figure S5. COSY spectrum (range 6.5–3.8 ppm) of complex Re-Cys-FA11 in DMSO-d₆ at 25 °C.



Figure S6. COSY spectrum (range 3.1–1.0 ppm) of complex Re-Cys-FA16 in DMSO-d₆ at 25 °C.