

Supplementary Materials: Evaluation of Tannins and Anthocyanins in Marquette, Frontenac, and St. Croix Cold-Hardy Grape Cultivars

Somchai Rice, Jacek A. Koziel, Murlidhar Dharmadhikari and Anne Fennell

Table S1. Chemical composition of Frontenac, Marquette, and St. Croix varieties at veraison and harvest, grown in Ames, Iowa (IA) and Brookings, South Dakota (SD) in 2012.

Location	Cultivar	Time Point	° Brix	pH	TA (g/L)
Iowa	Frontenac	Harvest	23.3	3.3	10.5
Iowa	Frontenac	Veraison	16.6	2.9	20.2
Iowa	Marquette	Harvest	23.5	3.2	9.5
Iowa	Marquette	Veraison	12.3	2.7	27.2
Iowa	St Croix	Harvest	18.4	3.4	5.8
Iowa	St Croix	Veraison	11.5	2.8	21.5
SD	Frontenac	Harvest	24.7	3.1	9.7
SD	Frontenac	Veraison	14.5	2.9	17.9
SD	Marquette	Harvest	26.4	3.2	6.2
SD	Marquette	Veraison	12.7	2.7	18.5
SD	St Croix	Harvest	18.9	3.6	4.4
SD	St Croix	Veraison	10.5	2.8	16.0