

Algicidal Properties of Microbial Fermentation Products on the Inhibiting the Growth of Harmful Dinoflagellate Species

Barathan Balaji Prasath^{1,2,3*}, Ying Wang¹, Yu Ping Su^{1 2,3*}, Minghua Chen⁴, Yi Zheng⁴


¹College of Environmental Science and Engineering, Fujian Normal University, Fuzhou 350007, PR China

²Fujian Key Laboratory of Pollution Control and Resource Recycling, Fuzhou 350007, PR China

³Fujian Province Research Centre for River and Lake Health Assessment, Fuzhou 350007, P. R. China

⁴College of Life Sciences, Fujian Normal University, Fuzhou 350007, PR China


Correspondence: *Yuping Su ypsu@fjnu.edu.cn, College of Environmental Science and Engineering, Fujian Normal University /Fujian Key Laboratory of Pollution Control and Resource Recycling, Fuzhou 350007, PR China.



1(a)


1(b)

1(c)




1(d)


Supplementary Figure S1. Identification of Ba3 by the morphology, plate shape (a); under microscope (b); SEM image (c), and 16S rRNA gene, which was shown as the phylogenetic tree (d).




Supplementary Figure S2. High performance liquid chromatography of elution component 10% ethanol.




Supplementary Figure S3. High performance liquid chromatography of elution component 20% ethanol.




Supplementary Figure S4. High performance liquid chromatography of elution component 30% ethanol.




Supplementary Figure S5. High performance liquid chromatography of elution component 40% ethanol.



Supplementary Figure S6. High performance liquid chromatography of elution component 50% ethanol.



Supplementary Figure S7. Ultraviolet absorption peak of 5.39 min.



Supplementary Figure S8. Ultraviolet absorption peak of 6.20 min.