



Relative Frequency of *Blastocystis* Subtypes 1, 2, and 3 in Urban and Periurban Human Populations of Arequipa, Peru

Kasandra Ascuña-Durand ^{1,*}, Renzo S. Salazar-Sánchez ^{1,2}, Ricardo Castillo-Neyra ^{2,3} and Jorge Ballón-Echegaray ^{2,4,*}

¹ Laboratorio de Microbiología Molecular, Facultad de Medicina, Universidad Nacional de San Agustín, Arequipa, Peru; kasandra.asc@gmail.com (K.A.D.), rendaths@gmail.com (R.S.S.S.)

² Zoonotic Disease Research Laboratory, One Health Unit, School of Public Health and Administration, Universidad Peruana Cayetano Heredia, 4314 Lima, Peru; cricardo@upenn.edu (R.C.N.), jballone@unsa.edu.pe (J.B.E.)

³ Department of Biostatistics, Epidemiology & Informatics, Perelman School of Medicine of the University of Pennsylvania, PA 19104, USA.

⁴ Departamento de Microbiología y Patología, Facultad de Medicina, Universidad Nacional de San Agustín, Santa Catalina 117, Arequipa, Peru; damp@unsa.edu.pe

* Correspondence: kasandra.asc@gmail.com (K.A.D.); jballone@unsa.edu.pe (J.B.E.)

Table A1. *Blastocystis* Subtypes Coinfection with Other Intestinal Protozoa.

Protozoa	Coinfection with Subtype 1	Coinfection with Subtype 3
<i>Entamoeba coli</i>	57.1% (4/7)	57.1% (8/14)
<i>Chilomastix mesnilli</i>	57.1% (4/7)	28.6% (4/14)
<i>Giardia duodenalis</i>	28.8% (2/7)	7.1% (1/14)
<i>Iodamoeba bütschlii</i>	14.3% (1/7)	21.4% (3/14)
<i>Entamoeba histolytica</i>	14.3% (1/7)	14.3% (2/14)
<i>Endolimax nana</i>	14.3% (1/7)	0% (0/14)