

Supplementary

Table S1. Clinical specimens collected for culture in patients diagnosed with histoplasmosis, INI 2000–2018.

Culture	Variables	Diagnosis of HIV		
		Negative	Positive	p-value
Lymphnode biopsy	No	34 (100)	58 (89.2)	0.092
	Yes	0 (0)	7 (10.8)	
Blood	No	30 (88.2)	44 (67.7)	0.025
	Yes	4 (11.8)	21 (32.3)	
Bone Marrow	No	31 (91.2)	41 (63.1)	0.003
	Yes	3 (8.8)	24 (36.9)	
Skin Biopsy	No	31 (91.2)	61 (93.8)	0.689
	Yes	3 (8.8)	4 (6.2)	
Bronchoalveolar Lavage	No	29 (85.3)	61 (93.8)	0.268
	Yes	5 (14.7)	4 (6.2)	
Sputum	No	31 (91.2)	60 (92.3)	1
	Yes	3 (8.8)	5 (7.7)	
Urine	No	34 (100)	64 (98.5)	1
	Yes	0 (0)	1 (1.5)	
Negative culture	No	27 (79.4)	54 (83.1)	0.653
	Yes	7 (20.6)	11 (16.9)	
Total		34	65	

Table S2. Temporal comparison between obtaining the result of the culture and the date of hospitalization until death, INI 2000–2018.

Variables	Diagnosis of HIV		
	Negative	Positive	p-value
Time to receive culture result (in days)	Median (IQR)	12.5 (7-19.5)	13.5 (8.8-27)
Hospitalization time resulting in death	Mean (SD)	12 (10,14)	6 (5,18.5)

IQR = Interquartile Range; SD = Standard Deviation