

Article

Some Double Generalized Weighted Fractional Integral Inequalities Associated with Monotone Chebyshev Functionals

Gauhar Rahman ¹, Saud Fahad Aldosary ², Muhammad Samraiz ³ and Kottakkaran Sooppy Nisar ^{2,*}

¹ Department of Mathematics and Statistics, Hazara University, Mansehra 21300, Pakistan; gauhar55uom@gmail.com or drgauhar.rahman@hu.edu.pk

² Department of Mathematics, College of Arts and Sciences, Prince Sattam Bin Abdulaziz University, Wadi Aldawaser 11991, Saudi Arabia; sau.aldosary@psau.edu.sa

³ Department of Mathematics, University of Sargodha, Sargodha 40100, Pakistan; muhammad.samraiz@uos.edu.pk

* Correspondence: n.sooppy@psau.edu.sa or ksnisar1@gmail.com

Abstract: In this manuscript, we study the unified integrals recently defined by Rahman et al. and present some new double generalized weighted type fractional integral inequalities associated with increasing, positive, monotone and measurable function \mathcal{F} . Also, we establish some new double-weighted inequalities, which are particular cases of the main result and are represented by corollaries. These inequalities are further refinement of all other inequalities associated with increasing, positive, monotone and measurable function existing in literature. The existing inequalities associated with increasing, positive, monotone and measurable function are also restored by applying specific conditions as given in Remarks. Many other types of fractional integral inequalities can be obtained by applying certain conditions on \mathcal{F} and Ψ given in the literature.



Citation: Rahman, G.; Aldosary, S.F.; Samraiz, M.; Nisar, K.S. Some Double Generalized Weighted Fractional Integral Inequalities Associated with Monotone Chebyshev Functionals. *Fractal Fract.* **2021**, *5*, 275. <https://doi.org/10.3390/fractfract5040275>

Academic Editor: Norbert Herencsar

Received: 12 November 2021

Accepted: 8 December 2021

Published: 15 December 2021

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

In the context of fractional differential equations, integral inequalities are very significant. This field has gained popularity during the last few decades. Various researchers, such as [1–3], have investigated the significant developments in this domain. By employing Riemann-Liouville (R-L) fractional integrals, the authors presented Grüss type and several other new inequalities in [4,5]. Certain inequalities for the generalised (k, ρ) -fractional integral operator are proposed in [6]. In [7], the modified Hermite-Hadamard type inequalities can be found. Dahmani [8] discovered various fractional integral inequalities employing a family of n positive functions. In [9], Srivastava et al. presented the Chebyshev inequality by employing general family of fractional integral operators. Some remarkable inequalities and their applications can be found in the work of [10–15].

In [16,17], the Chebyshev functional for the integrable functions \mathcal{Z}_1 and \mathcal{Z}_2 on $[v_1, v_2]$, is given by

$$\mathcal{H}(\mathcal{Z}_1, \mathcal{Z}_2, h_1) = \int_{v_1}^{v_2} h_1(\theta) d\theta \int_{v_1}^{v_2} h_1(\theta) \mathcal{Z}_1(\theta) \mathcal{Z}_2(\theta) d\theta - \int_{v_1}^{v_2} h_1(\theta) \mathcal{Z}_1(\theta) d\theta \int_{v_1}^{v_2} h_1(\theta) \mathcal{Z}_2(\theta) d\theta, \quad (1)$$

where the function h_1 is a positive and integrable on $[v_1, v_2]$.

The following extended Chebyshev functional for the integrable functions \mathcal{Z}_1 and \mathcal{Z}_2 on $[v_1, v_2]$ can be found in [5,18] by

$$\begin{aligned}\mathcal{H}(\mathcal{Z}_1, \mathcal{Z}_2, \hbar_1, \hbar_2) = & \int_{v_1}^{v_2} \hbar_2(\theta) d\theta \int_{v_1}^{v_2} \hbar_1(\theta) \mathcal{Z}_1(\theta) \mathcal{Z}_2(\theta) d\theta + \int_{v_1}^{v_2} \hbar_1(\theta) d\theta \int_{v_1}^{v_2} \hbar_2(\theta) \mathcal{Z}_1(\theta) \mathcal{Z}_2(\theta) d\theta \\ & - \int_{v_1}^{v_2} \hbar_1(\theta) \mathcal{Z}_1(\theta) d\theta \int_{v_1}^{v_2} \hbar_2(\theta) \mathcal{Z}_2(\theta) d\theta - \int_{v_1}^{v_2} \hbar_2(\theta) \mathcal{Z}_1(\theta) d\theta \int_{v_1}^{v_2} \hbar_1(\theta) \mathcal{Z}_2(\theta) d\theta.\end{aligned}\quad (2)$$

where the two functions \hbar_1 and \hbar_2 are positive and integrable on $[v_1, v_2]$.

Kuang [19] and Mitrinovic [18] proved that $\mathcal{H}(\mathcal{Z}_1, \mathcal{Z}_2, \hbar_1) \geq 0$ and $\mathcal{H}(\mathcal{Z}_1, \mathcal{Z}_2, \hbar_1, \hbar_2) \geq 0$ if the functions \mathcal{Z}_1 and \mathcal{Z}_2 are synchronous on $[v_1, v_2]$.

Remark 1. If we take $\hbar_1(\theta) = \hbar_2(\theta) = 1, \theta \in [v_1, v_2]$, then

$$\mathcal{H}(\mathcal{Z}_1, \mathcal{Z}_2, \hbar_1) = \frac{1}{2} \mathcal{H}(\mathcal{Z}_1, \mathcal{Z}_2, \hbar_1, \hbar_2).$$

Certain remarkable integral inequalities associated with the Chebyshev's functionals (1) and (2) can be found in the work of [20–25].

Awan et al. [26] proposed the following inequality by:

Theorem 1. Let the function g be an absolutely continuous on $[v_1, v_2]$, and \hbar_1 be integrable and positive function on $[v_1, v_2]$ and $(g')^2 \in L_1[v_1, v_2]$, then the following inequality holds;

$$\mathcal{H}(\Phi, \Phi, \hbar_1) \leq \frac{1}{G^2(v_2)} \int_{v_1}^{v_2} \left[\int_{v_1}^{\theta} \hbar_1(\varrho) d\varrho \int_{v_1}^{v_2} \varrho \hbar_1(\varrho) d\varrho - \int_{v_1}^{v_2} \hbar_1(\varrho) d\varrho \int_{v_1}^{\theta} \varrho \hbar_1(\varrho) d\varrho \right] [\Phi'(\theta)]^2 d\theta,$$

where $G(v_2) = \int_{v_1}^{v_2} \hbar_1(\varrho) d\varrho$.

In [27], Bezzou et al. proposed the below result for Riemann-Liouville fractional integral as follows:

Theorem 2. Assume that the function $g : [v_1, v_2] \rightarrow \mathbb{R}$ be an absolutely continuous function, and the function $\hbar_1 : [v_1, v_2] \rightarrow \mathbb{R}^+$ be an integrable, and $(g')^2 \in L_1[v_1, v_2]$. Then the following inequality for $\kappa > 0$ holds:

$$\mathcal{I}_{v_1+}^{\kappa} \hbar_1(v_2) \mathcal{I}_{v_1+}^{\kappa} \hbar_1 g^2(v_2) - (\mathcal{I}_{v_1+}^{\kappa} \hbar_1 g(v_2))^2 \leq \int_{v_1}^{v_2} Y(\theta) [g'(\theta)]^2 d\theta,$$

where

$$Y(\theta) = \frac{1}{2} \left[\mathcal{I}_{v_1+}^{\kappa} v_2 \hbar_1(v_2) \int_{v_1}^{\theta} (v_2 - \varrho)^{\kappa-1} \hbar_1(\varrho) d\varrho - \mathcal{I}_{v_1+}^{\kappa} \hbar_1(v_2) \int_{v_1}^{\theta} \varrho (v_2 - \varrho)^{\kappa-1} \hbar_1(\varrho) d\varrho \right].$$

Dahmani and Bounoua [28] proposed the following inequality for Riemann-Liouville fractional integral by:

Theorem 3. If the function $g : [v_1, v_2] \rightarrow \mathbb{R}$ be an absolutely continuous and let $\hbar_1 : [v_1, v_2] \rightarrow \mathbb{R}^+$ be an integrable function. If $(\Phi')^2 \in L_1[v_1, v_2]$, then for all $\kappa > 0$, and $\theta \in [v_1, v_2]$, the following inequality holds;

$$\frac{1}{\mathcal{I}_{v_1+}^{\kappa} \hbar_1(\theta)} \mathcal{I}_{v_1+}^{\theta} (\hbar_1 g^2)(\theta) - \left[\frac{1}{\mathcal{I}_{v_1+}^{\kappa} \hbar_1(\theta)} \mathcal{I}_{v_1+}^{\kappa} (\hbar_1 g)(\theta) \right]^2 \leq \frac{1}{[\mathcal{I}_{v_1+}^{\kappa} \hbar_1(\theta)]^2} \int_{v_1}^{\theta} P_x(\theta) [g'(\theta)]^2 d\theta,$$

with

$$P_x(\theta) = \frac{1}{\Gamma(\kappa)} \left[\mathcal{I}_{v_1+}^{\kappa} (x \hbar_1(x)) \int_{v_1}^{\theta} \hbar_1(\varrho) (x - \varrho)^{\kappa-1} d\varrho - \mathcal{J}_{v_1+}^{\kappa} \hbar_1(x) \int_{v_1}^{\theta} \varrho \hbar_1(\varrho) (x - \varrho)^{\kappa-1} d\varrho \right].$$

Definition 1 ([29]). Suppose that the function $\Psi : [0, \infty) \rightarrow [0, \infty)$ be satisfying the conditions given below:

$$\int_0^1 \frac{\Psi(\varrho)}{\varrho} d\varrho < \infty, \quad (3)$$

$$\frac{1}{P} \leq \frac{\Psi(\hbar_1)}{\Psi(\hbar_2)} \leq P, \frac{1}{2} \leq \frac{\hbar_1}{\hbar_2} \leq 2, \quad (4)$$

$$\frac{\Psi(\hbar_2)}{\hbar_2^2} \leq Q \frac{\Psi(\hbar_1)}{\hbar_1^2}, \hbar_1 \leq \hbar_2, \quad (5)$$

$$\left| \frac{\Psi(\hbar_2)}{\hbar_2^2} - \frac{\Psi(\hbar_1)}{\hbar_1^2} \right| \leq S |\hbar_2 - \hbar_1| \frac{\Psi(\hbar_2)}{\hbar_2^2}, \frac{1}{2} \leq \frac{\hbar_1}{\hbar_2} \leq 2, \quad (6)$$

where $P, Q, S > 0$ and are independent of $\hbar_1, \hbar_2 > 0$. If $\Psi(\hbar_2)\hbar_2^\alpha$ is increasing for some $\alpha > 0$ and $\frac{\Psi(\hbar_2)}{\hbar_2^\beta}$ is decreasing for some $\beta > 0$, then Ψ satisfies (3)–(6).

Next, we recall the following generalized weighted type fractional integral operators recently proposed by Rahman et al. [30].

Definition 2. The generalized weighted type fractional integral operators both left and right sided are respectively defined by:

$$\left({}_{\omega}^{\mathcal{F}} \mathcal{I}_{v_1+}^{\Psi} \mathcal{Z}_1 \right)(\theta) = \omega^{-1}(\theta) \int_{v_1}^{\theta} \frac{\Psi(\mathcal{F}(\theta) - \mathcal{F}(\varrho))}{\mathcal{F}(\theta) - \mathcal{F}(\varrho)} \omega(\varrho) \mathcal{F}'(\varrho) \mathcal{Z}_1(\varrho) d\varrho, v_1 < \theta \quad (7)$$

and

$$\left({}_{\omega}^{\mathcal{F}} \mathcal{I}_{v_2-}^{\Psi} \mathcal{Z}_1 \right)(\theta) = \omega^{-1}(\theta) \int_{\theta}^{v_2} \frac{\Psi(\mathcal{F}(\varrho) - \mathcal{F}(\theta))}{\mathcal{F}(\varrho) - \mathcal{F}(\theta)} \omega(\varrho) \mathcal{F}'(\varrho) \mathcal{Z}_1(\varrho) d\varrho, v_2 > \theta. \quad (8)$$

Remark 2. 1. If we consider $\Psi(\mathcal{F}(\theta)) = \mathcal{F}(\theta)$, the fractional integrals (7) and (8) reduce to the following:

$$\left({}_{\omega}^{\mathcal{F}} \mathcal{I}_{v_1+} \mathcal{Z}_1 \right)(\theta) = \omega^{-1}(\theta) \int_{v_1}^{\theta} \omega(\varrho) \mathcal{F}'(\varrho) \mathcal{Z}_1(\varrho) d\varrho, v_1 < \theta$$

and

$$\left({}_{\omega}^{\mathcal{F}} \mathcal{I}_{v_2-} \mathcal{Z}_1 \right)(\theta) = \omega^{-1}(\theta) \int_{\theta}^{v_2} \omega(\varrho) \mathcal{F}'(\varrho) \mathcal{Z}_1(\varrho) d\varrho, v_2 > \theta,$$

respectively.

2. If we consider $\mathcal{F}(\theta) = \theta$, the fractional integrals (7) and (8) reduce to the following respectively

$$\left({}_{\omega}^{\mathcal{F}} \mathcal{I}_{v_1+} \mathcal{Z}_1 \right)(\theta) = \omega^{-1}(\theta) \int_{v_1}^{\theta} \frac{\Psi(\theta - \varrho)}{\theta - \varrho} \omega(\varrho) \mathcal{Z}_1(\varrho) d\varrho, v_1 < \theta$$

and

$$\left({}_{\omega}^{\mathcal{F}} \mathcal{I}_{v_2-} \mathcal{Z}_1 \right)(\theta) = \omega^{-1}(\theta) \int_{\theta}^{v_2} \frac{\Psi(\varrho - \theta)}{\varrho - \theta} \omega(\varrho) \mathcal{Z}_1(\varrho) d\varrho, v_2 > \theta.$$

3. If we consider $\Psi(\mathcal{F}(\theta)) = \frac{\mathcal{F}(\theta)^\kappa}{\Gamma(\kappa)}$, the fractional integrals (7) and (8) reduce to the following respectively (see [31]):

$$\left({}_{\omega}^{\mathcal{F}} \mathcal{I}_{v_1}^{\kappa} \mathcal{Z}_1 \right)(\theta) = \frac{\omega^{-1}(\theta)}{\Gamma(\kappa)} \int_{v_1}^{\theta} (\mathcal{F}(\theta) - \mathcal{F}(\varrho))^{\kappa-1} \omega(\varrho) \mathcal{F}'(\varrho) \mathcal{Z}_1(\varrho) d\varrho, v_1 < \theta$$

and

$$\left({}_{\omega}^{\mathcal{F}} \mathcal{I}_{v_2-}^{\kappa} \mathcal{Z}_1 \right)(\theta) = \frac{\omega^{-1}(\theta)}{\Gamma(\kappa)} \int_{\theta}^{v_2} (\mathcal{F}(\varrho) - \mathcal{F}(\theta))^{\kappa-1} \omega(\varrho) \mathcal{F}'(\varrho) \mathcal{Z}_1(\varrho) d\varrho, v_2 > \theta,$$

where $\kappa \in \mathbb{C}$ with $\Re(\kappa) > 0$.

4. If we consider $\mathcal{F}(\theta) = \theta$ and $\Psi(\mathcal{F}(\theta)) = \frac{\theta^\kappa}{\Gamma(\kappa)}$, the fractional integrals (7) and (8) reduce to the following:

$$\left({}_{\omega} \mathcal{I}_{v_1+}^{\kappa} \mathcal{Z}_1 \right)(\theta) = \frac{\omega^{-1}(\theta)}{\Gamma(\kappa)} \int_{v_1}^{\theta} (\theta - \varrho)^{\kappa-1} \omega(\varrho) \mathcal{Z}_1(\varrho) d\varrho, v_1 < \theta$$

and

$$\left({}_{\omega} \mathcal{I}_{v_2-}^{\kappa} \mathcal{Z}_1 \right)(\theta) = \frac{\omega^{-1}(\theta)}{\Gamma(\kappa)} \int_{\theta}^{v_2} (\varrho - \theta)^{\kappa-1} \omega(\varrho) \mathcal{Z}_1(\varrho) d\varrho, v_2 > \theta,$$

respectively.

5. If we consider $\mathcal{F}(\theta) = \ln \theta$ and $\Psi(\mathcal{F}(\theta)) = \frac{(\ln \theta)^\kappa}{\Gamma(\kappa)}$, the fractional integrals (7) and (8) reduce to the following weighted Hadamard fractional integrals:

$$\left({}_{\omega} \mathcal{I}_{v_1+}^{\kappa} \mathcal{Z}_1 \right)(\theta) = \frac{\omega^{-1}(\theta)}{\Gamma(\kappa)} \int_{v_1}^{\theta} (\ln \theta - \ln \varrho)^{\kappa-1} \omega(\varrho) \mathcal{Z}_1(\varrho) \frac{d\varrho}{\varrho}, v_1 < \theta$$

and

$$\left({}_{\omega} \mathcal{I}_{v_2-}^{\kappa} \mathcal{Z}_1 \right)(\theta) = \frac{\omega^{-1}(\theta)}{\Gamma(\kappa)} \int_{\theta}^{v_2} (\ln \varrho - \ln \theta)^{\kappa-1} \omega(\varrho) \mathcal{Z}_1(\varrho) \frac{d\varrho}{\varrho}, v_2 > \theta.$$

6. If we consider $\mathcal{F}(\theta) = \theta^\eta$ and $\Psi(\mathcal{F}(\theta)) = \frac{\theta^\eta}{\eta}$, $\eta > 0$, the fractional integrals (7) and (8) reduce to the following weighted Katugampola fractional integrals,

$$\left({}_{\omega} \mathcal{I}_{v_1}^{\kappa} \mathcal{Z}_1 \right)(\theta) = \frac{\omega^{-1}(\theta)}{\Gamma(\kappa)} \int_{v_1}^{\theta} \left(\frac{\theta^\eta - \varrho^\eta}{\eta} \right)^{\kappa-1} \omega(\varrho) \mathcal{Z}_1(\varrho) \frac{d\varrho}{\varrho^{1-\eta}}, v_1 < \theta$$

and

$$\left({}_{\omega} \mathcal{I}_{v_2}^{\kappa} \mathcal{Z}_1 \right)(\theta) = \frac{\omega^{-1}(\theta)}{\Gamma(\kappa)} \int_{\theta}^{v_2} \left(\frac{\varrho^\eta - \theta^\eta}{\eta} \right)^{\kappa-1} \omega(\varrho) \mathcal{Z}_1(\varrho) \frac{d\varrho}{\varrho^{1-\eta}}, v_2 > \theta.$$

7. If we consider $\mathcal{F}(\theta) = \theta$ and $\Psi(\mathcal{F}(\theta)) = \frac{\theta}{\eta} \exp\left(-\frac{1-\eta}{\eta}\theta\right)$, $\eta \in (0, 1)$, the fractional integrals (7) and (8) reduce to the following weighted fractional integrals,

$$\left({}_{\omega} \mathcal{I}_{v_1+}^{\eta} \mathcal{Z}_1 \right)(\theta) = \frac{\omega^{-1}(\theta)}{\eta} \int_{v_1}^{\theta} \exp\left(-\frac{1-\eta}{\eta}(\theta - \varrho)\right) \omega(\varrho) \mathcal{Z}_1(\varrho) d\varrho, v_1 < \theta$$

and

$$\left({}_{\omega} \mathcal{I}_{v_2-}^{\eta} \mathcal{Z}_1 \right)(\theta) = \frac{\omega^{-1}(\theta)}{\eta} \int_{\theta}^{v_2} \exp\left(-\frac{1-\eta}{\eta}(\varrho - \theta)\right) \omega(\varrho) \mathcal{Z}_1(\varrho) d\varrho, v_2 > \theta.$$

Also, one can derive the weighted form of conformable fractional integrals introduced by [32–35]

The following special cases can be easily obtained by applying the conditions on $\omega(\theta)$ and $\Psi(\mathcal{F}(\theta))$.

Remark 3. 1. If we consider $\omega(\theta) = 1$ and $\Psi(\mathcal{F}(\theta)) = \mathcal{F}(\theta)$, the fractional integrals (7) and (8) reduce to the following:

$$({}^{\mathcal{F}}\mathcal{I}_{v_1+}\mathcal{Z}_1)(\theta) = \int_{v_1}^{\theta} \mathcal{F}'(\varrho) \mathcal{Z}_1(\varrho) d\varrho, v_1 < \theta$$

and

$$({}^{\mathcal{F}}\mathcal{I}_{v_2-}\mathcal{Z}_1)(\theta) = \int_{\theta}^{v_2} \mathcal{F}'(\varrho) \mathcal{Z}_1(\varrho) d\varrho, v_2 > \theta,$$

respectively.

2. If we consider $\omega(\theta) = 1$ and $\mathcal{F}(\theta) = \theta$, the fractional integrals (7) and (8) reduce to the following respectively (see [36]) as follows:

$$({}^{\mathcal{F}}\mathcal{I}_{v_1+}\mathcal{Z}_1)(\theta) = \int_{v_1}^{\theta} \frac{\Psi(\theta - \varrho)}{\theta - \varrho} \mathcal{Z}_1(\varrho) d\varrho, v_1 < \theta$$

and

$$({}^{\mathcal{F}}\mathcal{I}_{v_2-}\mathcal{Z}_1)(\theta) = \int_{\theta}^{v_2} \frac{\Psi(\varrho - \theta)}{\varrho - \theta} \mathcal{Z}_1(\varrho) d\varrho, v_2 > \theta.$$

3. If we consider $\omega(\theta) = 1$ and $\Psi(\mathcal{F}(\theta)) = \frac{\mathcal{F}(\theta)^{\kappa}}{\Gamma(\kappa)}$, the fractional integrals (7) and (8) reduce to the following respectively (see [37,38]):

$$({}^{\mathcal{F}}\mathcal{I}_{v_1}^{\kappa}\mathcal{Z}_1)(\theta) = \frac{1}{\Gamma(\kappa)} \int_{v_1}^{\theta} (\mathcal{F}(\theta) - \mathcal{F}(\varrho))^{\kappa-1} \mathcal{F}'(\varrho) \mathcal{Z}_1(\varrho) d\varrho, v_1 < \theta$$

and

$$({}^{\mathcal{F}}\mathcal{I}_{v_2}^{\kappa}\mathcal{Z}_1)(\theta) = \frac{1}{\Gamma(\kappa)} \int_{\theta}^{v_2} (\mathcal{F}(\varrho) - \mathcal{F}(\theta))^{\kappa-1} \mathcal{F}'(\varrho) \mathcal{Z}_1(\varrho) d\varrho, v_2 > \theta,$$

where $\kappa \in \mathbb{C}$ with $\Re(\kappa) > 0$.

4. If we consider $\omega(\theta) = 1$, $\mathcal{F}(\theta) = \theta$ and $\Psi(\mathcal{F}(\theta)) = \frac{\theta^{\kappa}}{\Gamma(\kappa)}$, the fractional integrals (7) and (8) reduce to the following (see [37,38]):

$$(\mathcal{I}_{v_1+}^{\kappa}\mathcal{Z}_1)(\theta) = \frac{1}{\Gamma(\kappa)} \int_{v_1}^{\theta} (\theta - \varrho)^{\kappa-1} \mathcal{Z}_1(\varrho) d\varrho, v_1 < \theta$$

and

$$(\mathcal{I}_{v_2-}^{\kappa}\mathcal{Z}_1)(\theta) = \frac{1}{\Gamma(\kappa)} \int_{\theta}^{v_2} (\varrho - \theta)^{\kappa-1} \mathcal{Z}_1(\varrho) d\varrho, v_2 > \theta,$$

respectively.

5. If we consider $\omega(\theta) = 1$, $\mathcal{F}(\theta) = \ln \theta$ and $\Psi(\mathcal{F}(\theta)) = \frac{(\ln \theta)^{\kappa}}{\Gamma(\kappa)}$, the fractional integrals (7) and (8) reduce to the following weighted Hadamard fractional integrals (see [37,38]):

$$(\mathcal{I}_{v_1+}^{\kappa}\mathcal{Z}_1)(\theta) = \frac{1}{\Gamma(\kappa)} \int_{v_1}^{\theta} (\ln \theta - \ln \varrho)^{\kappa-1} \mathcal{Z}_1(\varrho) \frac{d\varrho}{\varrho}, v_1 < \theta$$

and

$$(\mathcal{I}_{v_2-}^{\kappa} \mathcal{Z}_1)(\theta) = \frac{1}{\Gamma(\kappa)} \int_{\theta}^{v_2} (\ln \varrho - \ln \theta)^{\kappa-1} \mathcal{Z}_1(\varrho) \frac{d\varrho}{\varrho}, \quad v_2 > \theta.$$

6. If we consider $\omega(\theta) = 1$, $\mathcal{F}(\theta) = \theta^{\eta}$ and $\Psi(\mathcal{F}(\theta)) = \frac{\theta^{\eta}}{\eta}$, $\eta > 0$, the fractional integrals (7) and (8) reduce to the following Katugampola [39] fractional integrals respectively,

$$(\mathcal{I}_{v_1}^{\kappa} \mathcal{Z}_1)(\theta) = \frac{1}{\Gamma(\kappa)} \int_{v_1}^{\theta} \left(\frac{\theta^{\eta} - \varrho^{\eta}}{\eta} \right)^{\kappa-1} \mathcal{Z}_1(\varrho) \frac{d\varrho}{\varrho^{1-\eta}}, \quad v_1 < \theta$$

and

$$(\mathcal{I}_{v_2}^{\kappa} \mathcal{Z}_1)(\theta) = \frac{1}{\Gamma(\kappa)} \int_{\theta}^{v_2} \left(\frac{\varrho^{\eta} - \theta^{\eta}}{\eta} \right)^{\kappa-1} \mathcal{Z}_1(\varrho) \frac{d\varrho}{\varrho^{1-\eta}}, \quad v_2 > \theta.$$

7. If we consider $\omega(\theta) = 1$, $\mathcal{F}(\theta) = \theta$ and $\Psi(\mathcal{F}(\theta)) = \frac{\theta}{\eta} \exp\left(-\frac{1-\eta}{\eta}\theta\right)$, $\eta \in (0, 1)$, the fractional integrals (7) and (8) reduce to the following weighted fractional integrals,

$$(\mathcal{I}_{v_1+}^{\eta} \mathcal{Z}_1)(\theta) = \frac{1}{\eta} \int_{v_1}^{\theta} \exp\left(-\frac{1-\eta}{\eta}(\theta - \varrho)\right) \mathcal{Z}_1(\varrho) d\varrho, \quad v_1 < \theta$$

and

$$(\mathcal{I}_{v_2-}^{\eta} \mathcal{Z}_1)(\theta) = \frac{1}{\eta} \int_{\theta}^{v_2} \exp\left(-\frac{1-\eta}{\eta}(\varrho - \theta)\right) \mathcal{Z}_1(\varrho) d\varrho, \quad v_2 > \theta.$$

Similarly, (7) and (8) will lead to the fractional integrals defined by [32–35].

2. Some Double-Weighted Generalized Fractional Integral Inequalities

In this section, some double-weighted generalized fractional integral inequalities are presented. To this end, we begin by proving the following Lemma.

Lemma 1. Let the function \mathcal{F} be measurable, increasing, positive and monotone function on (v_1, v_2) , and has a continuous derivative \mathcal{F}' on $[v_1, v_2]$. If $\Phi : [v_1, v_2] \rightarrow \mathbb{R}$ is continuous on $[v_1, v_2]$, and $\hbar_1, \hbar_2 : [u, v] \rightarrow \mathbb{R}^+$ are positive integrable. Then, we have

$$\begin{aligned} & \left[\frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^{\Psi} \hbar_1(v_2) \right] \left[\frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^{\Psi} v_2(\hbar_2 \Phi)(v_2) \right] + \left[\frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^{\Psi} \Phi(v_2) \right] \left[\frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^{\Psi} v_2(\hbar_1 \Phi)(v_2) \right] \\ & - \left[\frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^{\Psi} v_2 \hbar_1(v_2) \right] \left[\frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^{\Psi} (\hbar_2 \Phi)(v_2) \right] - \left[\frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^{\Psi} v_2 \hbar_2(v_2) \right] \left[\frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^{\Psi} (\hbar_1 \Phi)(v_2) \right] \\ & = \frac{1}{\omega(\theta)} \int_{v_1}^{v_2} \left[\frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^{\Psi} v_2 \hbar_1(v_2) \right] \int_{v_1}^{\theta} \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\varrho))}{\mathcal{F}(v_2) - \mathcal{F}(\varrho)} \omega(\varrho) \mathcal{F}'(\varrho) \hbar_2(\varrho) d\varrho \\ & - \frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^{\Psi} \hbar_1(v_2) \int_{v_1}^{\theta} \varrho \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\varrho))}{\mathcal{F}(v_2) - \mathcal{F}(\varrho)} \omega(\varrho) \mathcal{F}'(\varrho) \hbar_2(\varrho) d\varrho] (\Phi'(\theta)) d\theta. \end{aligned} \quad (9)$$

Proof. Assume that $\mathcal{Z}_1 : [v_1, v_2] \rightarrow \mathbb{R}$ is a continuous function on $[v_1, v_2]$. Then, one may gets

$$\begin{aligned} & \left[\frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^{\Psi} \hbar_2(v_2) \right] \left[\frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^{\Psi} (\hbar_2 \mathcal{Z}_1 \Phi)(v_2) \right] + \left[\frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^{\Psi} \Phi(v_2) \right] \left[\frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^{\Psi} (\hbar_1 \mathcal{Z}_1 \Phi)(v_2) \right] \\ & - \left[\frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^{\Psi} (\hbar_1 \mathcal{Z}_1)(v_2) \right] \left[\frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^{\Psi} (\hbar_2 \Phi)(v_2) \right] - \left[\frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^{\Psi} (\hbar_2 \mathcal{Z}_1)(v_2) \right] \left[\frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^{\Psi} (\hbar_1 \Phi)(v_2) \right] \\ & = \frac{1}{\omega^2(\theta)} \int_{v_1}^{v_2} \int_{v_1}^{\varrho} \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\xi))}{\mathcal{F}(v_2) - \mathcal{F}(\xi)} \omega(\xi) \mathcal{F}'(\xi) \\ & \times \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\varrho))}{\mathcal{F}(v_2) - \mathcal{F}(\varrho)} \omega(\varrho) \mathcal{F}'(\varrho) \hbar_1(\xi) \hbar_2(\varrho) [(\mathcal{Z}_1(\xi) - \mathcal{Z}_1(\varrho))(\Phi(\xi) - \Phi(\varrho))] d\xi d\varrho. \end{aligned}$$

Consequently, it follows

$$\begin{aligned}
& \left[{}_{\omega}^{\mathcal{F}} \mathcal{I}_{v_1+}^{\Psi} h_1(v_2) \right] \left[{}_{\omega}^{\mathcal{F}} \mathcal{I}_{v_1+}^{\Psi} (h_2 \mathcal{Z}_1 \Phi)(v_2) \right] + \left[{}_{\omega}^{\mathcal{F}} \mathcal{I}_{v_1+}^{\Psi} \Phi(v_2) \right] \left[{}_{\omega}^{\mathcal{F}} \mathcal{I}_{v_1+}^{\Psi} (h_1 \mathcal{Z}_1 \Phi)(v_2) \right] \\
& - \left[{}_{\omega}^{\mathcal{F}} \mathcal{I}_{v_1+}^{\Psi} (h_1 \mathcal{Z}_1)(v_2) \right] \left[{}_{\omega}^{\mathcal{F}} \mathcal{I}_{v_1+}^{\Psi} (h_2 \Phi)(v_2) \right] - \left[{}_{\omega}^{\mathcal{F}} \mathcal{I}_{v_1+}^{\Psi} (h_2 \mathcal{Z}_1)(v_2) \right] \left[{}_{\omega}^{\mathcal{F}} \mathcal{I}_{v_1+}^{\Psi} (h_1 \Phi)(v_2) \right] \\
& = \frac{1}{\omega^2(\theta)} \int_{v_1}^{v_2} \int_{v_1}^{v_2} \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\xi))}{\mathcal{F}(v_2) - \mathcal{F}(\xi)} \omega(\xi) \mathcal{F}'(\xi) \\
& \times \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\varrho))}{\mathcal{F}(v_2) - \mathcal{F}(\varrho)} \omega(\varrho) \mathcal{F}'(\varrho) h_1(\xi) h_2(\varrho) (\mathcal{Z}_1(\xi) - \mathcal{Z}_1(\varrho)) \left(\int_{\xi}^{\varrho} \Phi'(\vartheta) d\vartheta \right) d\xi d\varrho. \quad (10)
\end{aligned}$$

By utilizing the given condition $v_1 \leq \varrho \leq \theta \leq \xi \leq v_2$, we get

$$\begin{aligned}
& \left[{}_{\omega}^{\mathcal{F}} \mathcal{I}_{v_1+}^{\Psi} h_1(v_2) \right] \left[{}_{\omega}^{\mathcal{F}} \mathcal{I}_{v_1+}^{\Psi} (h_2 \mathcal{Z}_1 \Phi)(v_2) \right] + \left[{}_{\omega}^{\mathcal{F}} \mathcal{I}_{v_1+}^{\Psi} \Phi(v_2) \right] \left[{}_{\omega}^{\mathcal{F}} \mathcal{I}_{v_1+}^{\Psi} (h_1 \mathcal{Z}_1 \Phi)(v_2) \right] \\
& - \left[{}_{\omega}^{\mathcal{F}} \mathcal{I}_{v_1+}^{\Psi} (h_1 \mathcal{Z}_1)(v_2) \right] \left[{}_{\omega}^{\mathcal{F}} \mathcal{I}_{v_1+}^{\Psi} (h_2 \Phi)(v_2) \right] - \left[{}_{\omega}^{\mathcal{F}} \mathcal{I}_{v_1+}^{\Psi} (h_2 \mathcal{Z}_1)(x_2) \right] \left[{}_{\omega}^{\mathcal{F}} \mathcal{I}_{v_1+}^{\Psi} (h_1 \Phi)(x_2) \right] \\
& = \frac{1}{\omega^2(\theta)} \int_{v_1}^{v_2} \left[\int_{v_1}^{\theta} \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\varrho))}{\mathcal{F}(v_2) - \mathcal{F}(\varrho)} \omega(\varrho) \mathcal{F}'(\varrho) h_2(\varrho) \right. \\
& \times \left. \int_{v_1}^{v_2} \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\xi))}{\mathcal{F}(v_2) - \mathcal{F}(\xi)} (\mathcal{Z}_1(\xi) - \mathcal{Z}_1(\varrho)) \omega(\xi) \mathcal{F}'(\xi) h_1(\xi) d\xi d\varrho \right] (\Phi'(\theta)) d\theta. \quad (11)
\end{aligned}$$

Applying (11) for the particular case when $\mathcal{Z}_1(v) = v$, then we can write

$$\begin{aligned}
& \left[{}_{\omega}^{\mathcal{F}} \mathcal{I}_{v_1+}^{\Psi} h_1(v_2) \right] \left[{}_{\omega}^{\mathcal{F}} \mathcal{I}_{v_1+}^{\Psi} v_2(h_2 \Phi)(v_2) \right] + \left[{}_{\omega}^{\mathcal{F}} \mathcal{I}_{v_1+}^{\Psi} \Phi(v_2) \right] \left[{}_{\omega}^{\mathcal{F}} \mathcal{I}_{v_1+}^{\Psi} v_2(h_1 \Phi)(v_2) \right] \\
& - \left[{}_{\omega}^{\mathcal{F}} \mathcal{I}_{v_1+}^{\Psi} v_2 h_1(v_2) \right] \left[{}_{\omega}^{\mathcal{F}} \mathcal{I}_{v_1+}^{\Psi} (h_2 \Phi)(v_2) \right] - \left[{}_{\omega}^{\mathcal{F}} \mathcal{I}_{v_1+}^{\Psi} v_2 h_2(v_2) \right] \left[{}_{\omega}^{\mathcal{F}} \mathcal{I}_{v_1+}^{\Psi} (h_1 \Phi)(v_2) \right] \\
& = \frac{1}{\omega^2(\theta)} \int_{v_1}^{v_2} \left[\int_{v_1}^{\theta} \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\varrho))}{\mathcal{F}(v_2) - \mathcal{F}(\varrho)} \omega(\varrho) \mathcal{F}'(\varrho) h_2(\varrho) \right. \\
& \times \left. \int_{v_1}^{v_2} \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\xi))}{\mathcal{F}(v_2) - \mathcal{F}(\xi)} (\xi - \varrho) \omega(\xi) \mathcal{F}'(\xi) h_1(\xi) d\xi d\varrho \right] (\Phi'(\theta)) d\theta \\
& = \frac{1}{\omega(\theta)} \int_{v_1}^{v_2} \left[\frac{1}{\omega(\theta)} \int_{v_1}^{v_2} \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\xi))}{\mathcal{F}(v_2) - \mathcal{F}(\xi)} \omega(\xi) \mathcal{F}'(\xi) \xi h_1(\xi) d\xi \right. \\
& \times \left. \int_{v_1}^{\theta} \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\varrho))}{\mathcal{F}(v_2) - \mathcal{F}(\varrho)} \omega(\varrho) \mathcal{F}'(\varrho) h_2(\varrho) d\varrho - \frac{1}{\omega(\theta)} \right. \\
& \times \left. \int_{v_1}^{v_2} \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\xi))}{\mathcal{F}(v_2) - \mathcal{F}(\xi)} \omega(\xi) \mathcal{F}'(\xi) h_1(\xi) d\xi \times \int_{v_1}^{\theta} \right. \\
& \times \left. \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\varrho))}{\mathcal{F}(v_2) - \mathcal{F}(\varrho)} \omega(\varrho) \mathcal{F}'(\varrho) \varrho h_2(\varrho) d\varrho \right] (\Phi'(\theta)) d\theta.
\end{aligned}$$

Thus with the aid of (7), the above equation gives,

$$\begin{aligned}
& \left[{}_{\omega}^{\mathcal{F}} \mathcal{I}_{v_1+}^{\Psi} h_1(v_2) \right] \left[{}_{\omega}^{\mathcal{F}} \mathcal{I}_{v_1+}^{\Psi} v_2(h_2 \Phi)(v_2) \right] + \left[{}_{\omega}^{\mathcal{F}} \mathcal{I}_{v_1+}^{\Psi} \Phi(v_2) \right] \left[{}_{\omega}^{\mathcal{F}} \mathcal{I}_{v_1+}^{\Psi} v_2(h_1 \Phi)(v_2) \right] \\
& - \left[{}_{\omega}^{\mathcal{F}} \mathcal{I}_{v_1+}^{\Psi} v_2 h_1(v_2) \right] \left[{}_{\omega}^{\mathcal{F}} \mathcal{I}_{v_1+}^{\Psi} (h_2 \Phi)(v_2) \right] - \left[{}_{\omega}^{\mathcal{F}} \mathcal{I}_{v_1+}^{\Psi} v_2 h_2(v_2) \right] \left[{}_{\omega}^{\mathcal{F}} \mathcal{I}_{v_1+}^{\Psi} (h_1 \Phi)(v_2) \right] \\
& = \frac{1}{\omega(\theta)} \int_{v_1}^{v_2} \left[{}_{\omega}^{\mathcal{F}} \mathcal{I}_{v_1+}^{\Psi} v_2 h_1(v_2) \int_{v_1}^{\theta} \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\varrho))}{\mathcal{F}(v_2) - \mathcal{F}(\varrho)} \omega(\varrho) \mathcal{F}'(\varrho) \varrho h_2(\varrho) d\varrho \right. \\
& \left. - {}_{\omega}^{\mathcal{F}} \mathcal{I}_{v_1+}^{\Psi} h_1(v_2) \int_{x_1}^{\theta} \varrho \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\varrho))}{\mathcal{F}(v_2) - \mathcal{F}(\varrho)} \omega(\varrho) \mathcal{F}'(\varrho) \varrho h_2(\varrho) d\varrho \right] (\Phi'(\theta)) d\theta,
\end{aligned}$$

which completes the proof. \square

Based on Lemma 1, we prove the following theorem.

Theorem 4. Suppose that the function \mathcal{F} be measurable, increasing, positive and monotone function on (v_1, v_2) , and has a continuous derivative \mathcal{F}' on (v_1, v_2) . Assume that $\Phi : [v_1, v_2] \rightarrow \mathbb{R}$ is absolutely continuous on $[v_1, v_2]$, and $\hbar_1, \hbar_2 : [v_1, v_2] \rightarrow \mathbb{R}^+$ are positive integrable functions. If $(\Phi')^2 \in L_1[v_1, v_2]$. Then, we have

$$\begin{aligned} & \left[{}_{\omega}^{\mathcal{F}} \mathcal{I}_{v_1+}^{\Psi} \hbar_1(v_2) \right] \left[{}_{\omega}^{\mathcal{F}} \mathcal{I}_{v_1+}^{\Psi} (\hbar_2 \Phi^2)(v_2) \right] + \left[{}_{\omega}^{\mathcal{F}} \mathcal{I}_{v_1+}^{\Psi} \hbar_2(v_2) \right] \left[{}_{\omega}^{\mathcal{F}} \mathcal{I}_{v_1+}^{\Psi} (\hbar_1 \Phi^2)(v_2) \right] \\ & - 2 \left[{}_{\omega}^{\mathcal{F}} \mathcal{I}_{v_1+}^{\Psi} (\hbar_1 \Phi)(v_2) \right] \left[{}_{\omega}^{\mathcal{F}} \mathcal{I}_{v_1+}^{\Psi} (\hbar_2 \Phi)(v_2) \right] \\ & \leq \frac{1}{\omega(\theta)} \int_{v_1}^{v_2} \left[{}_{\omega}^{\mathcal{F}} \mathcal{I}_{v_1+}^{\Psi} v_2 \hbar_1(v_2) \int_{v_1}^{\theta} \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\varrho))}{\mathcal{F}(v_2) - \mathcal{F}(\varrho)} \omega(\varrho) \mathcal{F}'(\varrho) \hbar_2(\varrho) d\varrho \right. \\ & \quad \left. - {}_{\omega}^{\mathcal{F}} \mathcal{I}_{v_1+}^{\Psi} \hbar_1(v_2) \int_{v_1}^{\theta} \varrho \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\varrho))}{\mathcal{F}(v_2) - \mathcal{F}(\varrho)} \omega(\varrho) \mathcal{F}'(\varrho) \hbar_2(\varrho) d\varrho \right] (\Phi'(\theta))^2 d\theta. \end{aligned} \quad (12)$$

Proof. By employing the definition (7) and Lemma 1, we obtain

$$\begin{aligned} & \left[{}_{\omega}^{\mathcal{F}} \mathcal{I}_{v_1+}^{\Psi} \hbar_1(v_2) \right] \left[{}_{\omega}^{\mathcal{F}} \mathcal{I}_{v_1+}^{\Psi} (\hbar_2 \Phi^2)(v_2) \right] + \left[{}_{\omega}^{\mathcal{F}} \mathcal{I}_{v_1+}^{\Psi} \hbar_2(v_2) \right] \left[{}_{\omega}^{\mathcal{F}} \mathcal{I}_{v_1+}^{\Psi} (\hbar_1 \Phi^2)(v_2) \right] \\ & - 2 \left[{}_{x_1}^{\Psi} \mathcal{I}_{\kappa}^{\theta} (\hbar_1 \Phi)(x_2) \right] \left[{}_{x_1}^{\Psi} \mathcal{I}_{\kappa}^{\theta} (\hbar_2 \Phi)(x_2) \right] \\ & = \frac{1}{\omega^2(\theta)} \int_{v_1}^{v_2} \int_{v_1}^{v_2} \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\xi))}{\mathcal{F}(v_2) - \mathcal{F}(\xi)} \\ & \times \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\varrho))}{\mathcal{F}(v_2) - \mathcal{F}(\varrho)} \omega(\xi) \mathcal{F}'(\xi) \hbar_1(\xi) \omega(\varrho) \mathcal{F}'(\varrho) \hbar_2(\varrho) (\Phi(\xi) - \Phi(\varrho))^2 d\xi d\varrho \\ & = \frac{1}{\omega^2(\theta)} \int_{v_1}^{v_2} \int_{v_1}^{v_2} \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\xi))}{\mathcal{F}(v_2) - \mathcal{F}(\xi)} \\ & \times \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\varrho))}{\mathcal{F}(v_2) - \mathcal{F}(\varrho)} \omega(\xi) \mathcal{F}'(\xi) \hbar_1(\xi) \omega(\varrho) \mathcal{F}'(\varrho) \hbar_2(\varrho) (\xi - \varrho)^2 \left(\frac{\Phi(\xi) - \Phi(\varrho)}{\xi - \varrho} \right)^2 d\xi d\varrho. \end{aligned}$$

Consequently, it follows that

$$\begin{aligned} & \left[{}_{\omega}^{\mathcal{F}} \mathcal{I}_{v_1+}^{\Psi} \hbar_1(v_2) \right] \left[{}_{\omega}^{\mathcal{F}} \mathcal{I}_{v_1+}^{\Psi} (\hbar_2 \Phi^2)(v_2) \right] + \left[{}_{\omega}^{\mathcal{F}} \mathcal{I}_{v_1+}^{\Psi} \hbar_2(v_2) \right] \left[{}_{\omega}^{\mathcal{F}} \mathcal{I}_{v_1+}^{\Psi} (\hbar_1 \Phi^2)(v_2) \right] \\ & - 2 \left[{}_{x_1}^{\Psi} \mathcal{I}_{\kappa}^{\theta} (\hbar_1 \Phi)(x_2) \right] \left[{}_{x_1}^{\Psi} \mathcal{I}_{\kappa}^{\theta} (\hbar_2 \Phi)(x_2) \right] \\ & = \frac{1}{\omega^2(\theta)} \int_{v_1}^{v_2} \int_{v_1}^{v_2} \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\xi))}{\mathcal{F}(v_2) - \mathcal{F}(\xi)} \\ & \times \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\varrho))}{\mathcal{F}(v_2) - \mathcal{F}(\varrho)} \omega(\xi) \mathcal{F}'(\xi) \hbar_1(\xi) \omega(\varrho) \mathcal{F}'(\varrho) \hbar_2(\varrho) (\xi - \varrho)^2 \left(\frac{\int_{\varrho}^{\xi} \Phi'(\theta) d\theta}{\xi - \varrho} \right)^2 d\xi d\varrho. \end{aligned}$$

By applying Cauchy-Schwartz inequality [40], we get

$$\begin{aligned}
& \left[{}_{\omega}^{\mathcal{F}} \mathcal{I}_{v_1+}^{\Psi} \hbar_1(v_2) \right] \left[{}_{\omega}^{\mathcal{F}} \mathcal{I}_{v_1+}^{\Psi} (\hbar_2 \Phi^2)(v_2) \right] + \left[{}_{\omega}^{\mathcal{F}} \mathcal{I}_{v_1+}^{\Psi} \hbar_2(v_2) \right] \left[{}_{\omega}^{\mathcal{F}} \mathcal{I}_{v_1+}^{\Psi} (\hbar_1 \Phi^2)(v_2) \right] \\
& - 2 \left[{}_{x_1}^{\Psi} \mathcal{I}_{\kappa}^{\theta} (\hbar_1 \Phi)(x_2) \right] \left[{}_{x_1}^{\Psi} \mathcal{I}_{\kappa}^{\theta} (\hbar_2 \Phi)(x_2) \right] \\
& = \frac{1}{\omega^2(\theta)} \int_{v_1}^{v_2} \int_{v_1}^{v_2} \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\xi))}{\mathcal{F}(v_2) - \mathcal{F}(\xi)} \\
& \times \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\varrho))}{\mathcal{F}(v_2) - \mathcal{F}(\varrho)} \omega(\xi) \mathcal{F}'(\xi) \hbar_1(\xi) \omega(\varrho) \mathcal{F}'(\varrho) \hbar_2(\varrho) (\xi - \varrho)^2 \left(\frac{\int_{\varrho}^{\xi} \Phi'(\theta) d\theta}{\xi - \varrho} \right)^2 d\xi d\varrho \\
& \leq \frac{1}{\omega^2(\theta)} \int_{v_1}^{v_2} \int_{v_1}^{v_2} \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\xi))}{\mathcal{F}(v_2) - \mathcal{F}(\xi)} \\
& \times \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\varrho))}{\mathcal{F}(v_2) - \mathcal{F}(\varrho)} \omega(\xi) \mathcal{F}'(\xi) \hbar_1(\xi) \omega(\varrho) \mathcal{F}'(\varrho) \hbar_2(\varrho) (\xi - \varrho)^2 \left(\frac{\left(\int_{\varrho}^{\xi} d\theta \right)^{\frac{1}{2}} \left(\int_{\varrho}^{\xi} (\Phi'(\theta))^2 d\theta \right)^{\frac{1}{2}}}{\xi - \varrho} \right)^2 d\xi d\varrho \\
& \leq \frac{1}{\omega^2(\theta)} \int_{v_1}^{v_2} \int_{v_1}^{v_2} \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\xi))}{\mathcal{F}(v_2) - \mathcal{F}(\xi)} \\
& \times \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\varrho))}{\mathcal{F}(v_2) - \mathcal{F}(\varrho)} \omega(\xi) \mathcal{F}'(\xi) \hbar_1(\xi) \omega(\varrho) \mathcal{F}'(\varrho) \hbar_2(\varrho) (\xi - \varrho)^2 \left(\frac{\left(\int_{\varrho}^{\xi} (\Phi'(\theta))^2 d\theta \right)}{\xi - \varrho} \right) d\xi d\varrho \\
& \leq \frac{1}{\omega^2(\theta)} \int_{v_1}^{v_2} \int_{v_1}^{v_2} \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\xi))}{\mathcal{F}(v_2) - \mathcal{F}(\xi)} \\
& \times \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\varrho))}{\mathcal{F}(v_2) - \mathcal{F}(\varrho)} \omega(\xi) \mathcal{F}'(\xi) \hbar_1(\xi) \omega(\varrho) \mathcal{F}'(\varrho) \hbar_2(\varrho) (\xi - \varrho) \left(\int_{\varrho}^{\xi} (\Phi'(\theta))^2 d\theta \right) d\xi d\varrho
\end{aligned} \tag{13}$$

Hence, using (11) and (13) concludes the proof. \square

The following new particular results of Theorem 4 can be easily obtained.

Corollary 1. Suppose that the function \mathcal{F} be measurable, increasing, positive and monotone function on (v_1, v_2) , and has a continuous derivative \mathcal{F}' on (v_1, v_2) . Assume that $\Phi : [v_1, v_2] \rightarrow \mathbb{R}$ is absolutely continuous on $[v_1, v_2]$, and $\hbar_1, \hbar_2 : [v_1, v_2] \rightarrow \mathbb{R}^+$ are positive integrable functions. If $(\Phi')^2 \in L_1[v_1, v_2]$. Then, we have

$$\begin{aligned}
& \left[{}_{\omega}^{\mathcal{F}} \mathcal{I}_{v_1+}^{\Psi} 1 \right] \left[{}_{\omega}^{\mathcal{F}} \mathcal{I}_{v_1+}^{\Psi} (\hbar_2 \Phi^2)(v_2) \right] + \left[{}_{\omega}^{\mathcal{F}} \mathcal{I}_{v_1+}^{\Psi} \hbar_2(v_2) \right] \left[{}_{\omega}^{\mathcal{F}} \mathcal{I}_{v_1+}^{\Psi} (\Phi^2)(v_2) \right] \\
& - 2 \left[{}_{\omega}^{\mathcal{F}} \mathcal{I}_{v_1+}^{\Psi} (\Phi)(v_2) \right] \left[{}_{\omega}^{\mathcal{F}} \mathcal{I}_{v_1+}^{\Psi} (\hbar_2 \Phi)(v_2) \right] \\
& \leq \frac{1}{\omega(\theta)} \int_{v_1}^{v_2} \left[{}_{\omega}^{\mathcal{F}} \mathcal{I}_{v_1+}^{\Psi} v_2 \int_{v_1}^{\theta} \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\varrho))}{\mathcal{F}(v_2) - \mathcal{F}(\varrho)} \omega(\varrho) \mathcal{F}'(\varrho) \hbar_2(\varrho) d\varrho \right. \\
& \quad \left. - {}_{\omega}^{\mathcal{F}} \mathcal{I}_{v_1+}^{\Psi} 1 \int_{v_1}^{\theta} \varrho \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\varrho))}{\mathcal{F}(v_2) - \mathcal{F}(\varrho)} \omega(\varrho) \mathcal{F}'(\varrho) \hbar_2(\varrho) d\varrho \right] (\Phi'(\theta))^2 d\theta.
\end{aligned}$$

Proof. By considering $\hbar_1(\theta) = 1, \theta \in [v_1, v_2]$ in Theorem 4, the desired result is obtained. \square

Corollary 2. Suppose that the function \mathcal{F} be measurable, increasing, positive and monotone function on (v_1, v_2) , and has a continuous derivative \mathcal{F}' on (v_1, v_2) . Assume that $\Phi : [v_1, v_2] \rightarrow \mathbb{R}$ is absolutely continuous on $[v_1, v_2]$, and $\hbar_1, \hbar_2 : [v_1, v_2] \rightarrow \mathbb{R}^+$ are positive integrable functions. If $(\Phi')^2 \in L_1[v_1, v_2]$. Then, we have

$$\begin{aligned}
& \left[\frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^{\Psi} \hbar_1(v_2) \right] \left[\frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^{\Psi} (\Phi^2)(v_2) \right] + \left[\frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^{\Psi} 1 \right] \left[\frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^{\Psi} (\hbar_1 \Phi^2)(v_2) \right] \\
& - 2 \left[\frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^{\Psi} (\hbar_1 \Phi)(v_2) \right] \left[\frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^{\Psi} (\Phi)(v_2) \right] \\
& \leq \frac{1}{\omega(\theta)} \int_{v_1}^{v_2} \left[\frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^{\Psi} v_2 \hbar_1(v_2) \int_{v_1}^{\theta} \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\varrho))}{\mathcal{F}(v_2) - \mathcal{F}(\varrho)} \omega(\varrho) \mathcal{F}'(\varrho) d\varrho \right. \\
& \quad \left. - \frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^{\Psi} \hbar_1(v_2) \int_{v_1}^{\theta} \varrho \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\varrho))}{\mathcal{F}(v_2) - \mathcal{F}(\varrho)} \omega(\varrho) \mathcal{F}'(\varrho) d\varrho \right] (\Phi'(\theta))^2 d\theta.
\end{aligned}$$

Proof. By considering $\hbar_2(\theta) = 1$, $\theta \in [v_1, v_2]$ in Theorem 4, desired corollary is proven. \square

Corollary 3. Suppose that the function \mathcal{F} be measurable, increasing, positive and monotone function on (v_1, v_2) , and has a continuous derivative \mathcal{F}' on (v_1, v_2) . Assume that $\Phi : [v_1, v_2] \rightarrow \mathbb{R}$ is absolutely continuous on $[v_1, v_2]$, and $\hbar_1, \hbar_2 : [v_1, v_2] \rightarrow \mathbb{R}^+$ are positive integrable functions. If $(\Phi')^2 \in L_1[v_1, v_2]$. Then, we have

$$\begin{aligned}
& \left[\frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^{\Psi} 1 \right] \left[\frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^{\Psi} (\Phi^2)(v_2) \right] - \left[\frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^{\Psi} (\Phi)(v_2) \right]^2 \\
& \leq \frac{1}{\omega(\theta)} \int_{v_1}^{v_2} \left[\frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^{\Psi} v_2 \int_{v_1}^{\theta} \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\varrho))}{\mathcal{F}(v_2) - \mathcal{F}(\varrho)} \omega(\varrho) \mathcal{F}'(\varrho) d\varrho \right. \\
& \quad \left. - \frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^{\Psi} 1 \int_{v_1}^{\theta} \varrho \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\varrho))}{\mathcal{F}(v_2) - \mathcal{F}(\varrho)} \omega(\varrho) \mathcal{F}'(\varrho) d\varrho \right] (\Phi'(\theta))^2 d\theta.
\end{aligned}$$

Proof. Taking $\hbar_1(\theta) = \hbar_2(\theta) = 1$, $\theta \in [v_1, v_2]$ in Theorem 4, the desired result is obtained. \square

Remark 4. If we consider $\omega(\theta) = 1$ and $\Psi(\mathcal{F}(\theta)) = \frac{\mathcal{F}(\theta)^{\kappa}}{\Gamma(\kappa)}$, then Theorem 4 and Corollaries 1–3 will reduce to the work of Bezziou et al. [27].

Remark 5. If we consider $\omega(\theta) = 1$ and $\Psi(\mathcal{F}(\theta)) = \frac{\mathcal{F}(\theta)^{\frac{\kappa}{\eta}}}{\Gamma_{\eta}(\kappa)}$, then Theorem 4 and Corollaries 1–3 will reduce to the work of Rahman et al. [41].

Theorem 5. Let \mathcal{F} be measurable, increasing, positive and monotone function on (v_1, v_2) , and having a continuous derivative \mathcal{F}' on (v_1, v_2) . Assume that $f_1, f_2 : [v_1, v_2] \rightarrow \mathbb{R}$ are absolutely continuous on $[v_1, v_2]$, and $\hbar_1, \hbar_2 : [v_1, v_2] \rightarrow \mathbb{R}^+$ are positive integrable. If $(f'_1)^2 \in L_1[v_1, v_2]$ and $(f'_2)^2 \in L_1[v_1, v_2]$. Then, we have

$$\begin{aligned}
& \left| \frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^{\Psi} \hbar_1(v_2) \frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^{\Psi} (\hbar_2 f_1 f_2)(v_2) + \frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^{\Psi} \hbar_2(v_2) \frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^{\Psi} (\hbar_1 f_1 f_2)(v_2) \right. \\
& \quad \left. - \frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^{\Psi} (\hbar_2 f_1)(v_2) \frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^{\Psi} (\hbar_1 f_2)(v_2) - \frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^{\Psi} (\hbar_1 f_1)(v_2) \frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^{\Psi} (\hbar_2 f_2)(v_2) \right| \\
& \leq \frac{1}{\omega(\theta)} \left(\int_{v_1}^{v_2} \left[\frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^{\Psi} v_2 \hbar_1(v_2) \int_{v_1}^{\theta} \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\varrho))}{\mathcal{F}(v_2) - \mathcal{F}(\varrho)} \omega(\varrho) \mathcal{F}'(\varrho) \hbar_2(\varrho) d\varrho \right. \right. \\
& \quad \left. \left. - \frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^{\Psi} \hbar_1(v_2) \int_{v_1}^{\theta} \varrho \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\varrho))}{\mathcal{F}(v_2) - \mathcal{F}(\varrho)} \omega(\varrho) \mathcal{F}'(\varrho) \hbar_2(\varrho) d\varrho \right] (f'_1(\theta))^2 d\theta \right)^{\frac{1}{2}} \\
& \quad \times \left(\int_{v_1}^{v_2} \left[\frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^{\Psi} v_2 \hbar_2(v_2) \int_{v_1}^{\theta} \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\varrho))}{\mathcal{F}(v_2) - \mathcal{F}(\varrho)} \omega(\varrho) \mathcal{F}'(\varrho) \hbar_1(\varrho) d\varrho \right. \right. \\
& \quad \left. \left. - \frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^{\Psi} \hbar_2(v_2) \int_{v_1}^{\theta} \varrho \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\varrho))}{\mathcal{F}(v_2) - \mathcal{F}(\varrho)} \omega(\varrho) \mathcal{F}'(\varrho) \hbar_1(\varrho) d\varrho \right] (f'_2(\theta))^2 d\theta \right)^{\frac{1}{2}}
\end{aligned} \tag{14}$$

Proof. Consider the left-hand side of (14), we have

$$\begin{aligned}
& \left| {}_{\omega}^{\mathcal{F}} \mathcal{I}_{v_1+}^{\Psi} h_2(v_2) {}_{\omega}^{\mathcal{F}} \mathcal{I}_{v_1+}^{\Psi} (h_2 f_1 f_2)(v_2) + {}_{\omega}^{\mathcal{F}} \mathcal{I}_{v_1+}^{\Psi} h_2(v_2) {}_{\omega}^{\mathcal{F}} \mathcal{I}_{v_1+}^{\Psi} (h_1 f_1 f_2)(v_2) \right. \\
& - \left. {}_{\omega}^{\mathcal{F}} \mathcal{I}_{v_1+}^{\Psi} (h_2 f_1)(v_2) {}_{\omega}^{\mathcal{F}} \mathcal{I}_{v_1+}^{\Psi} (h_1 f_2)(v_2) - {}_{\omega}^{\mathcal{F}} \mathcal{I}_{v_1+}^{\Psi} (h_1 f_1)(v_2) {}_{\omega}^{\mathcal{F}} \mathcal{I}_{v_1+}^{\Psi} (h_2 f_2)(v_2) \right| \\
& \leq \frac{1}{\omega^2(\theta)} \left(\int_{v_1}^{v_2} \int_{v_1}^{v_2} \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\xi))}{\mathcal{F}(v_2) - \mathcal{F}(\xi)} \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\varrho))}{\mathcal{F}(v_2) - \mathcal{F}(\varrho)} \right. \\
& \times \omega(\xi) \mathcal{F}'(\xi) \omega(\varrho) \mathcal{F}'(\varrho) h_1(\xi) h_2(\varrho) (f_1(\xi) - f_1(\varrho))^2 d\xi d\varrho \left. \right)^{\frac{1}{2}} \times \left(\int_{v_1}^{v_2} \int_{v_1}^{v_2} \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\xi))}{\mathcal{F}(v_2) - \mathcal{F}(\xi)} \right. \\
& \times \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\varrho))}{\mathcal{F}(v_2) - \mathcal{F}(\varrho)} \omega(\xi) \mathcal{F}'(\xi) \omega(\varrho) \mathcal{F}'(\varrho) h_1(\xi) h_2(\varrho) (f_2(\xi) - f_2(\varrho))^2 d\xi d\varrho \left. \right)^{\frac{1}{2}} \\
& \leq \frac{1}{\omega^2(\theta)} \left(\int_{v_1}^{v_2} \int_{v_1}^{v_2} \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\xi))}{\mathcal{F}(v_2) - \mathcal{F}(\xi)} \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\varrho))}{\mathcal{F}(v_2) - \mathcal{F}(\varrho)} \omega(\xi) \mathcal{F}'(\xi) \omega(\varrho) \mathcal{F}'(\varrho) h_1(\xi) h_2(\varrho) \right. \\
& \times \left(\int_{\varrho}^{\xi} f'_1(\theta) d\theta \right)^2 d\xi d\varrho \left. \right)^{\frac{1}{2}} \times \left(\int_{v_1}^{v_2} \int_{v_1}^{v_2} \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\xi))}{\mathcal{F}(v_2) - \mathcal{F}(\xi)} \right. \\
& \times \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\varrho))}{\mathcal{F}(v_2) - \mathcal{F}(\varrho)} \omega(\xi) \mathcal{F}'(\xi) \omega(\varrho) \mathcal{F}'(\varrho) h_1(\xi) h_2(\varrho) \left(\int_{\varrho}^{\xi} f'_2(\theta) d\theta \right)^2 d\xi d\varrho \left. \right)^{\frac{1}{2}}
\end{aligned}$$

Applying Cauchy-Schwartz inequality [40] to the above equation yields,

$$\begin{aligned}
& \left| {}_{\omega}^{\mathcal{F}} \mathcal{I}_{v_1+}^{\Psi} h_2(v_2) {}_{\omega}^{\mathcal{F}} \mathcal{I}_{v_1+}^{\Psi} (h_2 f_1 f_2)(v_2) + {}_{\omega}^{\mathcal{F}} \mathcal{I}_{v_1+}^{\Psi} h_2(v_2) {}_{\omega}^{\mathcal{F}} \mathcal{I}_{v_1+}^{\Psi} (h_1 f_1 f_2)(v_2) \right. \\
& - \left. {}_{\omega}^{\mathcal{F}} \mathcal{I}_{v_1+}^{\Psi} (h_2 f_1)(v_2) {}_{\omega}^{\mathcal{F}} \mathcal{I}_{v_1+}^{\Psi} (h_1 f_2)(v_2) - {}_{\omega}^{\mathcal{F}} \mathcal{I}_{v_1+}^{\Psi} (h_1 f_1)(v_2) {}_{\omega}^{\mathcal{F}} \mathcal{I}_{v_1+}^{\Psi} (h_2 f_2)(v_2) \right| \\
& \leq \frac{1}{\omega^2(\theta)} \left\{ \int_{v_1}^{v_2} \int_{v_1}^{v_2} \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\xi))}{\mathcal{F}(v_2) - \mathcal{F}(\xi)} \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\varrho))}{\mathcal{F}(v_2) - \mathcal{F}(\varrho)} \omega(\xi) \mathcal{F}'(\xi) \omega(\varrho) \mathcal{F}'(\varrho) h_1(\xi) h_2(\varrho) \right. \\
& \times \left(\left(\int_{\varrho}^{\xi} d\theta \right)^{\frac{1}{2}} \left(\int_{\varrho}^{\xi} (f'_1(\theta))^2 d\theta \right)^{\frac{1}{2}} \right)^2 d\xi d\varrho \left. \right\}^{\frac{1}{2}} \times \left\{ \int_{v_1}^{v_2} \int_{v_1}^{v_2} \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\xi))}{\mathcal{F}(v_2) - \mathcal{F}(\xi)} \right. \\
& \times \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\varrho))}{\mathcal{F}(v_2) - \mathcal{F}(\varrho)} \omega(\xi) \mathcal{F}'(\xi) \omega(\varrho) \mathcal{F}'(\varrho) h_1(\xi) h_2(\varrho) \left(\left(\int_{\varrho}^{\xi} d\theta \right)^{\frac{1}{2}} \left(\int_{\varrho}^{\xi} (f'_2(\theta))^2 d\theta \right)^{\frac{1}{2}} \right)^2 d\xi d\varrho \left. \right\}^{\frac{1}{2}} \\
& \leq \frac{1}{\omega^2(\theta)} \left\{ \int_{v_1}^{v_2} \int_{v_1}^{v_2} \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\xi))}{\mathcal{F}(v_2) - \mathcal{F}(\xi)} \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\varrho))}{\mathcal{F}(v_2) - \mathcal{F}(\varrho)} \omega(\xi) \mathcal{F}'(\xi) \omega(\varrho) \mathcal{F}'(\varrho) h_1(\xi) h_2(\varrho) \right. \\
& \times \omega(\varrho) \mathcal{F}'(\varrho) h_1(\xi) h_2(\varrho) (\xi - \varrho) \left(\int_{\varrho}^{\xi} (f'_1(\theta))^2 d\theta \right) d\xi d\varrho \left. \right\}^{\frac{1}{2}} \times \left\{ \int_{v_1}^{v_2} \int_{v_1}^{v_2} \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\xi))}{\mathcal{F}(v_2) - \mathcal{F}(\xi)} \right. \\
& \times \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\varrho))}{\mathcal{F}(v_2) - \mathcal{F}(\varrho)} \omega(\xi) \mathcal{F}'(\xi) \omega(\varrho) \mathcal{F}'(\varrho) h_1(\xi) h_2(\varrho) (\xi - \varrho) \left(\int_{\varrho}^{\xi} (f'_2(\theta))^2 d\theta \right) d\xi d\varrho \left. \right\}^{\frac{1}{2}} \\
& \leq \frac{1}{\omega^2(\theta)} \left\{ \int_{v_1}^{v_2} \left(\int_{v_1}^{v_2} \xi \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\xi))}{\mathcal{F}(v_2) - \mathcal{F}(\xi)} \omega(\xi) \mathcal{F}'(\xi) h_1(\xi) d\xi \right. \right. \\
& \times \int_{v_1}^{\theta} \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\varrho))}{\mathcal{F}(v_2) - \mathcal{F}(\varrho)} \omega(\varrho) \mathcal{F}'(\varrho) h_2(\varrho) d\varrho \\
& - \int_{v_1}^{v_2} \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\xi))}{\mathcal{F}(v_2) - \mathcal{F}(\xi)} \omega(\xi) \mathcal{F}'(\xi) h_1(\xi) d\xi \\
& \times \int_{x_1}^{\theta} \varrho \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\varrho))}{\mathcal{F}(v_2) - \mathcal{F}(\varrho)} \omega(\varrho) \mathcal{F}'(\varrho) h_2(\varrho) d\varrho \left(f'_1(\theta) \right)^2 \left. \right)^{\frac{1}{2}} \\
& - \left\{ \int_{v_1}^{v_2} \left(\int_{v_1}^{v_2} \xi \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\xi))}{\mathcal{F}(v_2) - \mathcal{F}(\xi)} \omega(\xi) \mathcal{F}'(\xi) h_1(\xi) d\xi \right. \right. \\
& \times \int_{v_1}^{\theta} \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\varrho))}{\mathcal{F}(v_2) - \mathcal{F}(\varrho)} \omega(\varrho) \mathcal{F}'(\varrho) h_2(\varrho) d\varrho \\
& - \int_{v_1}^{v_2} \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\xi))}{\mathcal{F}(v_2) - \mathcal{F}(\xi)} \omega(\xi) \mathcal{F}'(\xi) h_1(\xi) d\xi \\
& \times \int_{x_1}^{\theta} \varrho \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\varrho))}{\mathcal{F}(v_2) - \mathcal{F}(\varrho)} \omega(\varrho) \mathcal{F}'(\varrho) h_2(\varrho) d\varrho \left(f'_2(\theta) \right)^2 \left. \right]^{\frac{1}{2}}.
\end{aligned}$$

In view of (7), we get the desired proof of (14). \square

Corollary 4. Let \mathcal{F} be measurable, increasing, positive and monotone function on (v_1, v_2) , and having a continuous derivative \mathcal{F}' on (v_1, v_2) . Assume that $f_1, f_2 : [v_1, v_2] \rightarrow \mathbb{R}$ are absolutely continuous on $[v_1, v_2]$, and $\hbar_1, \hbar_2 : [v_1, v_2] \rightarrow \mathbb{R}^+$ are positive integrable. If $(f'_1)^2 \in L_1[v_1, v_2]$ and $(f'_2)^2 \in L_1[v_1, v_2]$. Then, we have

$$\begin{aligned} & \left| \frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^\Psi(1) \frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^\Psi(\hbar_2 f_1 f_2)(v_2) + \frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^\Psi \hbar_2(v_2) \frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^\Psi(f_1 f_2)(v_2) \right. \\ & \quad \left. - \frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^\Psi(\hbar_2 f_1)(v_2) \frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^\Psi(f_2)(v_2) - \frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^\Psi(f_1)(v_2) \frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^\Psi(\hbar_2 f_2)(v_2) \right| \\ & \leq \frac{1}{\omega(\theta)} \left(\int_{v_1}^{v_2} \left[\frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^\Psi v_2 \int_{v_1}^\theta \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\varrho))}{\mathcal{F}(v_2) - \mathcal{F}(\varrho)} \omega(\varrho) \mathcal{F}'(\varrho) \hbar_2(\varrho) d\varrho \right. \right. \\ & \quad \left. \left. - \frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^\Psi(1) \int_{v_1}^\theta \varrho \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\varrho))}{\mathcal{F}(v_2) - \mathcal{F}(\varrho)} \omega(\varrho) \mathcal{F}'(\varrho) \hbar_2(\varrho) d\varrho \right] (f'_1(\theta))^2 d\theta \right)^{\frac{1}{2}} \\ & \quad \times \left(\int_{v_1}^{v_2} \left[\frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^\Psi v_2 \int_{v_1}^\theta \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\varrho))}{\mathcal{F}(v_2) - \mathcal{F}(\varrho)} \omega(\varrho) \mathcal{F}'(\varrho) \hbar_2(\varrho) d\varrho \right. \right. \\ & \quad \left. \left. - \frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^\Psi(1) \int_{v_1}^\theta \varrho \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\varrho))}{\mathcal{F}(v_2) - \mathcal{F}(\varrho)} \omega(\varrho) \mathcal{F}'(\varrho) \hbar_2(\varrho) d\varrho \right] (f'_2(\theta))^2 d\theta \right)^{\frac{1}{2}} \end{aligned}$$

Proof. Applying Theorem 5 for $\hbar_1(\theta) = 1, \theta \in [v_1, v_2]$, the desired result is obtained. \square

Corollary 5. Let \mathcal{F} be measurable, increasing, positive and monotone function on (v_1, v_2) , and having a continuous derivative \mathcal{F}' on (v_1, v_2) . Assume that $f_1, f_2 : [v_1, v_2] \rightarrow \mathbb{R}$ are absolutely continuous on $[v_1, v_2]$, and $\hbar_1, \hbar_2 : [v_1, v_2] \rightarrow \mathbb{R}^+$ are positive integrable. If $(f'_1)^2 \in L_1[v_1, v_2]$ and $(f'_2)^2 \in L_1[v_1, v_2]$. Then, we have

$$\begin{aligned} & \left| \frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^\Psi \hbar_1(v_2) \frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^\Psi(f_1 f_2)(v_2) + \frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^\Psi(1) \frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^\Psi(\hbar_1 f_1 f_2)(v_2) \right. \\ & \quad \left. - \frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^\Psi(f_1)(v_2) \frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^\Psi(\hbar_1 f_2)(v_2) - \frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^\Psi(\hbar_1 f_1)(v_2) \frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^\Psi(f_2)(v_2) \right| \\ & \leq \frac{1}{\omega(\theta)} \left(\int_{v_1}^{v_2} \left[\frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^\Psi v_2 \hbar_1(v_2) \int_{v_1}^\theta \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\varrho))}{\mathcal{F}(v_2) - \mathcal{F}(\varrho)} \omega(\varrho) \mathcal{F}'(\varrho) d\varrho \right. \right. \\ & \quad \left. \left. - \frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^\Psi \hbar_1(v_2) \int_{v_1}^\theta \varrho \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\varrho))}{\mathcal{F}(v_2) - \mathcal{F}(\varrho)} \omega(\varrho) \mathcal{F}'(\varrho) d\varrho \right] (f'_1(\theta))^2 d\theta \right)^{\frac{1}{2}} \\ & \quad \times \left(\int_{v_1}^{v_2} \left[\frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^\Psi v_2 \hbar_1(v_2) \int_{v_1}^\theta \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\varrho))}{\mathcal{F}(v_2) - \mathcal{F}(\varrho)} \omega(\varrho) \mathcal{F}'(\varrho) d\varrho \right. \right. \\ & \quad \left. \left. - \frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^\Psi \hbar_1(v_2) \int_{v_1}^\theta \varrho \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\varrho))}{\mathcal{F}(v_2) - \mathcal{F}(\varrho)} \omega(\varrho) \mathcal{F}'(\varrho) d\varrho \right] (f'_2(\theta))^2 d\theta \right)^{\frac{1}{2}} \end{aligned}$$

Proof. Applying Theorem 5 for $\hbar_2(\theta) = 1, \theta \in [v_1, v_2]$, the desired result is obtained. \square

Corollary 6. Let \mathcal{F} be measurable, increasing, positive and monotone function on (v_1, v_2) , and having a continuous derivative \mathcal{F}' on (v_1, v_2) . Assume that $f_1, f_2 : [v_1, v_2] \rightarrow \mathbb{R}$ are absolutely continuous on $[v_1, v_2]$, and $\hbar_1, \hbar_2 : [v_1, v_2] \rightarrow \mathbb{R}^+$ are positive integrable. If $(f'_1)^2 \in L_1[v_1, v_2]$ and $(f'_2)^2 \in L_1[v_1, v_2]$. Then, we have

$$\begin{aligned} & \left| \frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^\Psi(1) \frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^\Psi(f_1 f_2)(v_2) - \frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^\Psi(f_1)(v_2) \frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^\Psi(f_2)(v_2) \right| \\ & \leq \frac{1}{\omega(\theta)} \left(\int_{v_1}^{v_2} \left[\frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^\Psi v_2 \int_{v_1}^\theta \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\varrho))}{\mathcal{F}(v_2) - \mathcal{F}(\varrho)} \omega(\varrho) \mathcal{F}'(\varrho) d\varrho \right. \right. \\ & \quad \left. \left. - \frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^\Psi(1) \int_{v_1}^\theta \varrho \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\varrho))}{\mathcal{F}(v_2) - \mathcal{F}(\varrho)} \omega(\varrho) \mathcal{F}'(\varrho) d\varrho \right] (f'_1(\theta))^2 d\theta \right)^{\frac{1}{2}} \\ & \quad \times \left(\int_{v_1}^{v_2} \left[\frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^\Psi v_2 \int_{v_1}^\theta \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\varrho))}{\mathcal{F}(v_2) - \mathcal{F}(\varrho)} \omega(\varrho) \mathcal{F}'(\varrho) d\varrho \right. \right. \\ & \quad \left. \left. - \frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^\Psi(1) \int_{v_1}^\theta \varrho \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\varrho))}{\mathcal{F}(v_2) - \mathcal{F}(\varrho)} \omega(\varrho) \mathcal{F}'(\varrho) d\varrho \right] (f'_2(\theta))^2 d\theta \right)^{\frac{1}{2}} \end{aligned}$$

Proof. Applying Theorem 5 for $\hbar_1(\theta) = \hbar_2(\theta) = 1$, $\theta \in [v_1, v_2]$, the desired result is obtained. \square

Theorem 6. Let \mathcal{F} be measurable, increasing, positive and monotone function on (v_1, v_2) , and having a continuous derivative \mathcal{F}' on (v_1, v_2) . Assume that $f_1 : [v_1, v_2] \rightarrow \mathbb{R}$ is absolutely continuous function on $[v_1, v_2]$, and $f_2 : [v_1, v_2] \rightarrow \mathbb{R}$ is non-decreasing on $[v_1, v_2]$. Moreover, suppose that both $\hbar_1, \hbar_2 : [v_1, v_2] \rightarrow \mathbb{R}^+$ are positive integrable. If $(f'_1) \in L^\infty[v_1, v_2]$, then we have

$$\begin{aligned} & \left| \frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^{\Psi} \hbar_1(v_2) \frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^{\Psi} (\hbar_2 f_1 f_2)(v_2) + \frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^{\Psi} \hbar_2(v_2) \frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^{\Psi} (\hbar_1 f_1 f_2)(v_2) \right. \\ & \quad \left. - \frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^{\Psi} (\hbar_1 f_1)(v_2) \frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^{\Psi} (\hbar_2 f_2)(v_2) - \frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^{\Psi} (\hbar_2 f_1)(v_2) \frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^{\Psi} (\hbar_1 f_2)(v_2) \right| \\ & \leq \frac{\|f'_1\|_\infty}{\omega(\theta)} \left[\frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^{\Psi} v_2 \hbar_1(v_2) \int_{v_1}^\theta \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\varrho))}{\mathcal{F}(v_2) - \mathcal{F}(\varrho)} \omega(\varrho) \mathcal{F}'(\varrho) \hbar_2(\varrho) d\varrho \right. \\ & \quad \left. - \frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^{\Psi} \hbar_1(v_2) \int_{v_1}^\theta \varrho \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\varrho))}{\mathcal{F}(v_2) - \mathcal{F}(\varrho)} \omega(\varrho) \mathcal{F}'(\varrho) \hbar_2(\varrho) d\varrho \right] \int_{v_1}^{v_2} f'_2(\theta) d\theta. \end{aligned} \quad (15)$$

Proof. Consider the left-hand side of (15), we have

$$\begin{aligned} & \left| \frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^{\Psi} \hbar_1(v_2) \frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^{\Psi} (\hbar_2 f_1 f_2)(v_2) + \frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^{\Psi} \hbar_2(v_2) \frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^{\Psi} (\hbar_1 f_1 f_2)(v_2) \right. \\ & \quad \left. - \frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^{\Psi} (\hbar_2 f_1)(v_2) \frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^{\Psi} (\hbar_1 f_2)(v_2) - \frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^{\Psi} (\hbar_1 f_1)(v_2) \frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^{\Psi} (\hbar_2 f_2)(v_2) \right| \\ & \leq \frac{1}{\omega^2(\theta)} \left| \int_{v_1}^{v_2} \int_{v_1}^{v_2} \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\xi))}{\mathcal{F}(v_2) - \mathcal{F}(\xi)} \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\varrho))}{\mathcal{F}(v_2) - \mathcal{F}(\varrho)} \omega(\xi) \mathcal{F}'(\xi) \omega(\varrho) \mathcal{F}'(\varrho) \hbar_1(\xi) \hbar_2(\varrho) \right. \\ & \quad \times [(f_1(\xi) - f_1(\varrho))(f_2(\xi) - f_2(\varrho))] d\xi d\varrho \left. \right| \\ & \leq \frac{1}{\omega^2(\theta)} \left| \int_{v_1}^{v_2} \int_{v_1}^{v_2} \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\xi))}{\mathcal{F}(v_2) - \mathcal{F}(\xi)} \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\varrho))}{\mathcal{F}(v_2) - \mathcal{F}(\varrho)} \omega(\xi) \mathcal{F}'(\xi) \omega(\varrho) \mathcal{F}'(\varrho) \hbar_1(\xi) \hbar_2(\varrho) \right. \\ & \quad \times \left| \frac{(f_1(\xi) - f_1(\varrho))}{\xi - \varrho} \right| |(\xi - \varrho)(f_2(\xi) - f_2(\varrho))| d\xi d\varrho \left. \right| \\ & \leq \frac{\|f'_1\|_\infty}{\omega^2(\theta)} \left| \int_{v_1}^{v_2} \int_{v_1}^{v_2} \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\xi))}{\mathcal{F}(v_2) - \mathcal{F}(\xi)} \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\varrho))}{\mathcal{F}(v_2) - \mathcal{F}(\varrho)} \omega(\xi) \mathcal{F}'(\xi) \omega(\varrho) \mathcal{F}'(\varrho) \hbar_1(\xi) \hbar_2(\varrho) \right. \\ & \quad \times (\xi - \varrho) \left(\int_{v_1}^{v_2} f'_2(\vartheta) d\vartheta \right) d\xi d\varrho \left. \right| \\ & \leq \frac{\|f'_1\|_\infty}{\omega^2(\theta)} \left[\int_{v_1}^{v_2} \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\xi))}{\mathcal{F}(v_2) - \mathcal{F}(\xi)} \xi \mathcal{F}'(\xi) \omega(\xi) \hbar_2(\xi) d\xi \int_{v_1}^\theta \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\xi))}{\mathcal{F}(v_2) - \mathcal{F}(\varrho)} \mathcal{F}'(\varrho) \omega(\varrho) \hbar_2(\varrho) d\varrho \right. \\ & \quad \left. - \int_{v_1}^{v_2} \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\xi))}{\mathcal{F}(v_2) - \mathcal{F}(\xi)} \mathcal{F}'(\xi) \omega(\varrho) \hbar_2(\xi) d\xi \int_{v_1}^\theta \varrho \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\xi))}{\mathcal{F}(v_2) - \mathcal{F}(\varrho)} \mathcal{F}'(\varrho) \omega(\varrho) \hbar_2(\varrho) d\varrho \right] \int_{v_1}^{v_2} f'_2(\theta) d\theta \end{aligned}$$

Hence taking (7) into account, the proof of (15) is completed. \square

Corollary 7. Let \mathcal{F} be measurable, increasing, positive and monotone function on (v_1, v_2) , and having a continuous derivative \mathcal{F}' on (v_1, v_2) . Assume that $f_1 : [v_1, v_2] \rightarrow \mathbb{R}$ is absolutely continuous function on $[v_1, v_2]$, and $f_2 : [v_1, v_2] \rightarrow \mathbb{R}$ is non-decreasing on $[v_1, v_2]$. Moreover, suppose that both $\hbar_1, \hbar_2 : [v_1, v_2] \rightarrow \mathbb{R}^+$ are positive integrable. If $(f'_1) \in L^\infty[v_1, v_2]$, then we have

$$\begin{aligned}
& \left| \mathcal{F} \mathcal{I}_{v_1+}^{\Psi} (1) \mathcal{F} \mathcal{I}_{v_1+}^{\Psi} (\hbar_2 f_1 f_2)(v_2) + \mathcal{F} \mathcal{I}_{v_1+}^{\Psi} \hbar_2(v_2) \mathcal{F} \mathcal{I}_{v_1+}^{\Psi} (f_1 f_2)(v_2) \right. \\
& - \left. \mathcal{F} \mathcal{I}_{v_1+}^{\Psi} (f_1)(v_2) \mathcal{F} \mathcal{I}_{v_1+}^{\Psi} (\hbar_2 f_2)(v_2) - \mathcal{F} \mathcal{I}_{v_1+}^{\Psi} (\hbar_2 f_1)(v_2) \mathcal{F} \mathcal{I}_{v_1+}^{\Psi} (f_2)(v_2) \right| \\
& \leq \frac{\|f'_1\|_{\infty}}{\omega(\theta)} \left[\mathcal{F} \mathcal{I}_{v_1+}^{\Psi} v_2 \int_{v_1}^{\theta} \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\varrho))}{\mathcal{F}(v_2) - \mathcal{F}(\varrho)} \omega(\varrho) \mathcal{F}'(\varrho) \hbar_2(\varrho) d\varrho \right. \\
& \left. - \mathcal{F} \mathcal{I}_{v_1+}^{\Psi} (1) \int_{v_1}^{\theta} \varrho \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\varrho))}{\mathcal{F}(v_2) - \mathcal{F}(\varrho)} \omega(\varrho) \mathcal{F}'(\varrho) \hbar_2(\varrho) d\varrho \right] \int_{v_1}^{v_2} f'_2(\theta) d\theta.
\end{aligned}$$

Proof. Applying Theorem 6 for $\hbar_1(\theta) = 1, \theta \in [v_1, v_2]$, the desired result is obtained. \square

Corollary 8. Let \mathcal{F} be measurable, increasing, positive and monotone function on (v_1, v_2) , and having a continuous derivative \mathcal{F}' on (v_1, v_2) . Assume that $f_1 : [v_1, v_2] \rightarrow \mathbb{R}$ is absolutely continuous function on $[v_1, v_2]$, and $f_2 : [v_1, v_2] \rightarrow \mathbb{R}$ is non-decreasing on $[v_1, v_2]$. Moreover, suppose that both $\hbar_1, \hbar_2 : [v_1, v_2] \rightarrow \mathbb{R}^+$ are positive integrable. If $(f'_1) \in L^\infty[v_1, v_2]$, then we have

$$\begin{aligned}
& \left| \mathcal{F} \mathcal{I}_{v_1+}^{\Psi} \hbar_1(v_2) \mathcal{F} \mathcal{I}_{v_1+}^{\Psi} (f_1 f_2)(v_2) + \mathcal{F} \mathcal{I}_{v_1+}^{\Psi} (1) \mathcal{F} \mathcal{I}_{v_1+}^{\Psi} (f_1 f_2)(v_2) \right. \\
& - \left. \mathcal{F} \mathcal{I}_{v_1+}^{\Psi} (\hbar_1 f_1)(v_2) \mathcal{F} \mathcal{I}_{v_1+}^{\Psi} (f_2)(v_2) - \mathcal{F} \mathcal{I}_{v_1+}^{\Psi} (f_1)(v_2) \mathcal{F} \mathcal{I}_{v_1+}^{\Psi} (\hbar_1 f_2)(v_2) \right| \\
& \leq \frac{\|f'_1\|_{\infty}}{\omega(\theta)} \left[\mathcal{F} \mathcal{I}_{v_1+}^{\Psi} v_2 \hbar_1(v_2) \int_{v_1}^{\theta} \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\varrho))}{\mathcal{F}(v_2) - \mathcal{F}(\varrho)} \omega(\varrho) \mathcal{F}'(\varrho) d\varrho \right. \\
& \left. - \mathcal{F} \mathcal{I}_{v_1+}^{\Psi} \hbar_1(v_2) \int_{v_1}^{\theta} \varrho \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\varrho))}{\mathcal{F}(v_2) - \mathcal{F}(\varrho)} \omega(\varrho) \mathcal{F}'(\varrho) d\varrho \right] \int_{v_1}^{v_2} f'_2(\theta) d\theta.
\end{aligned}$$

Proof. Applying Theorem 6 for $\hbar_2(\theta) = 1, \theta \in [v_1, v_2]$, the desired result is obtained. \square

Corollary 9. Let \mathcal{F} be measurable, increasing, positive and monotone function on (v_1, v_2) , and having a continuous derivative \mathcal{F}' on (v_1, v_2) . Assume that $f_1 : [v_1, v_2] \rightarrow \mathbb{R}$ is absolutely continuous function on $[v_1, v_2]$, and $f_2 : [v_1, v_2] \rightarrow \mathbb{R}$ is non-decreasing on $[v_1, v_2]$. Moreover, suppose that both $\hbar_1, \hbar_2 : [v_1, v_2] \rightarrow \mathbb{R}^+$ are positive integrable. If $(f'_1) \in L^\infty[v_1, v_2]$, then we have

$$\begin{aligned}
& \left| \mathcal{F} \mathcal{I}_{v_1+}^{\Psi} (1) \mathcal{F} \mathcal{I}_{v_1+}^{\Psi} (f_1 f_2)(v_2) - \mathcal{F} \mathcal{I}_{v_1+}^{\Psi} (f_1)(v_2) \mathcal{F} \mathcal{I}_{v_1+}^{\Psi} (f_2)(v_2) \right| \\
& \leq \frac{\|f'_1\|_{\infty}}{2\omega(\theta)} \left[\mathcal{F} \mathcal{I}_{v_1+}^{\Psi} v_2 \int_{v_1}^{\theta} \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\varrho))}{\mathcal{F}(v_2) - \mathcal{F}(\varrho)} \omega(\varrho) \mathcal{F}'(\varrho) d\varrho \right. \\
& \left. - \mathcal{F} \mathcal{I}_{v_1+}^{\Psi} (1) \int_{v_1}^{\theta} \varrho \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\varrho))}{\mathcal{F}(v_2) - \mathcal{F}(\varrho)} \omega(\varrho) \mathcal{F}'(\varrho) d\varrho \right] \int_{v_1}^{v_2} f'_2(\theta) d\theta.
\end{aligned}$$

Proof. Applying Theorem 6 for $\hbar_1(\theta) = \hbar_2(\theta) = 1, \theta \in [v_1, v_2]$, the desired result is obtained. \square

Theorem 7. Let \mathcal{F} be measurable, increasing, positive and monotone function on (v_1, v_2) and having continuous derivative \mathcal{F}' on (v_1, v_2) . Assume that $f_1, f_2 : [v_1, v_2] \rightarrow \mathbb{R}$ are absolutely continuous on $[v_1, v_2]$ and $f_2 : [v_1, v_2] \rightarrow \mathbb{R}$ is nondecreasing on $[v_1, v_2]$. Suppose that $\hbar_1, \hbar_2 : [v_1, v_2] \rightarrow \mathbb{R}^+$ are positive integrable. If $f'_1, f'_2 \in L^\infty[v_1, v_2]$, then we have

$$\begin{aligned}
& \left| \frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^{\Psi} \hbar_1(v_2) \frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^{\Psi} (\hbar_2 f_1 f_2)(v_2) + \frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^{\Psi} \hbar_2(v_2) \frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^{\Psi} (\hbar_1 f_1 f_2)(v_2) \right. \\
& - \frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^{\Psi} (\hbar_2 f_1)(v_2) \frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^{\Psi} (\hbar_1 f_2)(x_2) - \frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^{\Psi} (\hbar_1 f_1)(v_2) \frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^{\Psi} (\hbar_2 f_2)(v_2) \Big| \\
& \leq \frac{\|f'_1\|_{\infty} \|f'_2\|_{\infty}}{\omega(\theta)} \left[\frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^{\Psi} v_2 \hbar_1(v_2) \int_{v_1}^{\theta} \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\varrho))}{\mathcal{F}(v_2) - \mathcal{F}(\varrho)} \omega(\varrho) \mathcal{F}'(\varrho) \hbar_2(\varrho) d\varrho \right. \\
& - 2 \frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^{\Psi} v_2 \hbar_1(v_2) \int_{v_1}^{\theta} \varrho \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\varrho))}{\mathcal{F}(v_2) - \mathcal{F}(\varrho)} \omega(\varrho) \mathcal{F}'(\varrho) \hbar_2(\varrho) d\varrho \\
& \left. + \frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^{\Psi} \hbar_1(v_2) \int_{v_1}^{\theta} \varrho^2 \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\varrho))}{\mathcal{F}(v_2) - \mathcal{F}(\varrho)} \omega(\varrho) \mathcal{F}'(\varrho) \hbar_2(\varrho) d\varrho \right]. \tag{16}
\end{aligned}$$

Proof. Consider the left-hand side of (16), we have

$$\begin{aligned}
& \left| \frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^{\Psi} \hbar_2(v_2) \frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^{\Psi} (\hbar_2 f_1 f_2)(v_2) + \frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^{\Psi} \hbar_2(v_2) \frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^{\Psi} (\hbar_1 f_1 f_2)(v_2) \right. \\
& - \frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^{\Psi} (\hbar_2 f_1)(v_2) \frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^{\Psi} (\hbar_1 f_2)(x_2) - \frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^{\Psi} (\hbar_1 f_1)(v_2) \frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^{\Psi} (\hbar_2 f_2)(v_2) \Big| \\
& \leq \frac{1}{\omega^2(\theta)} \left| \int_{v_1}^{v_2} \int_{v_1}^{v_2} \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\xi))}{\mathcal{F}(v_2) - \mathcal{F}(\xi)} \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\varrho))}{\mathcal{F}(v_2) - \mathcal{F}(\varrho)} \right. \\
& \times \omega(\xi) \mathcal{F}'(\xi) \hbar_1(\xi) \omega(\varrho) \mathcal{F}'(\varrho) \hbar_2(\varrho) [(f_1(\xi) - f_1(\varrho))(f_2(\xi) - f_2(\varrho))] d\xi d\varrho \Big| \\
& \leq \frac{1}{\omega^2(\theta)} \int_{v_1}^{v_2} \int_{v_1}^{v_2} \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\xi))}{\mathcal{F}(v_2) - \mathcal{F}(\xi)} \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\varrho))}{\mathcal{F}(v_2) - \mathcal{F}(\varrho)} \omega(\xi) \mathcal{F}'(\xi) \hbar_1(\xi) \omega(\varrho) \mathcal{F}'(\varrho) \hbar_2(\varrho) \\
& \times \left| \frac{(f_1(\xi) - f_1(\varrho))}{\xi - \varrho} \right| \left| \frac{(f_2(\xi) - f_2(\varrho))}{\xi - \varrho} \right| (\xi - \varrho)^2 d\xi d\varrho \\
& \leq \frac{\|f'_1\|_{\infty} \|f'_2\|_{\infty}}{\omega^2(\theta)} \int_{v_1}^{v_2} \int_{v_1}^{\theta} \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\xi))}{\mathcal{F}(v_2) - \mathcal{F}(\xi)} \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\varrho))}{\mathcal{F}(v_2) - \mathcal{F}(\varrho)} \omega(\xi) \mathcal{F}'(\xi) \hbar_1(\xi) \omega(\varrho) \mathcal{F}'(\varrho) \hbar_2(\varrho) \\
& \times (\xi^2 - 2\xi\varrho + \varrho^2) d\xi d\varrho \\
& \leq \frac{\|f'_1\|_{\infty} \|f'_2\|_{\infty}}{\omega^2(\theta)} \left[\int_{v_1}^{v_2} \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\xi))}{\mathcal{F}(v_2) - \mathcal{F}(\xi)} \xi^2 \omega(\xi) \mathcal{F}'(\xi) \hbar_1(\xi) d\xi \int_{v_1}^{\theta} \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\varrho))}{\mathcal{F}(v_2) - \mathcal{F}(\varrho)} \omega(\varrho) \mathcal{F}'(\varrho) \hbar_2(\varrho) d\varrho \right. \\
& - 2 \int_{v_1}^{v_2} \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\xi))}{\mathcal{F}(v_2) - \mathcal{F}(\xi)} \xi \omega(\xi) \mathcal{F}'(\xi) \hbar_1(\xi) d\xi \int_{v_1}^{\theta} \varrho \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\varrho))}{\mathcal{F}(v_2) - \mathcal{F}(\varrho)} \omega(\varrho) \mathcal{F}'(\varrho) \hbar_2(\varrho) d\varrho \\
& \left. + \int_{v_1}^{v_2} \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\xi))}{\mathcal{F}(v_2) - \mathcal{F}(\xi)} \omega(\xi) \mathcal{F}'(\xi) \hbar_1(\xi) d\xi \int_{v_1}^{\theta} \varrho^2 \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\varrho))}{\mathcal{F}(v_2) - \mathcal{F}(\varrho)} \omega(\varrho) \mathcal{F}'(\varrho) \hbar_2(\varrho) d\varrho \right].
\end{aligned}$$

Hence, by using (7), the proof of the theorem is completed. \square

Corollary 10. Let \mathcal{F} be measurable, increasing, positive and monotone function on (v_1, v_2) and having continuous derivative \mathcal{F}' on (v_1, v_2) . Assume that $f_1, f_2 : [v_1, v_2] \rightarrow \mathbb{R}$ are absolutely continuous on $[v_1, v_2]$ and $f_2 : [v_1, v_2] \rightarrow \mathbb{R}$ is nondecreasing on $[v_1, v_2]$. Suppose that $\hbar_1, \hbar_2 : [v_1, v_2] \rightarrow \mathbb{R}^+$ are positive integrable. If $f'_1, f'_2 \in L^{\infty}[v_1, v_2]$, then we have

$$\begin{aligned}
& \left| \frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^{\Psi} (1) \frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^{\Psi} (\hbar_2 f_1 f_2)(v_2) + \frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^{\Psi} \hbar_2(v_2) \frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^{\Psi} (f_1 f_2)(v_2) \right. \\
& - \frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^{\Psi} (\hbar_2 f_1)(v_2) \frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^{\Psi} (f_2)(x_2) - \frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^{\Psi} (f_1)(v_2) \frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^{\Psi} (\hbar_2 f_2)(v_2) \Big| \\
& \leq \frac{\|f'_1\|_{\infty} \|f'_2\|_{\infty}}{\omega(\theta)} \left[\frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^{\Psi} v_2 \int_{v_1}^{\theta} \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\varrho))}{\mathcal{F}(v_2) - \mathcal{F}(\varrho)} \omega(\varrho) \mathcal{F}'(\varrho) \hbar_2(\varrho) d\varrho \right. \\
& - 2 \frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^{\Psi} v_2 \int_{v_1}^{\theta} \varrho \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\varrho))}{\mathcal{F}(v_2) - \mathcal{F}(\varrho)} \omega(\varrho) \mathcal{F}'(\varrho) \hbar_2(\varrho) d\varrho \\
& \left. + \frac{\mathcal{F}}{\omega} \mathcal{I}_{v_1+}^{\Psi} (1) \int_{v_1}^{\theta} \varrho^2 \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\varrho))}{\mathcal{F}(v_2) - \mathcal{F}(\varrho)} \omega(\varrho) \mathcal{F}'(\varrho) \hbar_2(\varrho) d\varrho \right].
\end{aligned}$$

Proof. Setting $\hbar_1(\theta) = 1$, $\theta \in [v_1, v_2]$ in Theorem 7, then the desired result is obtained. \square

Corollary 11. Let \mathcal{F} be measurable, increasing, positive and monotone function on (v_1, v_2) and having continuous derivative \mathcal{F}' on (v_1, v_2) . Assume that $f_1, f_2 : [v_1, v_2] \rightarrow \mathbb{R}$ are absolutely continuous on $[v_1, v_2]$ and $f_2 : [v_1, v_2] \rightarrow \mathbb{R}$ is nondecreasing on $[v_1, v_2]$. Suppose that $\hbar_1, \hbar_2 : [v_1, v_2] \rightarrow \mathbb{R}^+$ are positive integrable. If $f'_1, f'_2 \in L^\infty[v_1, v_2]$, then we have

$$\begin{aligned} & \left| {}_{\omega}^{\mathcal{F}}\mathcal{I}_{v_1+}^{\Psi} \hbar_1(v_2) {}_{\omega}^{\mathcal{F}}\mathcal{I}_{v_1+}^{\Psi}(f_1 f_2)(v_2) + {}_{\omega}^{\mathcal{F}}\mathcal{I}_{v_1+}^{\Psi}(1) {}_{\omega}^{\mathcal{F}}\mathcal{I}_{v_1+}^{\Psi}(\hbar_1 f_1 f_2)(v_2) \right. \\ & \quad \left. - {}_{\omega}^{\mathcal{F}}\mathcal{I}_{v_1+}^{\Psi}(f_1)(v_2) {}_{\omega}^{\mathcal{F}}\mathcal{I}_{v_1+}^{\Psi}(\hbar_1 f_2)(x_2) - {}_{\omega}^{\mathcal{F}}\mathcal{I}_{v_1+}^{\Psi}(\hbar_1 f_1)(v_2) {}_{\omega}^{\mathcal{F}}\mathcal{I}_{v_1+}^{\Psi}(f_2)(v_2) \right| \\ & \leq \frac{\|f'_1\|_\infty \|f'_2\|_\infty}{\omega(\theta)} \left[{}_{\omega}^{\mathcal{F}}\mathcal{I}_{v_1+}^{\Psi} v_2 \hbar_1(v_2) \int_{v_1}^{\theta} \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\varrho))}{\mathcal{F}(v_2) - \mathcal{F}(\varrho)} \omega(\varrho) \mathcal{F}'(\varrho) d\varrho \right. \\ & \quad \left. - 2 {}_{\omega}^{\mathcal{F}}\mathcal{I}_{v_1+}^{\Psi} v_2 \hbar_1(v_2) \int_{v_1}^{\theta} \varrho \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\varrho))}{\mathcal{F}(v_2) - \mathcal{F}(\varrho)} \omega(\varrho) \mathcal{F}'(\varrho) d\varrho \right. \\ & \quad \left. + {}_{\omega}^{\mathcal{F}}\mathcal{I}_{v_1+}^{\Psi} \hbar_1(v_2) \int_{v_1}^{\theta} \varrho^2 \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\varrho))}{\mathcal{F}(v_2) - \mathcal{F}(\varrho)} \omega(\varrho) \mathcal{F}'(\varrho) d\varrho \right]. \end{aligned}$$

Proof. Setting $\hbar_2(\theta) = 1$, $\theta \in [v_1, v_2]$ in Theorem 7, then the desired result is proven. \square

Corollary 12. Let \mathcal{F} be measurable, increasing, positive and monotone function on (v_1, v_2) and having continuous derivative \mathcal{F}' on (v_1, v_2) . Assume that $f_1, f_2 : [v_1, v_2] \rightarrow \mathbb{R}$ are absolutely continuous on $[v_1, v_2]$ and $f_2 : [v_1, v_2] \rightarrow \mathbb{R}$ is nondecreasing on $[v_1, v_2]$. Suppose that $\hbar_1, \hbar_2 : [v_1, v_2] \rightarrow \mathbb{R}^+$ are positive integrable. If $f'_1, f'_2 \in L^\infty[v_1, v_2]$, then we have

$$\begin{aligned} & \left| {}_{\omega}^{\mathcal{F}}\mathcal{I}_{v_1+}^{\Psi}(1) {}_{\omega}^{\mathcal{F}}\mathcal{I}_{v_1+}^{\Psi}(f_1 f_2)(v_2) + {}_{\omega}^{\mathcal{F}}\mathcal{I}_{v_1+}^{\Psi}(1) {}_{\omega}^{\mathcal{F}}\mathcal{I}_{v_1+}^{\Psi}(f_1 f_2)(v_2) \right. \\ & \quad \left. - {}_{\omega}^{\mathcal{F}}\mathcal{I}_{v_1+}^{\Psi}(f_1)(v_2) {}_{\omega}^{\mathcal{F}}\mathcal{I}_{v_1+}^{\Psi}(f_2)(x_2) - {}_{\omega}^{\mathcal{F}}\mathcal{I}_{v_1+}^{\Psi}(f_1)(v_2) {}_{\omega}^{\mathcal{F}}\mathcal{I}_{v_1+}^{\Psi}(f_2)(v_2) \right| \\ & \leq \frac{\|f'_1\|_\infty \|f'_2\|_\infty}{\omega(\theta)} \left[{}_{\omega}^{\mathcal{F}}\mathcal{I}_{v_1+}^{\Psi} v_2 \int_{v_1}^{\theta} \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\varrho))}{\mathcal{F}(v_2) - \mathcal{F}(\varrho)} \omega(\varrho) \mathcal{F}'(\varrho) d\varrho \right. \\ & \quad \left. - 2 {}_{\omega}^{\mathcal{F}}\mathcal{I}_{v_1+}^{\Psi} v_2 \int_{v_1}^{\theta} \varrho \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\varrho))}{\mathcal{F}(v_2) - \mathcal{F}(\varrho)} \omega(\varrho) \mathcal{F}'(\varrho) d\varrho \right. \\ & \quad \left. + {}_{\omega}^{\mathcal{F}}\mathcal{I}_{v_1+}^{\Psi} \int_{v_1}^{\theta} \varrho^2 \frac{\Psi(\mathcal{F}(v_2) - \mathcal{F}(\varrho))}{\mathcal{F}(v_2) - \mathcal{F}(\varrho)} \omega(\varrho) \mathcal{F}'(\varrho) d\varrho \right]. \end{aligned}$$

Proof. Setting $\hbar_1(\theta) = \hbar_2(\theta) = 1$, $\theta \in [v_1, v_2]$ in Theorem 7, then the desired result is obtained. \square

Remark 6. One can easily derive some new inequalities by applying the following conditions.

- i. Setting $\hbar_1(\theta) = \hbar_2(\theta)$, $\mathcal{F}(\theta) = \theta$ and $\Psi(\mathcal{F}(\theta)) = \theta$ throughout in the paper.
- ii. Setting $\hbar_1(\theta) = \hbar_2(\theta) = 1$ and $\mathcal{F}(\theta) = \theta$ and $\Psi(\mathcal{F}(\theta)) = \theta$ throughout in the paper.

Remark 7. Throughout in this article, if we put $\omega(\theta) = 1$ and $\Psi(\mathcal{F}(\theta)) = \frac{\mathcal{F}(\theta)^\kappa}{\Gamma(\kappa)}$, then all the inequalities will reduce to the work of Bezzou et al. [27].

Remark 8. Throughout in this article, if we consider $\omega(\theta) = 1$ and $\Psi(\mathcal{F}(\theta)) = \frac{\mathcal{F}(\theta)^{\frac{\kappa}{\eta}}}{\Gamma_\eta(\kappa)}$, then all the inequalities will reduce to the work of Rahman et al. [41].

Remark 9. Taking $\hbar_1(\theta) = \hbar_2(\theta)$, $\omega(\theta) = 1$, $\Psi(\mathcal{F}(\theta)) = \theta$ and $\Psi(\mathcal{F}(\theta)) = \frac{(\theta)^\kappa}{\Gamma(\kappa)}$ in Theorems 4–7, the results of Bezzou et al. [42] are restored.

3. Concluding Remarks

In the study of mathematics and related subjects, mathematical inequalities are extremely important. Fractional integral inequalities are now useful in determining the uniqueness of fractional partial differential equation solutions. They also guarantee the boundedness of fractional boundary value problem solutions. These suggestions have promoted the future research in the subject of integral inequalities to investigate the extensions of integral inequalities using fractional calculus operators. In the present investigation, we have proposed some double-weighted generalized fractional integral inequalities by utilizing more generalized class of fractional integrals associated with integrable, measurable, positive and monotone function \mathcal{F} in its kernel. The derived inequalities are more general than the existing inequalities cited therein. All the classical inequalities can be easily restored by applying specific conditions on \mathcal{F} and $\Psi(\theta)$ given in Remark 3. Also, we can derive some new weighted type double fractional integral inequalities by applying specific conditions on \mathcal{F} and $\Psi(\theta)$ given in Remark 2. In future research, some new other type of inequalities will be derived by employing the proposed operator. The special cases of the obtained result can be found in [24,25,27,41,42].

Author Contributions: Conceptualization, G.R. and M.S.; methodology, G.R.; software, G.R. and M.S.; validation, G.R. and K.S.N.; formal analysis, G.R. and K.S.N.; investigation, G.R. and M.S.; resources, K.S.N.; data curation, G.R.; writing—original draft preparation, G.R.; writing—review and editing, G.R., S.F.A. and K.S.N.; visualization, K.S.N.; supervision, G.R., S.F.A. and K.S.N.; project administration, S.F.A. and K.S.N.; funding acquisition, S.F.A. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: Not applicable.

Conflicts of Interest: The authors declare that they have no competing interest.

References

1. Mitrovic, D.S.; Pecaric, J.E.; Fink, A. *Clasical and New Inequalities in Analysis*; Kluwer Academic: Dordrecht, The Netherlands, 1993.
2. Qi, F. Several integral inequalities. *J. Inequ. Pure Appl. Math.* **2000**, *1*, 1039–1042.
3. Sarikaya, M.Z.; Yildirim, H.; Saglam, A. On Hardy type integral inequality associated with the generalised translation. *Int. J. Contemp. Math. Sci.* **2006**, *1*, 333–340. [[CrossRef](#)]
4. Dahmani, Z.; Tabharit, L. On weighted Grüss type inequalities via fractional integration. *J. Adv. Res. Pure Math.* **2010**, *2*, 31–38. [[CrossRef](#)]
5. Dahmani, Z. New inequalities in fractional integrals. *Int. J. Nonlinear Sci.* **2010**, *9*, 493–497.
6. Rahman, G.; Nisar, K.S.; Mubeen, S.; Choi, J. Certain inequalities involving the (k, η) -fractional integral operator. *Far East J. Math. Sci. (FJMS)* **2018**, *103*, 1879–1888. [[CrossRef](#)]
7. Set, E.; Noor, M.A.; Awan, M.U.; Gözpınar, A. Generalized Hermite-Hadamard type inequalities involving fractional integral operators. *J. Inequ. Appl.* **2017**, *169*, *10*. [[CrossRef](#)]
8. Dahmani, Z. New classes of integral inequalities of fractional order. *Le Mat.* **2014**, *69*, 237–247.
9. Srivastava, S.M.; Kashuri, A.; Mohammed, P.O.; Nonlaopon, K. Certain Inequalities Pertaining to Some New Generalized Fractional Integral Operators. *Fractal Fract.* **2021**, *5*, 160. [[CrossRef](#)]
10. Treană, S.; Singh, S. Weak sharp solutions associated with a multidimensional variational-type inequality. *Positivity* **2021**, *25*, 329–351. [[CrossRef](#)]
11. Treană, S.; Mititelu, Ş. Efficiency for variational control problems on Riemann manifolds with geodesic quasiinvex curvilinear integral functionals. *RACSAM* **2020**, *114*, 113. [[CrossRef](#)]
12. Ning, R.-J.; Liu, X.-Y.; Liu, Z. Conversion calculation method of multivariate integrals. *AIMS Math.* **2021**, *6*, 3009–3024. [[CrossRef](#)]
13. Usta, F.; Sarikaya, M.Z.; Qi, F. Explicit bounds on certain integral inequalities via conformable fractional calculus. *Cogent Math.* **2017**, *4*, 1–8. [[CrossRef](#)]
14. Usta, F.; Sarikaya, M.Z. Some Improvements of Conformable Fractional Integral Inequalities. *Int. J. Anal. Appl.* **2017**, *14*, 162–166. [[CrossRef](#)]

15. Usta, F.; Sarikaya, M.Z. On bivariate retarded integral inequalities and their applications. *Facta Univ. Ser. Math. Inform.* **2019**, *34*, 553–561.
16. Chebyshev, P.L. Sur les expressions approximatives des intégrales définies par les autres prises entre les mêmes limites. *Proc. Math. Soc. Charkov* **1882**, *2*, 93–98. [[CrossRef](#)]
17. Dahmani, Z. The Riemann-Liouville Operator to Generate Some New Inequalities. *Int. J. Nonlinear Sci.* **2011**, *12*, 452–455.
18. Mitrovic, D.S. *Analytic Inequalities*; Springer: Berlin, Germany, 1970.
19. Kuang, J.C. *Applied Inequalities*, 3rd ed.; Shandong Science and Technology Press: Jinan, China, 2004. (In Chinese)
20. Belarbi, S.; Dahmani, Z. On some new fractional integral inequalities. *J. Inequ. Pure Appl. Math.* **2009**, *10*, 1–12.
21. Dahmani, Z.; Benzidane, A. New inequalities using Q-fractional theory. *Bull. Math. Anal. Appl.* **2012**, *4*, 190–196.
22. Mercer, A.M. An improvement of the Gruss inequality. *J. Inequ. Pure Appl. Math.* **2005**, *10*, 93.
23. Ostrowski, A.M. On an integral inequality. *Aequ. Math.* **1970**, *4*, 358–373.
24. Rahman, G.; Ullah, Z.; Khan, A.; Set, E.; Nisar, K.S. Certain Chebyshev type inequalities involving fractional conformable integral operators. *Math. Math.* **2019**, *7*, 364. [[CrossRef](#)]
25. Tassaddiq, A.; Rahman, G.; Nisar, K.S.; Samraiz, M. Certain fractional conformable inequalities for the weighted and the extended Chebyshev functionals. *Adv. Differ. Equ.* **2020**, *2020*, 96. [[CrossRef](#)]
26. Awan, K.M.; Pecaric, J.; Rehman, A. Steffensen's generalization of Chebyshev inequality. *J. Math. Inequ.* **2015**, *9*, 155–163. [[CrossRef](#)]
27. Bezzou, M.; Dahmani, Z.; Khameli, A. On some double-weighted fractional integral inequalities. *Sarajevo J. Math.* **2019**, *15*, 23–36. [[CrossRef](#)]
28. Dahmani, Z.; Bounoua, M.D. Further results on Chebyshev and Steffensen inequalities. *Kyungpook Math. J.* **2018**, *58*, 55–66.
29. Sarikaya, M.Z.; Yildirim, H. On generalization of the Riesz potential. *Indian J. Math. Math. Sci.* **2007**, *3*, 231–235.
30. Rahman, G.; Hussain, A.; Ali, A.; Nisar, K.S.; Mohamed, R.N. More General Weighted-Type Fractional Integral Inequalities via Chebyshev Functionals. *Fractal Fract.* **2021**, *5*, 232. [[CrossRef](#)]
31. Jarad, F.; Abdeljawad, T.; Shah, K. On the weighted fractional operators of a function with respect to another function. *Fractals* **2020**, *28*, 2040011. [[CrossRef](#)]
32. Jarad, F.; Ugurlu, E.; Abdeljawad, T.; Baleanu, D. On a new class of fractional operators. *Adv. Differ. Equ.* **2017**, *2017*, 247. [[CrossRef](#)]
33. Khalil, R.; Horani, M.A.; Yousef, A.; Sababheh, M. A new definition of fractional derivative. *J. Comput. Appl. Math.* **2014**, *264*, 6570. [[CrossRef](#)]
34. Khan, T.U.; Khan, M.A. Generalized conformable fractional integral operators. *J. Comput. Appl. Math.* **2018**, *346*, 378–389. [[CrossRef](#)]
35. Mubeen, S.; Habibullah, G.M. k-Fractional integrals and application. *Int. J. Contemp. Math. Sci.* **2012**, *7*, 89–94. [[CrossRef](#)]
36. Sarikaya, M.Z.; Ertugral, F. On the generalized Hermite-Hadamard inequalities. *Ann. Univ. Craiova Math. Comput. Sci. Ser.* **2020**, *47*, 193–213.
37. Kilbas, A.A.; Srivastava, H.M.; Trujillo, J.J. *Theory and Application of Fractional Differential Equations*; Elsevier: Amersterdam, The Netherlands, 2006.
38. Samko, S.G.; Kilbas, A.A.; Marichev, O.I. *Fractional Integrals and Derivatives, Theory and Applications*; Nikol'ski, S.M., Ed.; Translated from the 1987 Russian Original, Revised by the Authors; Gordon and Breach Science: Yverdon, Switzerland, 1993.
39. Katugampola, U.N. New approach to a generalized fractional integral. *Appl. Math. Comput.* **2011**, *218*, 860–865.
40. Dahmani, Z. About some integral inequalities using Riemann-Liouville integrals. *Gen. Math.* **2012**, *20*, 63–69. [[CrossRef](#)]
41. Rahman, G.; Nisar, K.S.; Ghanbari, B.; Abdeljawad, T. On generalized fractional integral inequalities for the monotone weighted Chebyshev functionals. *Adv. Differ. Equ.* **2020**, *2020*, 368. [[CrossRef](#)]
42. Bezzou, M.; Dahmani, Z.; Khameli, A. Some weighted inequalities of Chebyshev type via RL-approach. *Mathematica* **2018**, *60*, 12–20. [[CrossRef](#)]