

Abstract

Screening of Some Apiaceae and Asteraceae Plants for Their Cytotoxic Potential [†]

Perihan Gürbüz ^{1,*}, Kübra Uzun ¹, Ebru Öztürk ² and Mükerrerem Betül Yerer ²

¹ Pharmacognosy Department, Faculty of Pharmacy, Erciyes University, Kayseri 38039, Turkey

² Pharmacology Department, Faculty of Pharmacy, Erciyes University, Kayseri 38039, Turkey

* Correspondence: pgurbuz@erciyes.edu.tr; Tel.: +90-553-355-3870

[†] Presented at the 2nd International Conference on Natural Products for Cancer Prevention and Therapy, Kayseri, Turkey, 8–11 November 2017.

Published: 13 November 2017

Abstract: Terpenoids, especially sesquiterpene lactones (SLS) and furanocoumarins are diverse secondary metabolite groups which show a promising efficiency on different cancer cell lines. Therefore the screening of terpenoid and/or coumarin rich plants from the plant kingdom remains still importance. Breast cancer and lung adenocarcinoma are the most common cancer types leading to cancer-related deaths in the world. To investigate the cytotoxic effects of some Asteraceae plants [*Jurinea spec.*, *Ptilostemon spec.*, *Cirsium spec.*, *Centaurea spec.*, *Tanacetum spec.*, *Tragopogon spec.*, *Pulicaria spec.*, *Cyanus spec.*] and Apiaceae [*Heracleum spec.*, *Pastinaca spec.*] plant extracts, real time cell analyzer xCELLigence system was used. A549 (12.500 cells/well) and MCF-7 (5000 cells/well) cells were seeded in E-plate then approximately 24 h post-seeding when the cells were in the log growth phase, the cells were treated with the samples. Cell viability was observed during 48 h after treatment and IC₅₀ values have been calculated. According to results, *Jurinea macrocephala*, *Tanacetum parthenium* and *Pulicaria dysenterica* MeOH extracts showed significant cytotoxic effects on both two cell lines. The IC₅₀ values of those extracts on MCF-7 cell line were calculated as 17 µg/mL, 21.47 µg/mL and 27.05 µg/mL respectively after the end of the 24 h. To understand the secondary metabolite profiles of the extracts; crude ¹H NMR experiments were performed. For each active extracts the major compounds and/or group of compounds were discussed.

Keywords: Apiaceae; Asteraceae; A549; MCF-7; SLS; furanocoumarin

Conflicts of Interest: The authors declare no conflict of interest.



© 2017 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).