



Proceedings Changes in Blocking Characteristics during the First Part of the 21st Century ⁺

Anthony R. Lupo 1,*, Andrew D. Jensen 2, Igor I. Mokhov 3,4 and Mirseid G. Akperov 3

- ¹ Atmospheric Science Program, School of Natural Resources, University of Missouri, Columbia, MO 65211, USA
- ² Department of Mathematics and Meteorology, Northland College, Ashland, WI 54806, USA; ajensen@northland.edu
- ³ A.M. Obukhov Institute of Atmospheric Physics, Russian Academy of Science, 3 Pyzhevsky, Moscow 119017, Russia; mokhov@ifaran.ru (I.I.M.); akperovm@ifaran.ru (M.G.A.)
- ⁴ Faculty of Physics, Lomonosov State University, Moscow 119991, Russia
- * Correspondence: lupoa@missouri.edu; Tel.: +1-573-489-8457
- + Presented at the 2nd International Electronic Conference on Atmospheric Sciences, 16–31 July 2017; Available online: http://sciforum.net/conference/ecas2017.

Published: 19 July 2017

Abstract: A global blocking climatology published by this group for events that occurred during the late 20th century examined the comprehensive list of characteristics that included block intensity. In addition to confirming the results of other published climatologies, they found that Northern Hemisphere blocking was stronger than Southern Hemisphere events and winter events are stronger than summer ones. This work also examined the interannual variability of blocking as related to El Niño. Since this time, there is evidence that the occurrence of blocking has increased globally. A comparison of blocking characteristics during the first part of the 21st century to those in the late 20th century shows that the number of blocking events and their duration have increased in the Northern and Southern Hemisphere. The intensity of blocking has decreased by about nine percent in the Northern Hemisphere, but there was little change in the intensity of Southern Hemisphere events. Additionally, there is little or no change in the genesis regions of blocking. An examination of variability related to El Niño and Southern Oscillation reveals that the variability found in the earlier work has reversed. This could either be the result of interdecadal variability or a change in the climate.

Keywords: blocking; ENSO; climate change; Pacific Decadal Oscillation

1. Introduction

Many recent studies have examined the climatological behavior of blocking in both the northern and Southern Hemispheres in the later part of the 20th century [1–7]. Only a few have examined blocking in the 21st century (e.g., [8,9]). Many have also examined projections for the future occurrence of blocking [7,10–13]. Also, there is some evidence to suggest that there has been an increase in the occurrence of blocking since 2000 in both the northern and Southern Hemispheres.

The study of [1,7] studies examine a long list of blocking characteristic including the number of events, days, duration, intensity, and location at onset. Most other climatologies use only the occurrence, day, and/or durations of blocking events. The work of [5] suggested interannual and interdecadal variability in block occurrences. Then [6] implies that the number of Northern Hemisphere blocking events is increasing, but that the period 1970–1999 was likely a period of low blocking occurrence. Many authors projected that the occurrence of blocking many increase in a warmer world [7,10,12,13]. Others projected that far into the future the number of events may

decrease slightly [11]. Others, such as [8,9] suggest that the primary location for block occurrences would also change in a warmer world.

The goal of this work is to make a comprehensive comparison of the global occurrence of blocking since the end of the climatology of [1] to their work for the late 20th century. This work will examine the more comprehensive characteristics that they researched. This study will also determine if there have been any changes in the trends or interannual variability of any of these variables, as well as the possibility of interdecadal variability in these variables. This study is unique in that it is the only study that includes an examination of block intensity (BI). BI has been shown to be related to the 500 hPa height gradients in each hemisphere [1]. This work will primarily use the National Centers of Environmental Prediction/National Center for Atmospheric Research (NCEP/NCAR) reanalyses.

2. Data and Methods

2.1. Data

The data used here were the NCEP/NCAR reanalyses 500 hPa height fields on a 2.5° latitude by 2.5° longitude gridded data set available at 6-h intervals [14,15]. The 1200 UTC data was used primarily in the calculation of intensity since these tended to have the most number of observation data. This study also used data archived at [16] which contains a list of all blocking events since the beginning of the [1] study. The period of study in the Northern Hemisphere is from July 1998 to June 2017 (19 additional seasons), while for the Southern Hemisphere included the period from January 2000 to December 2016 (17 additional seasons). These time periods (spatial regions) follow the blocking year (geographic regions) established by [1].

2.2. Methodology

The blocking criterion and block intensity used here were established by [1] and references therein. The definition for ENSO used here is described in [17] and references therein and a short description is given here. The Japanese Meteorological Agency (JMA) El Niño and Southern Oscillation (ENSO) index is available through the Center for Ocean and Atmospheric Prediction Studies (COAPS) from 1868 to present [18]. The JMA classifies ENSO phases using SST within the bounded region of 4° S to 4° N, 150° W to 90° W, and defines the inception of an ENSO year as 1 October, and its conclusion on 30 September of the next year. This index is widely used in other published works (see [17] and references therein) and a list of years is given in (Table 1). Also, The work of [19] found that while the JMA index is more sensitive to La Niña events than other definitions, it is less sensitive than other indices to El Niño events.

El Niño	Neutral	La Niña
1969	1968	1967
1972	1977–1981	1970–1971
1976	1983–1985	1973–1975
1982	1989–1990	1988
1986–1987	1992–1996	1998–1999
1991	2000-2001	2007
1997	2003-2005	2010
2002	2008	
2006	2011-2013	
2009	2016	
2014-2015		

Table 1. List of ENSO years used here. The years are taken from [18].

The Pacific Decadal Oscillation (PDO) positive and negative modes are catalogued also by the Center for Ocean-Atmospheric Prediction Studies (COAPS). The most important effect of PDO is how

it interacts with ENSO during certain phases to create an enhanced effect on temperatures and precipitation variability (e.g., [20,21]). The characteristics of these modes are less pronounced than those for ENSO due to the fifty-to seventy-year cycle of PDO [22,23]. The positive phase of the PDO is recognized as the period 1977–1998, and the negative phase is recognized as the years 1947–1976, and 1999–2017.

3. The Climatology of Blocking in the Early 21st Century

3.1. The Northern Hemisphere

The noteworthy character of blocking at the start of the 21st century (Table 2) is the increases in the occurrence, number of blocking days and durations. In all regions and seasons, these increases are statistically significant at the 95% confidence level, using the Z-score test for means [24]. The Atlantic region showed the weakest increases which is consistent with the results of [7]. The large increase in the occurrence and days within Pacific Region Blocking is consistent with the results of [17] who examined spring and summer blocking in this region. The work of [25] also showed a large increase in blocking for the eastern Europe and western Russia region for the spring and summer seasons as well. Only the block intensity (BI) showed a decrease over the first part of the 21st century. This decrease is not significant at any standard level of statistical significance.

Table 2. Character of Northern Hemisphere blocking events per year as a function of region and season for all characteristics since the study of [1]. The change in the value is expressed as a percentage.

Region	Occurrence	Duration	Days	Intensity
Atlantic	15.6/+21.9	10.5/+24.5	163.8/+42.3	
Pacific	11.8/+77.9	9.5/+25.0	112.1/+111.4	
Continental	10.1/+84.7	9.0/+10.8	90.9/+81.1	
Total	37.6/+56.0	9.5/+11.8	356.3/+67.9	3.02/-6.3
Season	Occurrence	Duration	Days	Intensity
Summer	7.9/+51.0	10.0/+30.7	79.3/+85.5	
Fall	8.3/+52.7	9.7/+19.8	80.5/+71.3	
Winter	10.1/+42.3	9.7/+9.0	98.0/+46.1	
Spring	11.2/+53.4	9.7/+23.4	109.2/+72.4	

Such increases across all regions and seasons were suprising, however, as shown above, several studies have examined blocking over limted regions and seasons and shown similar strong inceases. While [17,25] showed increases within the warm season in recent decades, on the other hand, the study of [6] showed that over the Atlantic for the winter season that blocking days over the same period covered by [1] was associated with a significant minimum. Additionally, the work of [7] showed an increase in blocking since 2000 as shown with three blocking indexes including the index used here. Taken together, these provide confidence that the increases in Northern Hemisphere blocking occurrence is likely real. Finally, the mean number of simultaneous blocking days (defined as the number of days with two or more blocking events occurring concurrently in any sectors) per year since [1] is 102.1 days or 28.0% of the year. This represents a large increase over [1] who found the number to be approximately 8.7% of the year.

3.2. The Southern Hemisphere

Examining blocking in the Southern Hemisphere (Table 3) and comparing with the work of [1] shows similar results to the Northern Hemisphere above. The weakest increases occurred within the Pacific Region, and unlike the Northern Hemisphere this region is the predominant blocking region accounting for about three-quarters of all events when counting events and blocking days. The Pacific Region increases in the Southern Hemisphere were weaker than their Northern Hemisphere counterparts, but still statistically significant at the 95% confdence level. Increases within the Atlantic

or Indian Ocean Regions were very large, however, blocking in this region is not as prominent. Only BI showed no significant change in the Southern Hemisphere. Lastly, the number of simultaneous blocking days ber year since [1] in the Southern Hemisphere is 15.8 days per year (4.3%) which is higher than the 1.5% found by [1] and this will be discussed more below.

Region	Occurrence	Duration	Days	Intensity
Atlantic	1.8/+81.4	6.6/+11.4	11.7/+101.7	
Pacific	11.3/+43.1	8.4/+11.4	95.0/+59.9	
Indian	2.5/+181.1	7.3/+9.8	18.4/+206.7	
Total	15.6/+599	8.0/+9.6	124.7/+75.1	2.86/+1.2
Season	Occurrence	Duration	Days	Intensity
Summer	1.9/+52.8	7.0/+9.4	13.1/+67.9	
Fall	5.1/+47.3	8.2/+8.5	41.8/+60.2	
Winter	5.4/+65.6	8.5/+11.2	47.2/+90.3	
Spring	3.2/+80.0	7.5/+10.0	24.2/+96.7	

Table 3. Character of Southern Hemisphere blocking events per year as a function of region and season for all characteristics since the study of [1]. The change in the value is expressed as a percentage.

4. Interannual and Interdecadal Variability

In this section, the change in ENSO variability as well as PDO variability will be examined. In order to accomplish the latter, the data from [1] will be reanalyzed by stratifying the blocking years using the PDO epochs defined in Section 2.2. The change in phase of the PDO occurred near the end of the [1] study, thus it might be reasonable to assume that the recent increases in blocking are related to the change in phase of the PDO. The results from the Northern Hemisphere are shown in Table 4. The results show that the total number of blocking events, days, and durations were higher for the –PDO, but the BI indicated weaker events similar to the occurrence of blocking after the [1] study. Additionally, during the +PDO, the ENSO variability matched that of [1] which implied that La Niña years were associated with more blocking events and days and which were slightly stronger. Then during the –PDO phase the ENSO variability was the opposite in that El Niño years were associated with the greater number of blocking events, days, and slightly stronger. Thus, by examining the overall set of years (49) there was little ENSO variability overall.

In the Southern Hemisphere (Table 5), as in the Northern Hemisphere, the occurrence of blocking was greater during the –PDO phase. In this part of the world, however, the ENSO variability did not change across the phase of the PDO. During both PDO epochs, blocking occurred more often, was more persistent and more intense in El Niño and Neutral years as compared to La Niña years. This variability is reflected across each region as well. The study of [1] speculated that blocking was on the decrease during the late 20th century. Lastly, the work of [26] showed that there was a reversal in the proportion of east Pacific blocking occurrences when compared to those of the west Pacific Region. During the +PDO phase there were a larger fraction of east Pacific Region blocking during La Niña and Neutral years, while the opposite was true during the –PDO epoch (La Niña years showed the smallest fraction of east Pacific blocking).

The work of [1] speculated that the greater occurrence of simultaneous blocking events in the Northern Hemisphere versus those in the Southern Hemisphere was due to the greater number of occurrence of blocking since previous work demonstrated that the occurrence of blocking was caused local synoptic processes rather than hemisphere-wide planetary-scale processes. The results here show a similar result since the fewest number of blocking events occurred during the Southern Hemisphere +PDO epoch as well as the fewest number of simultaneous days. The opposite was true in that the largest number of simultaneous blocking days occurred during the Northern Hemisphere -PDO epoch which showed the largest number of blocking occurrences. The correlation between the number of simultaneous blocking days and the number of events was 0.89 in the Northern

Hemisphere and 0.76 for the Southern Hemisphere and both results are statistically significant at the 99% confidence level.

+PDO	Occurrence	Duration	Days	Intensity	% Simult.
El Niño (6)	23.5	8.1	190.4	3.06	7.6
Neutal (15)	24.4	8.2	200.1	3.26	8.9
La Niña (2)	30.5	8.3	253.3	3.11	12.7
Total (23)	24.7	8.2	202.5	3.20	8.9
-PDO	Occurrence	Duration	Days	Intensity	% Simult.
El Niño (7)	37.0	9.6	355.2	3.13	24.4
Neutral (11)	36.4	10.0	364.0	3.01	28.2
La Niña (8)	29.1	8.4	244.4	3.08	13.9
Total (26)	32.8	9.9	323.7	3.07	21.7

Table 4. Character of Northern Hemisphere blocking events per year as a function of ENSO and PDO. The number of years in each category is shown in parenthesis.

Table 5. Character of Southern Hemisphere blocking events per year as a function of ENSO and PDO. The number of years in each category is shown in parenthesis.

+PDO	Occurrence	Duration	Days	Intensity	% Simult.
El Niño (5)	9.0	7.0	63.0	3.02	1.7
Neutal (15)	9.5	7.1	67.5	2.76	1.4
La Niña (2)	6.0	6.7	40.2	1.87	0.0
Total (22)	9.0	7.1	63.9	2.81	1.3
-PDO	Occurrence	Duration	Days	Intensity	% Simult.
El Niño (8)	14.8	8.4	124.3	2.96	3.9
Neutral (9)	16.0	7.8	124.8	2.80	4.2
La Niña (8)	12.0	7.6	91.2	2.75	2.7
Total (25)	14.3	8.0	114.4	2.85	3.6

5. Discussion, Summary, and Conclusions

This study examined the occurrence of blocking over the entire globe since the study of [1], and including such characteristics as occurrence, duration, blocking days, block intensity (BI), and the number of simultaneous blocking days. The data set used here was the NCEP/NCAR re-analyses and the archive of blocking events are found at [16]. Then, the long-term trends as well as interannual and interdecadal variability were examined.

The study here showed statistically significant increases in block occurrences, days, and duration since the end of the 20th century. In the Northern Hemisphere, there was a slight decrease in BI, which was not statistically significant. There was little change in the Southern Hemisphere BI. In the Northern Hemisphere, the increases found here were consistent with the results of others [6,7,17,25] who examined 'partial' climatologies examining only certain regions during certain seasons. The work of [6] implied that the period 1970–1999 showed a relative minimum in Atlantic Region winter season blocking. Thus, there is strong evidence to support the results here. In the Southern Hemisphere, Pacific Region blocking showed increases that were not as strong as their Northern Hemisphere counterparts.

Separating the occurrence of blocking by ENSO and PDO showed that the +PDO epoch interannual variability in the occurrence, duration, and BI were similar to those found in [1] as expected. In the Southern Hemisphere, the interannual variability was the same in the –PDO epoch albeit with increases in occurrence, duration, and days. But, within the Pacific Region, the relative occurrence of East Pacific blocking was greater in La Niña and Neutral years than during El Niño years for the +PDO epoch. The opposite occurred during the –PDO phase. In the Northern Hemisphere, the interannual variability of these characteristics was opposite during the –PDO epoch.

Finally, an examination of the occurrence of simultaneous blocking events was greater with a greater frequency in blocking occurrence, supporting the conjecture of [1].

Acknowledgments: The authors acknowledge the reviewer for their time and effort in examining this manuscript.

Author Contributions: All four authors conceived and designed the experiments; performed the experiments; all four co-authors analyzed the data; and A.R. Lupo wrote the paper.

Conflicts of Interest: The authors declare no conflict of interest.

References

- 1. Wiedenmann, J.M.; Lupo, A.R.; Mokhov, I.I.; Tikhonova, E.A. The climatology of blocking anticyclones for the Northern and Southern Hemisphere: Block intensity as a diagnostic. *J. Clim.* **2002**, *15*, 3459–3474.
- 2. Pelly, J.L.; Hoskins, B.J. A new perspective on blocking. J. Atmos. Sci. 2003, 60, 743–755.
- 3. Schwierz, C.; Croci-Maspoli, M.; Davies, H. Perspicacious indicators of atmospheric blocking. *Geophys. Res. Lett.* **2004**, *31*, L06125.
- 4. Croci-Maspoli, M.; Schweirz, C.; Davies, H. Atmospheric blocking: space-time links to the NAO and PNA. *Clim. Dyn.* **2007**, *29*, 713–725**.**
- 5. Barriopedro, D.; Garcia-Herrera, R.; Lupo, A.R.; Hernandez, E. A climatology of Northern Hemisphere Blocking. *J. Clim.* **2006**, *19*, 1042–1063.
- 6. Häkkinen, S.; Rhines, P.B.; Worthen, D.L. Atmospheric Blocking and Atlantic Multidecadal Ocean Variability. *Science* **2011**, *334*, 655–659.
- Mokhov, I.I.; Akperov, M.G.; Prokofyeva, M.A.; Timazhev, A.V.; Lupo, A.R.; Le Treut, H. Blockings in the Northern Hemisphere and Euro-Atlantic region: Estimates of changes from reanalyses data and model simulations. *Doklady* 2012, 449, 430–433.
- 8. Luo, D.; Xiao, Y.; Yao, Y.; Dai, A.; Simmonds, I.; Franzke, C. Impact of Ural Blocking on Winter Warm Arctic-Cold Eurasian Anomalies, Part I: Blocking-Induced Amplification. *J. Clim.* **2016**, *29*, 3925–3947, doi:10.1175/JCLI-D-15-0611.1.
- 9. Luo, D.; Xiao, Y.; Diao, Y.; Dai, A.; Franzke, C.; Simmonds, I. Impact of Ural Blocking on Winter Warm Arctic-Cold Eurasian Anomalies, Part II: The link to the North Atlantic Oscillation. *J. Clim.* **2016**, *29*, 3949–3971, doi:10.1175/JCLI-D-15-0612.1.
- Lupo, A.R.; Oglesby, R.J.; Mokhov, I.I. Climatological features of blocking anticyclones: A study of Northern Hemisphere CCM1 model blocking events in present-day and double CO₂ concentration atmospheres. *Clim. Dyn.* 1997, 13, 181–195.
- 11. Sillmann, J.; Croci-Maspoli, M. Present and future atmospheric blocking and its impact on European and extreme climate. *Geophys. Res. Lett.* **2009**, *36*, L10702.
- 12. Sciafe, A.A.; Copsey, D.; Gordon, C.; Harris, C.; Hinton, T.; Keeley, S.; O'Neill, A.; Roberts, M.; Williams, K. Improved Atlantic Winter Blocking in a Climate Model. *Geophys. Res. Lett.* **2010**, *38*, L23703.
- 13. Mokhov, I.I.; Timazhev, A.V.; Lupo, A.R. Changes in atmospheric blocking characteristics within Euro-Atlantic region and Northern Hemisphere as a whole in the 21st century from model simulations using RCP anthropogenic scenarios. *Glob. Planet. Chang.* **2014**, *122*, 265–270.
- 14. Kalnay, E.; Kanamitsu, M.; Kistler, R.; Collins, W.; Deaven, D.; Gandin, L.; Iredell, M.; Saha, S.; White, G.; Woollen, J. The NCEP/NCAR 40-year reanalysis project. *Bull. Am. Meteorol. Soc.* **1996**, *77*, 437–471.
- 15. NCEP/NCAR Reanalyses Project. 2016. Available online: http://www.esrl.noaa.gov/psd/data/reanalysis/ reanalysis.shtml (accessed on 11 May 2017).
- 16. University of Missouri Blocking Archive. Available online: http://weather.missouri.edu/gcc (accessed on 26 June 2017).
- Newberry, R.G.; Lupo, A.R.; Jensen, A.D.; Rodriges-Zalipynis, R.A. An analysis of the spring-to-summer transition in the West Central Plains for application to long range forecasting. *Atmos. Clim. Sci.* 2016, *6*, 375– 393.
- 18. Center for Ocean and Atmosphere Prediction Studies. Available online: http://www.coaps.fsu.edu (accessed on 11 May 2017).
- 19. Hanley, D.E.; Bourassa, M.A.; O'Brien, J.J.; Smith, S.R.; Spade, E.R. A Quantitative Evaluation of ENSO Indices. J. Clim. 2003, 16, 1249–1258, doi:10.1175/1520-0442(2003)16<1249:AQEOEI>2.0.CO;2.

- 20. Bove, M.C.; Elsner, J.B.; Landsea, C.W.; Niu, X.; O'Brien, J.J. Effects of El Niño on U.S. Landfalling Hurricanes, Revisited. *Bull. Am. Meteorol. Soc.* **1998**, *79*, 2477–2482.
- 21. Hu, Z.Z.; Huang, B. Interferential Impact of ENSO and PDO on Dry and Wet Conditions in the U.S. Great Plains. *J. Clim.* **2009**, *19*, 5500–5518.
- 22. Mantua, N.J.; Hare, S.R.; Zhang, Y.; Wallace, J.M.; Francis, R.C. A Pacific Interdecadal Climate Oscillation with Impacts on Salmon Production. *Bull. Am. Meteorol. Soc.* **1997**, *78*, 1069–1079, doi:10.1175/1520-0477(1997)078<1069:APICOW>2.0.CO;2.
- 23. Minobe, S. A 50–70-Year Climatic Oscillation over the North Pacific and North America. *Geophys. Res. Lett.* **1997**, 24, 683–686.
- 24. Neter, J.; Wasserman, W.; Whitmore, G.A. *Applied Statistics*, 3rd ed.; Allyn and Bacon Inc.: Boston, MA, United State 1988; p. 1066.
- 25. Lupo, A.R.; Mokhov, I.I.; Chendev, Y.G.; Lebedeva, M.G.; Akperov, M.G.; Hubbart, J.A. Studying summer season drought in western Russia. *Adv. Meteorol.* **2014**, 1–9, doi:10.1155/2014/942027.
- 26. Lupo, A.R.; Garcia, M.; Rojas, K.; Gilles, J. ENSO Related Seasonal Range Prediction over South America. In Proceedings of the 2nd International Electronic Conference on Atmospheric Sciences, 16–31 July 2017.



© 2017 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (http://creativecommons.org/licenses/by/4.0/).