

*Supplementary Materials*

## Role of *Helicobacter Pylori* in Upper Gastrointestinal Bleeding Among Ischemic Stroke Hospitalizations: A Nationwide Study of Outcomes

**Table S1.** ICD-9-CM codes used in this analysis.

Condition	ICD-9 CM Codes
Gastrointestinal bleeding	530.04, 531.00, 531.01, 531.40, 531.41, 531.10, 531.20, 531.21, 531.50, 531.51, 531.60, 531.61, 532.00, 532.01, 532.40, 532.41, 532.10, 532.11, 532.20, 532.21, 532.50, 532.51, 532.60, 532.61
<i>Helicobacter Pylori</i>	041.86
Hypertension	401–405
Diabetes mellitus	249.00, 250
Atrial fibrillation	427.31
Hypercholesterolemia	272.0, 272.1, 272.2
Hemorrhagic conversion	430, 431
Alcohol abuse/dependent	V11.3, 303, 305.0
Drug abuse	304, 305.2–305.9
Tobacco current/past use	V15.82, 305.1
Use of anticoagulant	V58.61
Use of antiplatelet	V58.63
Chronic use of NSAIDs	V58.64
Use of aspirin	V58.66
Use of IV tPA during same hospitalization or in a different institution within the last 24 hours prior to admission to facility	99.10, V45.88

**Table S2.** Deyo's modification of Charlson's co-morbidity index (CCI).

Condition	ICD-9-CM Codes	Charlson Score
Myocardial infarction	410–410.9	1
Congestive heart failure	428–428.9	1
Peripheral vascular disease	433.9, 441–441.9, 785.4, V43.4	1
Cerebrovascular disease	430–438	1
Dementia	290–290.9	1
Chronic pulmonary disease	490–496, 500–505, 506.4	1
Rheumatologic disease	710.0, 710.1, 710.4, 714.0–714.2, 714.81, 725	1
Peptic ulcer disease	531–534.9	1
Mild liver disease	571.2, 571.5, 571.6, 571.4–571.49	1
Diabetes	250–250.3, 250.7	1
Diabetes with chronic complications	250.4–250.6	2
Hemiplegia or paraplegia	344.1, 342–342.9	2
Renal disease	582–582.9, 583–583.7, 585, 586, 588–588.9	2
Any malignancy including leukemia and lymphoma	140–172.9, 174–195.8, 200–208.9	2
Moderate or severe liver disease	572.2–572.8	3
Metastatic solid tumor	196–199.1	6
AIDS	042–044.9	6