



## New Monocrystal Catalysts and Complex Photocatalysts: Synthesis and Characterization

Guest Editor:

**Prof. Dr. Jingfei Luan**

1. School of Physics, Changchun Normal University, Changchun 130032, China

2. State Key Laboratory of Pollution Control and Resource Reuse, School of the Environment, Nanjing University, Nanjing 210093, China

Deadline for manuscript submissions:

**30 June 2024**

### Message from the Guest Editor

Dear Colleagues,

This topic is for the preparation of new visible light-responsive photocatalysts which will be detected by all kinds of advanced experimental instruments, such as transmission electron microscope. Different preparation methods, such as the solvothermal method, will be utilized to prepare heterojunction photocatalysts, precious metal-doped composite and monocrystal photocatalysts. New photocatalysts will be used for the degradation of organic pollutants under visible light irradiation or under ultraviolet light illumination. The removal rate of poisonous organic pollutants and the removal rates of the total organic carbon will be analyzed and arranged. The intermediate product during photocatalytic degradation poisonous organic pollutants will be investigated. The degradation pathways of poisonous organic pollutants such as pentachlorophenol will be provided. The degradation mechanism using hydroxyl radical, superoxide anion and photoinduced holes will be revealed. Hydrogen production via water splitting will be realized by using new monocrystal photocatalysts or heterojunction photocatalysts under visible light irradiation or under ultraviolet light illumination.

