



## Cyclic Nucleotide Signaling and the Cardiovascular System

Guest Editors:

**Prof. Dr. Thomas Brand**

National Heart & Lung Institute,  
Imperial College London,  
Hammersmith Campus, London  
W12 0NN, UK

**Dr. Enno Klussmann**

1. Max-Delbrück-Center for  
Molecular Medicine in the  
Helmholtz Association (MDC),  
13125 Berlin, Germany  
2. DZHK (German Center for  
Cardiovascular Research),  
partner site Berlin, 10785 Berlin,  
Germany

Deadline for manuscript  
submissions:

**closed (20 December 2017)**

### Message from the Guest Editors

Cyclic nucleotides 3',5'-adenosine monophosphate and 3',5'-cyclic guanosine monophosphate play important roles in the control of cardiovascular function under physiological and pathological conditions. They are produced by adenylate and guanylate cyclases, respectively, bound by different effector proteins, and are subsequently degraded by phosphodiesterases. These proteins form nanodomains in specific locations in cardiac myocytes, such as the plasma membrane, t-tubules, and the nuclear envelope. Thereby, a highly-compartmentalized regulation of essential functions of cardiac myocytes, such as calcium cycling, excitation-contraction coupling, and cell-cell cohesion, is achieved. In cardiac myocytes, several effector proteins are expressed. Through the use of effector protein-specific agonists and antagonists and alternatively, with the help of genetic experiments, insight into their individual roles and cross-talk have been obtained.

The importance of the cyclic nucleotide pathway in both health and disease cannot be underestimated and up-to-date reviews of this important scientific field will be provided.

