

Supporting information (SI)

Recognition of 8-Oxo-2'-Deoxyguanosine in DNA Using the Triphosphate of 2'-Deoxycytidine Connecting the 1,3-Diazaphenoxazine Unit, dCdapTP

Takato Sakurada ¹, Yuta Chikada ^{1,2}, Ryo Miyahara ¹ and Yosuke Taniguchi ^{1,2,*}


1 Graduate School of Pharmaceutical Sciences, Kyushu University, 3-1-1 Maidashi, Higashi-ku, Fukuoka 812-8582, Japan; takatorres.31@gmail.com (T.S.); chikada.yuta.840@s.kyushu-u.ac.jp (Y.C.); c51339rm@gmail.com (R.M.)

2 Faculty of Medicine, Dentistry and Pharmaceutical Sciences, Okayama University, 1-1-1 Tsushima-naka, Kita-ku, Okayama 700-8530, Japan



* Correspondence: y-taniguchi@okayama-u.ac.jp; Tel.: +81-86-251-7964


Contents:

1. NMR spectra of new compounds (1–4, 8, 13–16)	p2
2. NMR spectra of corresponding triphosphate compounds (17–20)	p12





Compound 8







Compound 11






Compound 12








Compound **13**








Compound 14








Compound 15











Compound 1








Compound 2








Compound **3**








Compound 4








Compound 17








Compound 18





Compound 19





Compound 20

