


## Supplementary material



**Figure S1.** Type of CP.

**Table S1.** Pairwise Comparisons of Type of CP.

Sample 1-Sample 2	Test Statistic	Std. Error	Std. Test Statistic	Sig.	Adj. Sig. <sup>a</sup>
Spastic quadriplegia-Spastic diplegia	19.602	8.863	2.212	0.027	0.162
Spastic quadriplegia-Spastic hemiparesis	27.329	17.747	1.540	0.124	0.741
Spastic quadriplegia-Dyskinebtic	55.765	28.499	1.957	0.050	0.302
Spastic diplegia-Spastic hemiparesis	-7.727	17.780	-0.435	0.664	1.000
Spastic diplegia-Dyskinebtic	36.163	28.520	1.268	0.205	1.000
Spastic hemiparesis-Dyskinebtic	28.436	32.401	0.878	0.380	1.000

Each row tests the null hypothesis that the Sample 1 and Sample 2 distributions are the same.

A symptotic significances (2-sided tests) are displayed. The significance level is .05.

a. Significance values have been adjusted by the Bonferroni correction for multiple tests.

**Table S2.** Behavior related to age of the CP child and type of CP

Model		Unstandardized		Coefficients <sup>a</sup>		95.0% Confidence Interval for	
		B	Std. Error	Standardized Coefficients Beta	t	Sig.	B Lower Bound Upper Bound
1	(Constant)	2.814	0.102		27.501	0.000	2.612 3.016
	Age of CP child	-0.102	0.048	-0.141	-2.115	0.036	-0.197 -0.007
	Type of CP	-0.059	0.024	-0.163	-2.443	0.015	-0.106 -0.011

a. Dependent Variable: Behaviour