

Table S2. Multivariate Analysis of Risk Factors for Severity in COVID-19 Patients

Independent Variable	aOR	95% CL		Univariate p-value	Global p-value
Gender (REF=Female)	NA**	NA**	NA**	0.9489	0.03489*
Comorbidity (REF=No)	NA**	NA**	NA**	0.8901	
Diabetes (REF=No)	11.6	NA**	NA**	0.9857	
Hypertension (REF=No)	0.082	NA**	NA**	0.9848	
Age	2.018	0.002	NA**	0.8435	
Haemoglobin (g/L)	1.197	0.102	13.990	0.886	
Haematocrit (%)	NA**	NA**	NA**	0.8464	
White Blood Cell ($\times 10^9/L$)	NA**	NA**	NA**	0.8536	
Nutrophils ($\times 10^9/L$)	NA**	NA**	NA**	0.855	
Lymphocytes ($\times 10^9/L$)	NA**	NA**	NA**	0.8372	
Prothrombin Time (seconds)	NA**	NA**	NA**	0.9253	
The International Normalized Ratio	NA**	NA**	NA**	0.9536	
Blood Urea Nitrogen (mmol/L)	0.565	NA**	NA**	0.983	
Lactate Dehydrogenase (U/L)	1.005	0.611	1.652	0.9856	
Creatinine (umol/L)	1.007	0.876	1.15	0.9173	

* Statistically significant value

** Not available, not enough sample size distributed to calculate ORs.